

2. J. W. S. Cassels, "On the Representation of Integers as the Sums of Distinct Summands Taken from a Fixed Set," Szeged, Vol. 21 (1960), pp. 111-124.
3. P. Erdős, "On the Representation of Integers as Sums of Distinct Summands taken from a Fixed Set," Acta Arithmetica, Vol. VII (1962), pp. 345-354.
4. J. Folkman, "On the Representation of Integers as Sums of Distinct Terms from a Fixed Sequence," Can. J. Math., Vol. 18 (1966), pp. 643-655.
5. R. L. Graham, "Complete Sequences of Polynomial Values," Duke Math. Journal, Vol. 31 (1963), pp. 275-285.
6. R. L. Graham, "On a Conjecture of Erdős in Additive Number Theory," Acta Arithmetica, Vol. X (1964), pp. 63-70.
7. C. G. Lekkerkerker, "Representation of Natural Numbers as a Sum of Fibonacci Numbers," Simon Stevin, Vol. 29 (1952), pp. 190-195.
8. N. S. Mendelsohn, "A Linear Diophantine Equation with Applications to Non-negative Matrices," Proc. 1970 Int'l. Conf. on Comb. Math., Annals N. Y. Acad. Sci., 175, No. 1 (1970), pp. 287-294.
9. K. Roth and G. Szekeres, "Some Asymptotic Formulae in the Theory of Partitions," Quart. J. Math., Oxford Ser., Vol. 2, No. 5 (1954), pp. 241-259.



[Continued from page 261.]

A GENERAL Q-MATRIX

2. D. A. Lind, "The Q Matrix as a Counterexample in Group Theory," Fibonacci Quarterly, Vol. 5, No. 1, Feb. 1967, p. 44.
3. E. P. Miles, "Fibonacci Numbers and Associated Matrices," American Mathematical Monthly, Oct. 1960, p. 748.
4. Stephen Smale, "Differentiable Dynamical Systems," Bull. American Math. Society, 73, 1967, pp. 747-817.

