

THE COEFFICIENTS OF A FIBONACCI POWER SERIES

Federico Ardila

Department of Mathematics, Massachusetts Institute of Technology
77 Massachusetts Avenue, Room 2-333, Cambridge, MA 02139
e-mail: fardila@math.mit.edu

(Submitted September 2001-Final Revision February 2002)

Consider the infinite product

$$\begin{aligned} A(x) &= \prod_{k \geq 2} (1 - x^{F_k}) = (1 - x)(1 - x^2)(1 - x^3)(1 - x^5)(1 - x^8) \dots \\ &= 1 - x - x^2 + x^4 + x^7 - x^8 + x^{11} - x^{12} - x^{13} + x^{14} + x^{18} + \dots \end{aligned}$$

regarded as a formal power series. In [4], N. Robbins proved that the coefficients of $A(x)$ are all equal to $-1, 0$ or 1 . We shall give a short proof of this fact, and a very simple recursive description of the coefficients of $A(x)$.

Following the notation of [4], let $a(m)$ be the coefficient of x^m in $A(x)$. It is clear that $a(m) = r_E(m) - r_O(m)$, where $r_E(m)$ is equal to the number of partitions of m into an even number of distinct positive Fibonacci numbers, and $r_O(m)$ is equal to the number of m into an odd number of distinct positive Fibonacci numbers. We call these partitions “even” and “odd” respectively.

Proposition 1: Let $n \geq 5$ be an integer. Consider the coefficients $a(m)$ for m in the interval $[F_n, F_{n+1})$. Split this interval into the three subintervals $[F_n, F_n + F_{n-3} - 2]$, $[F_n + F_{n-3} - 1, F_n + F_{n-2} - 1]$ and $[F_n + F_{n-2}, F_{n+1} - 1]$.

1. The numbers $a(F_n), a(F_n + 1), \dots, a(F_n + F_{n-3} - 2)$ are equal to the numbers $(-1)^{n-1}a(F_{n-3} - 2), (-1)^{n-1}a(F_{n-3} - 3), \dots, (-1)^{n-1}a(0)$ in that order.
2. The numbers $a(F_n + F_{n-3} - 1), a(F_n + F_{n-3}), \dots, a(F_n + F_{n-2} - 1)$ are equal to 0.
3. The numbers $a(F_n + F_{n-2}), a(F_n + F_{n-2} + 1), \dots, a(F_{n+1} - 1)$ are equal to the numbers $a(0), a(1), \dots, a(F_{n-3} - 1)$ in that order.

This description gives a very fast method for computing the coefficients $a(m)$ recursively. Once we have computed them for $0 \leq m < F_n$ we can immediately compute them for $F_n \leq m < F_{n+1}$ using Proposition 1.

Also, since the coefficient of x^m in $A(x)$ is equal to $-1, 0$ or 1 for all non-negative integers $m < F_5$, it follows inductively that the coefficients in each interval $[F_n, F_{n+1})$ are also all equal to $-1, 0$ or 1 . This will prove Robbins’s result.

Proof of Proposition 1: It will be convenient to prove Proposition 1.2 first. Let $F_n + F_{n-3} - 1 \leq m \leq F_n + F_{n-2} - 1$, and consider the partitions of m into distinct positive Fibonacci numbers. It is clear that the largest part in such a partition cannot be F_{n+1} or larger. It cannot be F_{n-2} or smaller either, because $F_{n-2} + F_{n-3} + \dots + F_2 = F_n - 2 < m$. Therefore, it must be F_n or F_{n-1} .

If the largest part is F_n , then the second largest part cannot be F_{n-1} or F_{n-2} . If, on the other hand, it is F_{n-1} , then the second largest part must be F_{n-2} , because $F_{n-1} + F_{n-3} + F_{n-4} + \dots + F_2 = 2F_{n-1} - 2 = F_n + F_{n-3} - 2 < m$.

This means that we can split the set of partitions into pairs. Each pair consists of two partitions of the form $F_n + F_a + F_b + \dots$ and $F_{n-1} + F_{n-2} + F_a + F_b + \dots$, where $n - 3 \geq a > b >$

... In each pair, one of the partitions is even and the other is odd. Therefore $r_E(m) = r_O(m)$ and $a(m) = 0$ as claimed.

Now we use a similar analysis to prove Proposition 1.3. Let $F_n + F_{n-2} \leq m \leq F_{n+1} - 1$. As before, the largest part of a partition of m must be F_n or F_{n-1} . If it is F_n , the second largest part cannot be F_{n-1} . If, on the other hand, it is F_{n-1} , then the second largest part must be F_{n-2} .

Again, we can split a subset of the set of partitions into pairs. Each pair consists of two partitions of the form $F_n + F_a + F_b + \dots$ and $F_{n-1} + F_{n-2} + F_a + F_b + \dots$, where $n - 3 \geq a > b > \dots$. In each pair there is an even and an odd partition.

The remaining partitions are of the form $F_n + F_{n-2} + F_a + F_b + \dots$, where $n - 3 \geq a > b > \dots$. To each one of these partitions we can assign a partition of $m' = m - F_n - F_{n-2}$, by just removing the parts F_n and F_{n-2} . This is in fact a bijection. Since $m' < F_{n-2}$, any partition of m' has largest part less than or equal to F_{n-3} ; therefore it can be obtained in that way from a partition of m .

It is clear that, under this bijection, odd partitions of m go to odd partitions of m' and even partitions of m go to even partitions of m' . It follows that $a(m) = a(m - F_n - F_{n-2})$, as claimed.

Finally we prove Proposition 1.1. Consider $F_n \leq m \leq F_n + F_{n-3} - 2$. The parts of a partition of m come from the list F_2, F_3, \dots, F_n . To each partition π of m , assign the partition π' of $m' = F_{n+2} - 2 - m$ consisting of all the numbers on the above list that do not appear in π . Any partition of m' can be obtained in such a way from a partition of m : the partitions of m' also have all their parts less than or equal to F_n , because it is easily seen that $m' < F_{n+1}$.

So the partitions of m are in bijection with the partitions of m' . If a partition π of m has k parts, the corresponding partition π' of m' has $n - 1 - k$ parts. Therefore, if n is odd, the bijection takes odd partitions to odd partitions and even partitions to even partitions, and $a(m) = a(m')$. If n is even, the bijection takes odd partitions to even partitions, and even partitions to odd partitions, and $a(m) = -a(m')$. In any case, $a(m) = (-1)^{n-1}a(m')$.

Now, it is easily seen that $F_n + F_{n-2} \leq m' \leq F_{n+1} - 2$. Therefore Proposition 1.3 applies, and $a(m') = a(m' - F_n - F_{n-2}) = a(F_n + F_{n-3} - 2 - m)$. Hence $a(m) = (-1)^{n-1}a(F_n + F_{n-3} - 2 - m)$, which is what we wanted to show.

Proposition 2: Given an integer n , pick an integer m uniformly at random from the interval $[0, n]$. Let p_n be the probability that $a(m) = 0$ or, equivalently, that $r_E(m) = r_O(m)$.

Then $\lim_{n \rightarrow \infty} p_n = 1$.

Proof: Let α_n be the number of non-zero coefficients among the first F_n coefficients $a(0), a(1), \dots, a(F_n - 1)$, so that $p_{(F_n-1)} = 1 - \alpha_n/F_n$. Notice that for $F_{n-1} \leq m < F_n$ there are at most α_n non-zero coefficients among $a(0), a(1), \dots, a(m)$, so $p_m \geq 1 - \alpha_n/(m+1) > 1 - 2\alpha_n/F_n$. We shall now prove that $\lim_{n \rightarrow \infty} \alpha_n/F_n = 0$, from which Proposition 2 follows.

First we obtain a recurrence relation for α_n . Consider the non-zero coefficients $a(m)$ for $F_n \leq m \leq F_{n+1} - 1$. We know that there are $\alpha_{n+1} - \alpha_n$ such coefficients. Now split the interval $[F_n, F_{n+1} - 1]$ into the three subintervals $[F_n, F_n + F_{n-3} - 2]$, $[F_n + F_{n-3} - 1, F_n + F_{n-2} - 1]$ and $[F_n + F_{n-2}, F_{n+1} - 1]$. Proposition 1.2 shows that there are no non-zero coefficients in the second subinterval, and Proposition 1.3 shows that there are α_{n-3} non-zero coefficients in the third subinterval. Because $a(F_{n-3} - 1)$ is non-zero for all $n \geq 5$ (this follows inductively from Proposition 1.3), Proposition 1.1 shows that there are $\alpha_{n-3} - 1$ non-zero coefficients in the first subinterval. We conclude that $\alpha_{n+1} - \alpha_n = 2\alpha_{n-3} - 1$.

The characteristic polynomial of this recurrence relation is $x^4 - x^3 - 2 = 0$, and its roots are approximately $r_1 \approx 1.54, r_2 = -1, r_3 \approx 0.23 + 1.12i$ and $r_4 \approx 0.23 - 1.12i$. It follows from standard results on linear recurrences that $\alpha_n = O(r_1^n)$, while $F_n = \Theta(\lambda^n)$, where $\lambda = (\sqrt{5} + 1)/2 \approx 1.62$. Since $r_1 < \lambda$, we conclude that $\lim_{n \rightarrow \infty} \alpha_n / F_n \equiv 0$.

ACKNOWLEDGMENT

The author would like to thank Richard Stanley for encouraging him to work on this problem, and for pointing out [4].

REFERENCES

- [1] L. Carlitz. "Fibonacci Representations." *The Fibonacci Quarterly* **6.4** (1968): 193-220.
- [2] H. H. Ferns. "On the Representations of Integers as Sums of Distinct Fibonacci Numbers." *The Fibonacci Quarterly* **3.1** (1965): 21-30.
- [3] D. Klarner. "Partitions of N into Distinct Fibonacci Numbers." *The Fibonacci Quarterly* **6.4** (1968): 235-243.
- [4] N. Robbins. "Fibonacci Partitions." *The Fibonacci Quarterly* **34.4** (1996): 306-313.

AMS Classification Numbers: 05A15, 05A17, 11B39, 11P81

