

REFERENCES

1. P. F. Byrd, "Expansion of Analytic Functions in Polynomials Associated with Fibonacci Numbers," *The Fibonacci Quarterly*, Vol. 1, No. 1 (1963), pp. 16-28.
2. A. Erdélyi, et al., Higher Transcendental Functions, Vol. 2, McGraw-Hill, New York, 1953.
3. R. G. Buschman, "Fibonacci Numbers, Chebyshev Polynomials, Generalizations and Difference Equations," *The Fibonacci Quarterly*, Vol. 1, No. 4 (1963), pp. 1-7.
4. G. N. Watson, A Treatise on the Theory of Bessel Functions, Cambridge, 2nd Edition, 1944.
5. W. Groebner, and N. Hofreiter, Integraltafel (zweiter Teil, bestimmte Integrale), Springer-Verlag, Vienna, 1950.
6. H. W. Gould, "Generating Functions for Products of Powers of Fibonacci Numbers," *The Fibonacci Quarterly*, Vol. 1, No. 2 (1963), pp. 1-16.

XXXXXXXXXXXXXXXXXXXX

INFORMATION AND AN OMISSION

Reference related to H-37.

References by R. E. Greenwood: Problem 4047, Amer. Math. Mon. (issue of Feb. 1944, pp. 102-104), proposed by T. R. Running, solved by E. P. Starke. Problem #65, Nat. Math. Mag. (now just Math. Mag.) issue of November 1934, p. 63.

Omission H-37. Also solved by J. A. H. Hunter.

CORRECTION

H-28 Let

$$S_n(r, a, b) = \sum_{j=0}^{\infty} C_j(r, n) a^j b^{rn-n-j} = b^{(r-1)n} \sum_{N=0}^{r^{n-1}} \left(\frac{a}{b}\right)^{N_0+N_1+\dots+N_{n-1}}$$