

RECURRENCE SEQUENCES AND NÖRLUND-EULER POLYNOMIALS

Wang Tianming

Institute of Math. Sciences, Dalian University of Technology, Dalian, 116024, China

Zhang Zhizheng

Department of Math., Luoyang Teachers' College, Luoyang, 471022, China

(Submitted October 1994)

1. It is well known that a general linear sequence $S_n(p, q)$ ($n = 0, 1, 2, \dots$) of order 2 is defined by the law of recurrence,

$$S_n(p, q) = pS_{n-1}(p, q) - qS_{n-2}(p, q),$$

with S_0, S_1, p , and q arbitrary, provided that $\Delta = p^2 - 4q > 0$, see [1].

In particular, if $S_0 = 0$ and $S_1 = 1$ or if $S_0 = 2$ and $S_1 = p$, we have generalized Fibonacci and Lucas sequences, respectively, in symbols $U_n(p, q)$ and $V_n(p, q)$.

By the roots $x_1 > x_2$ of the generating equation $x^2 - px + q = 0$, it is proved that

$$U_n(p, q) = \frac{x_1^n - x_2^n}{x_1 - x_2} \quad \text{and} \quad V_n(p, q) = x_1^n + x_2^n; \quad (1)$$

moreover, the general term of the recurrence sequence $S_n(p, q)$ is expressed as a sum of the general terms of generalized Fibonacci and Lucas sequences by the formula

$$S_n(p, q) = \left(S_1 - \frac{1}{2} pS_0 \right) U_n(p, q) + \frac{1}{2} S_0 V_n(p, q). \quad (2)$$

We assume

$$S_0 = \omega,$$

$$S_1 = \frac{1}{2} p\omega + \left(x - \frac{1}{2} \omega \right) \Delta^{\frac{1}{2}},$$

and, according to (1) and (2), we deduce

$$S_n(x; p, q) = \left(x - \frac{1}{2} \omega \right) \Delta^{\frac{1}{2}} \cdot U_n(p, q) + \frac{1}{2} \omega V_n(p, q) \quad (3)$$

and

$$S_n(x; p, q) = x x_1^n + (\omega - x) x_2^n. \quad (4)$$

From this point on, we shall use the brief notation U_n, V_n , and $S_n(x)$ to denote $U_n(p, q), V_n(p, q)$, and $S_n(x; p, q)$, respectively.

2. From (3), we have

$$S_n^m(x) + S_n^m(\omega - x) = \frac{1}{2^{m-1}} \sum_{r=0}^{\left[\frac{m}{2} \right]} \begin{bmatrix} m \\ 2r \end{bmatrix} \Delta^r U_n^{2r} V_n^{m-2r} (2x - \omega)^{2r}, \quad (5)$$

and from (4), we have

$$\begin{aligned} S_n^m(x) + S_n^m(\omega - x) &= \sum_{r=0}^m \begin{bmatrix} m \\ r \end{bmatrix} [x_1^{nr} x_2^{n(m-r)} + x_1^{n(m-r)} x_2^{nr}] x^r (\omega - x)^{m-r} \\ &= \sum_{r=0}^m \begin{bmatrix} m \\ r \end{bmatrix} q^{nr} [x_1^{n(m-2r)} + x_2^{n(m-2r)}] x^r (\omega - x)^{m-r}. \end{aligned}$$

Then we have

$$\begin{aligned} S_n^{2m}(x) + S_n^{2m}(\omega - x) &= \sum_{r=0}^m \begin{bmatrix} 2m \\ r \end{bmatrix} q^{nr} [x_1^{2n(m-r)} + x_2^{2n(m-r)}] x^r (\omega - x)^{2m-r} + \sum_{s=0}^{m-1} \begin{bmatrix} 2m \\ s \end{bmatrix} q^{ns} [x_1^{2n(m-s)} + x_2^{2n(m-s)}] x^{2m-s} (\omega - x)^s \\ &= 2 \begin{bmatrix} 2m \\ m \end{bmatrix} q^{mn} x^m (\omega - x)^m + \sum_{r=0}^{m-1} \begin{bmatrix} 2m \\ r \end{bmatrix} q^{nr} [x_1^{2n(m-r)} + x_2^{2n(m-r)}] [x^r (\omega - x)^{2m-r} + x^{2m-r} (\omega - x)^r] \\ &= 2 \begin{bmatrix} 2m \\ m \end{bmatrix} q^{mn} x^m (\omega - x)^m + \sum_{r=0}^{m-1} \begin{bmatrix} 2m \\ r \end{bmatrix} q^{nr} V_{2n(m-r)} [x^r (\omega - x)^{2m-r} + x^{2m-r} (\omega - x)^r]. \end{aligned} \quad (6)$$

Similarly, we have the analogous formula

$$S_n^{2m+1}(x) + S_n^{2m+1}(\omega - x) = \sum_{r=0}^m \begin{bmatrix} 2m+1 \\ r \end{bmatrix} q^{nr} V_{n(2m-2r+1)} [x^r (\omega - x)^{2m-r+1} + x^{2m-r+1} (\omega - x)^r]. \quad (7)$$

We now have the difference formulas

$$S_n^m(x) - S_n^m(\omega - x) = \frac{\Delta^{\frac{1}{2}}}{2^{m-1}} \sum_{r=0}^{\left[\frac{m-1}{2}\right]} \begin{bmatrix} m \\ 2r+1 \end{bmatrix} \Delta^r U_n^{2r+1} V^{m-2r-1} \omega^{m-2r-1} (2x - \omega)^{2r+1}, \quad (8)$$

and

$$S_n^m(x) - S_n^m(\omega - r) = \Delta^{\frac{1}{2}} \sum_{r=0}^{\left[\frac{m-1}{2}\right]} \begin{bmatrix} m \\ r \end{bmatrix} q^{nr} U_{n(m-2r)} [x^{m-r} (\omega - x)^r - x^r (\omega - x)^{m-r}]. \quad (9)$$

We shall end this section by giving the generating functions

$$\sum_{r=0}^{\infty} \frac{t^r}{r!} U_{nr} = \frac{1}{\Delta^{\frac{1}{2}}} (\exp(tx_1^n) - \exp(tx_2^n)) \quad (10)$$

and

$$\sum_{r=0}^{\infty} \frac{t^r}{r!} V_{nr} = \exp(tx_1^n) + \exp(tx_2^n). \quad (11)$$

3. First, we recall the Nörlund-Euler polynomials $E_n^{(k)}(x|\omega, \dots, \omega_k)$ defined by the generating expansion (see [2], [6]):

$$\sum_{r=0}^{\infty} E_r^{(k)}(x|\omega, \dots, \omega_k) \frac{t^r}{r!} = \frac{2^k e^{xt}}{(e^{\omega_1 t} + 1) \cdots (e^{\omega_k t} + 1)}. \quad (12)$$

In particular, the Nörlund-Euler numbers of order k are given by

$$E_n^{(k)}[\omega_1, \dots, \omega_k] = 2^n E_n^{(k)}\left(\frac{\omega_1 + \dots + \omega_k}{2} \mid \omega_1, \dots, \omega_k\right).$$

If $\omega_1 = \dots = \omega_k = 1$, then $E_n^{(k)}[1, 1, \dots, 1] = E_n^{(k)}$ (the Euler numbers of order k , see [3]), and we note that

$$E_n^{(k)}(\omega_1 + \dots + \omega_k - x \mid \omega_1, \dots, \omega_k) = (-1)^n E_n^{(k)}(x \mid \omega_1, \dots, \omega_k). \quad (13)$$

From (12), replacing t by $\Delta^{\frac{1}{2}} U_n t$, we have

$$\begin{aligned} \sum_{r=0}^{\infty} \frac{(\Delta^{\frac{1}{2}} U_n t)^r}{r!} E_r^{(k)}(x \mid \omega_1, \dots, \omega_k) &= \frac{2^k e^{x \Delta^{\frac{1}{2}} U_n t}}{(e^{\omega_1 \Delta^{\frac{1}{2}} U_n t} + 1) \dots (e^{\omega_k \Delta^{\frac{1}{2}} U_n t} + 1)} \\ &= \frac{2^k}{(e^{\omega_1 t x_1^n} + e^{\omega_1 t x_2^n}) \dots (e^{\omega_k t x_1^n} + e^{\omega_k t x_2^n})} e^{t S_n(x)}, \end{aligned}$$

therefore,

$$(e^{\omega_1 t x_1^n} + e^{\omega_1 t x_2^n}) \dots (e^{\omega_k t x_1^n} + e^{\omega_k t x_2^n}) \sum_{r=0}^{\infty} \frac{(\Delta^{\frac{1}{2}} U_n t)^r}{r!} E_r^{(k)}(x \mid \omega_1, \dots, \omega_k) = 2^k e^{t S_n(x)}.$$

Using (11), we obtain

$$\sum_{r_1=0}^{\infty} \frac{\omega_1^{r_1} t^{r_1}}{r_1!} V_{nr_1} \dots \sum_{r_k=0}^{\infty} \frac{\omega_k^{r_k} t^{r_k}}{r_k!} V_{nr_k} \sum_{r=0}^{\infty} \frac{(\Delta^{\frac{1}{2}} U_n t)^r}{r!} E_r^{(k)}(x \mid \omega_1, \dots, \omega_k) = 2^k e^{t S_n(x)}$$

i.e.,

$$\left[\sum_{r=0}^{\infty} \left(\sum_{r_1+\dots+r_k=r} \frac{\omega_1^{r_1} V_{nr_1}}{r_1!} \dots \frac{\omega_k^{r_k} V_{nr_k}}{r_k!} \right) t^r \right] \sum_{r=0}^{\infty} \frac{(\Delta^{\frac{1}{2}} U_n t)^r}{r!} E_r^{(k)}(x \mid \omega_1, \dots, \omega_k) = 2^k e^{t S_n(x)}.$$

Expanding the product, figuring in the first member, into a power series of t , and comparing with the expansion of the second member, we find

$$\sum_{r=0}^m \left[\begin{matrix} m \\ r \end{matrix} \right] \Delta^{\frac{r}{2}} U_n^r E_r^{(k)}(x \mid \omega_1, \dots, \omega_k) (m-r)! \sum_{r_1+\dots+r_k=m-r} \frac{\omega_1^{r_1} V_{nr_1}}{r_1!} \dots \frac{\omega_k^{r_k} V_{nr_k}}{r_k!} = 2^k S_n^m(x). \quad (14)$$

And if we replace x by $\omega_1 + \dots + \omega_k - x$ in (14), and using (13), we have

$$\begin{aligned} \sum_{r=0}^m \left[\begin{matrix} m \\ r \end{matrix} \right] \Delta^{\frac{r}{2}} U_n^r (-1)^r E_r^{(k)}(x \mid \omega_1, \dots, \omega_k) (m-r)! \sum_{r_1+\dots+r_k=m-r} \frac{\omega_1^{r_1} V_{nr_1}}{r_1!} \dots \frac{\omega_k^{r_k} V_{nr_k}}{r_k!} \\ = 2^k S_n^m(\omega_1 + \dots + \omega_k - x). \end{aligned} \quad (15)$$

Taking (14) + (15), and using $\omega_1 + \dots + \omega_k$ to replace ω in (5), (6), and (7), we obtain

$$\begin{aligned} & \sum_{r=0}^{\left[\frac{m}{2}\right]} \binom{m}{2r} \Delta^r U_n^{2r} E_{2r}^{(k)}(x|\omega_1, \dots, \omega_k) (m-2r)! \sum_{r_1+\dots+r_k=m-2r} \frac{\omega_1^{r_1} V_{nr_1}}{r_1!} \dots \frac{\omega_k^{r_k} V_{nr_k}}{r_k!} \\ &= \frac{1}{2^{m-k}} \sum_{r=0}^{\left[\frac{m}{2}\right]} \binom{m}{2r} \Delta^r U_n^{2r} V_n^{m-2r} (\omega_1 + \dots + \omega_k)^{m-2r} (2x - (\omega_1 + \dots + \omega_k))^{2r} \end{aligned} \quad (16)$$

$$\begin{aligned} &= (1 + (-1)^m) 2^{k-1} \binom{m}{m/2} q^{mn/2} (x(\omega_1 + \dots + \omega_k - x))^{m/2} + 2^{k-1} \sum_{r=0}^{\left[\frac{m-1}{2}\right]} \binom{m}{r} q^{nr} \\ &\cdot V_{n(m-2r)} [x^r (\omega_1 + \dots + \omega_k - x)^{m-r} + x^{m-r} (\omega_1 + \dots + \omega_k - x)^r]. \end{aligned} \quad (17)$$

Taking (14) – (15), and using $\omega_1 + \dots + \omega_k$ to replace ω in (8) and (9), we get

$$\begin{aligned} & \sum_{r=0}^{\left[\frac{m-1}{2}\right]} \binom{m}{2r+1} \Delta^r U_n^{2r+1} E_{2r+1}^{(k)}(x|\omega_1, \dots, \omega_k) (m-2r-1)! \sum_{r_1+\dots+r_k=m-2r-1} \left(\frac{\omega_1^{r_1} V_{nr_1}}{r_1!} \dots \frac{\omega_k^{r_k} V_{nr_k}}{r_k!} \right) \\ &= \frac{1}{2^{m-k}} \sum_{r=0}^{\left[\frac{m-1}{2}\right]} \binom{m}{2r+1} \Delta^r U_n^{2r+1} V_n^{m-2r-1} (\omega_1 + \dots + \omega_k)^{m-2r-1} (2x - (\omega_1 + \dots + \omega_k))^{2r+1} \end{aligned} \quad (18)$$

$$= 2^{k-1} \sum_{r=0}^{\left[\frac{m-1}{2}\right]} \binom{m}{r} q^{nr} U_{n(m-2r)} [x^{m-r} (\omega_1 + \dots + \omega_k - x)^r - x^r (\omega_1 + \dots + \omega_k - x)^{m-r}]. \quad (19)$$

4. If we take $x = \frac{\omega_1 + \dots + \omega_k}{2}$ in (16), then

$$\begin{aligned} & \sum_{r=0}^{\left[\frac{m}{2}\right]} \binom{m}{2r} \Delta^r U_n^{2r} \frac{1}{2^{2r}} E_{2r}^{(k)} [\omega_1, \dots, \omega_k] (m-2r)! \sum_{r_1+\dots+r_k=m-2r} \frac{\omega_1^{r_1} V_{nr_1}}{r_1!} \dots \frac{\omega_k^{r_k} V_{nr_k}}{r_k!} \\ &= \frac{1}{2^{m-k}} (\omega_1 + \dots + \omega_k)^m V_n^m. \end{aligned} \quad (20)$$

Now, setting $\omega_1 = \dots = \omega_k = 1$ in (20), we have

$$\sum_{r=0}^{\left[\frac{m}{2}\right]} \binom{m}{2r} \Delta^r U_n^{2r} \frac{E_{2r}^{(k)}}{2^{2r}} (m-2r)! \sum_{r_1+\dots+r_k=m-2r} \frac{V_{nr_1}}{r_1!} \dots \frac{V_{nr_k}}{r_k!} = \frac{1}{2^{m-k}} k^m V_n^m. \quad (21)$$

Again, if we take $k = 1$, then

$$\sum_{r=0}^{\left[\frac{m}{2}\right]} \binom{m}{2r} \Delta^r U_n^{2r} \frac{E_{2r}}{2^{2r}} V_{n(m-2r)} = \frac{1}{2^{m-1}} V_n^m. \quad (22)$$

If we set $k = 1$ in (18), we obtain

$$\begin{aligned} & \sum_{r=0}^{\left[\frac{m-1}{2}\right]} \binom{m}{2r+1} \Delta^r U_n^{2r+1}(x|\omega_1) V_{n(m-2r-1)} \omega_1^{m-2r-1} \\ &= \frac{1}{2^{m-1}} \sum_{r=0}^{\left[\frac{m-1}{2}\right]} \binom{m}{2r+1} \Delta^r U_n^{2r+1} V_n^{m-2r-1} \omega_1^{m-2r-1} (2x - \omega_1)^{2r+1}. \end{aligned} \quad (23)$$

Now, taking $\omega_1 = 1$ and $x = 0$ or $x = \frac{1}{3}$, and using the following relations (see [1]),

$$E_{n-1}(0) = 2(1 - 2^n) \frac{B_n}{n},$$

$$E_{2n-1}\left(\frac{1}{3}\right) = (2^{2n} - 1) \left(\frac{1}{3^{2n} - 1}\right) \frac{B_{2n}}{2n},$$

where B_n is a Bernoulli number, we have

$$\begin{aligned} & \sum_{r=0}^{\left[\frac{m-1}{2}\right]} \binom{m}{2r+1} \Delta^r U_n^{2r+1} \frac{1}{r+1} (2^{2r+2} - 1) B_{2r+2} V_{n(m-2r-1)} \\ &= \frac{1}{2^{m-1}} \sum_{r=0}^{\left[\frac{m-1}{2}\right]} \binom{m}{2r+1} \Delta^r U_n^{2r+1} V_n^{m-2r-1}, \end{aligned} \quad (24)$$

and

$$\begin{aligned} & \sum_{r=0}^{\left[\frac{m-1}{2}\right]} \binom{m}{2r+1} \Delta^r U_n^{2r+1} (2^{2r+2} - 1) \left(1 - \frac{1}{3^{2r+1}}\right) \frac{B_{2r+2}}{2r+2} V_{n(m-2r-1)} \\ &= \frac{1}{2^{m-1}} \sum_{r=0}^{\left[\frac{m-1}{2}\right]} \binom{m}{2r+1} \Delta^r U_n^{2r+1} V_n^{m-2r-1} \frac{1}{3^{2r+1}}. \end{aligned} \quad (25)$$

Assuming $p = 1$ and $q = -1$, we have the so-called Fibonacci and Lucas sequences

$$U_n = F_n \quad \text{and} \quad V_n = L_n,$$

respectively. And from (22), (24), and (25), it follows that

$$\sum_{r=0}^{\left[\frac{m-1}{2}\right]} \binom{m}{2r} 5^r F_n^{2r} \frac{E_{2r}}{2^{2r}} L_{n(m-2r)} = \frac{1}{2^{m-1}} L_n^m, \quad (26)$$

$$\begin{aligned} & \sum_{r=0}^{\left[\frac{m-1}{2}\right]} \binom{m}{2r+1} 5^r F_n^{2r+1} \frac{1}{r+1} (2^{2r+2} - 1) B_{2r+2} L_{n(m-2r-1)} \\ &= \frac{1}{2^{m-1}} \sum_{r=0}^{\left[\frac{m-1}{2}\right]} \binom{m}{2r+1} 5^r F_n^{2r+1} L_n^{m-2r-1}, \end{aligned} \quad (27)$$

and

$$\begin{aligned}
& \sum_{r=0}^{\left[\frac{m-1}{2}\right]} \binom{m}{2r+1} 5^r F_n^{2r+1} (2^{2r+2} - 1) \left(1 - \frac{1}{3^{2r+1}}\right) \frac{B_{2r+2}}{2r+2} L_{n(m-2r-1)} \\
& = \frac{1}{2^{m-1}} \sum_{r=0}^{\left[\frac{m}{2}\right]} \binom{m}{2r} 5^r F_n^{2r+1} L_n^{(m-2r-1)} \frac{1}{3^{2r+1}}, \tag{28}
\end{aligned}$$

where (26) is a generalization of P. F. Byrd's result (see [5]):

$$\sum_{r=0}^{\left[\frac{m}{2}\right]} 5^r \binom{m}{2r} B_{2r} F_n^{2r} F_{n(m-2r)} = \frac{m}{2} F_n L_{n(m-1)}.$$

ACKNOWLEDGMENT

The authors wish to thank the referees for their patience and for suggestions that significantly improved the appearance of this paper.

REFERENCES

1. L. Toscano. "Recurring Sequences and Bernoulli-Euler Polynomials." *J. Comb. Inform. & System Sci.* **4** (1979):303-08.
2. L. Carlitz. "Eulerian Numbers and Polynomials of Higher Order." *Duke Math. J.* **27** (1960): 401-23.
3. L. C. Hsu. "Finding Some Strange Identities via Fa  di Bruno's Formula." *J. Math. Research & Exposition* **13** (1993):159-65.
4. R. P. Kelisky. "On Formulas Involving Both the Bernoulli and Fibonacci Numbers." *Scripta Math.* **23** (1957):27-35.
5. P. F. Byrd. "Relations between Euler and Lucas Numbers." *The Fibonacci Quarterly* **13.2** (1975):111-14.
6. N. E. N rlund. *Vorlesungen ober differenzenrechnung*. Berlin, 1924.
7. Zhang Zhizheng. "Recurrence Sequences and Bernoulli Polynomials of Higher Order." *The Fibonacci Quarterly* **33.4** (1995):359-62.

AMS Classification Numbers: 11B39, 11B37

