

REFERENCES

1. G. Djordjević. "On Some Properties of Generalized Hermite Polynomials." *The Fibonacci Quarterly* **34.1** (1996):2-6.
2. G. Djordjević, & G. V. Milovanović. "A Generalization of One Class of Panda's Polynomials." In *IV Conference on Applied Mathematics*, pp. 42-47. Belgrade, 1989.
3. R. Panda. "On a New Class of Polynomials." *Glasgov Math. J.* **18** (1977):105-08.
4. R. C. Singh Chandel. "A Note on Some Generating Functions." *Indiana J. Math.* **25** (1983): 185-88.
5. H. M. Srivastava. "A Note on a Generating Function for the Generalized Hermite Polynomials." *Nederl. Akad. Wetensch. Proc. Ser A*, **79** (1976):457-61.
6. H. M. Srivastava. "On the Product of Two Laguerre Polynomials." *Rollettino della Unione Matematica Italiano* **5.4** (1972):1-6.

AMS Classification Numbers: 11B39, 11B83



**Errata for "Generalizations of Some Identities Involving the Fibonacci Numbers"**

**by Fengzhen Zhao & Tianming Wang**

*The Fibonacci Quarterly* **39.2** (2001):165-167

On page 166, (10) should be

$$\sum_{a+b+c=n} U_{ak}U_{bk}U_{ck} = \frac{U_k^2}{2(V_k^2 - 4q^k)^2} ((n-1)(n-2)V_k^2 U_{nk} - q^k V_k (4n^2 - 6n - 4)U_{(n-1)k} + q^{2k} (4n^2 - 4)U_{(n-2)k}), \quad n \geq 2.$$

Hence, on page 167, (13) should be

$$\sum_{a+b+c=n} F_{ak}F_{bk}F_{ck} = \frac{F_k^2}{2(L_k^2 - 4(-1)^k)^2} ((n-1)(n-2)L_k^2 F_{nk} - (-1)^k L_k (4n^2 - 6n - 4)F_{(n-1)k} + (4n^2 - 4)F_{(n-2)k}), \quad n \geq 2.$$

In the meantime, line 14 and line 16 of page should be, respectively,

$$\sum_{a+b+c=n} F_{2a}F_{2b}F_{2c} = \frac{1}{50} (9(n-1)(n-2)F_{2n} - 3(4n^2 - 6n - 4)F_{2n-2} + (4n^2 - 4)F_{2n-4}).$$

$$\sum_{a+b+c=n} F_{2a}F_{2b}F_{2c} = \frac{1}{50} ((15n^2 - 63n + 66)F_{2n-3} + (10n^2 - 36n + 44)F_{2n-4}).$$

Line 19 of page 167 should be:  $+(4n^2 - 4)P_{(n-2)k}), \quad n \geq 2.$