

# TSCHEBYSCHIEFF AND OTHER FUNCTIONS ASSOCIATED WITH THE SEQUENCE $\{w_n(a,b; p,q)\}$

A. F. HORADAM  
University of New England, Armidale, Australia,  
and University of Leeds, England

## 1. INTRODUCTION

Previously in this journal [5] and [6], I have defined a generalized sequence  $\{w_n(a,b; p,q)\}$  and established its fundamental general arithmetical properties, as well as certain special properties of it. In this article, the sequence is related to Tschebyscheff functions and to some combinatorial functions used by Riordan [8]. This is the third of a series of articles developing the theory of  $\{w_n(a,b; p,q)\}$ , as envisaged in [5]. Notation and content of [5] and [6] are assumed when the occasion warrants.

For subsequent reference, we reproduce the Lucas results [7]

$$(1.1) \quad u_n(p,q) = \sum_{k=0}^{\lfloor n/2 \rfloor} (-1)^k \binom{n-k}{p} p^{n-2k} q^k$$

and

$$(1.2) \quad v_n(p,q) = \sum_{k=0}^{\lfloor n/2 \rfloor} (-1)^k \frac{n}{n-k} \binom{n-k}{k} p^{n-2k} q^k$$

with reciprocals [3]

$$(1.3) \quad p_n = \sum_{k=0}^{\lfloor n/2 \rfloor} \left[ \binom{n}{k} - \binom{n}{k-1} \right] u_{n-2k}(p,q) q^k$$

and

$$(1.4) \quad p^n = \sum_{k=0}^{\lfloor n/2 \rfloor} \binom{n}{k} v_{n-2k}(p, q) q^k \quad (v_0(p, q) = 1) ,$$

respectively. Consequently, it follows that  $(p = -q = 1)$ .

$$(1.5) \quad f_n = \sum_{k=0}^{\lfloor n/2 \rfloor} \binom{n-k}{k}$$

from (1.1), and

$$(1.6) \quad l_n = \sum_{k=0}^{\lfloor n/2 \rfloor} \frac{n}{n-k} \binom{n-k}{k}$$

from (1.2), with appropriate reciprocals from (1.3) and (1.4).

Making use of (1.1) above together with the first of the forms given in (2.14) [5], we may express  $w_n$  as

$$(1.7) \quad w_n(a, b, p, q) = a \sum_{k=0}^{\lfloor \frac{n}{2} \rfloor} (-1)^k \binom{n-k}{k} p^{n-2k} q^k + (b - pa) \sum_{k=0}^{\lfloor \frac{n-1}{2} \rfloor} (-1)^k \binom{n-1-k}{k} p^{n-1-2k} q^k$$

## 2. TSCHEBYSCHIEFF FUNCTIONS

Write

$$(2.1) \quad x = \cos \theta$$

$$(2.2) \quad p = 2x, \quad q = 1$$

so that

$$(2.3) \quad d = 2i \sin \theta \quad (i = \sqrt{-1}) .$$

Define

$$(2.4) \quad w_n = w_n(a, 2x; 2x, 1) = a \cos n\theta + (2 - a) \sin n\theta \cot \theta .$$

Using Simpson's formulae (reference Lucas [7]),

$$(2.5) \quad \begin{cases} \sin (n+2)\theta = 2 \cos \theta \sin (n+1)\theta - \sin n\theta \\ \cos (n+2)\theta = 2 \cos \theta \cos (n+1)\theta - \cos n\theta \end{cases} ,$$

we deduce that

$$(2.6) \quad w_{n+2} = pw_{n+1} - w_n ,$$

as required by the definition of  $w_n(a, b; p, q)$  given in [5], in conjunction with (2.1) and (2.2). Notice that (2.1) and (2.2) ensure [5] that

$$(2.7) \quad e = 4(a - 1) \cos^2 \theta - a^2 ,$$

whence, for  $\{u_n\}$ , for which  $a = 1$ ,

$$(2.8) \quad e = -1 ,$$

while for  $\{v_n\}$ , for which  $a = 2$ ,

$$(2.9) \quad e = -4 \sin^2 \theta .$$

Immediately from (2.4) we have the Lucas substitutions [7]

$$(2.10) \quad u_n(2x, 1) = \frac{\sin (n+1)\theta}{\sin \theta} = \sum_{k=0}^{\lfloor n/2 \rfloor} (-1)^k \binom{n-k}{k} (2x)^{n-2k}$$

and

$$(2.11) \quad v_n(2x, 1) = 2 \cos n\theta = \sum_{k=0}^{\lfloor n/2 \rfloor} (-1)^k \frac{n}{n-k} \binom{n-k}{k} (2x)^{n-2k}$$

with reciprocals

$$(2.12) \quad (2x)^n = \sum_{k=0}^{\lfloor n/2 \rfloor} \left[ \binom{n}{k} - \binom{n}{k-1} \right] u_{n-2k}(2x, 1)$$

and

$$(2.13) \quad (2x)^n = \sum_{k=0}^{\lfloor n/2 \rfloor} \binom{n}{k} v_{n-2k}(2x, 1) ,$$

where we have used (1.1)-(1.4).

But, the expressions in (2.10) and (2.11) exactly describe the Tschebyscheff functions  $U_n(x)$  and  $2T_n(x) = t_n(x)$  respectively ( $T_0 = \frac{1}{2}t_0 = 1$ ). That is,

$$(2.14) \quad w_n(1, 2x; 2x, 1) = u_n(2x, 1) = U_n(x) = 2xU_{n-1}(x) - U_{n-1}(x)$$

and

$$(2.15) \quad w_n(2, 2x; 2x, 1) = v_n(2x, 1) = 2T_n(x) = 2(xU_{n-1}(x) - U_{n-2}(x)) .$$

Special cases are

$$(2.16) \quad w_n(1, 1; 1, 1) = u_n(1, 1) = U_n\left(\frac{1}{2}\right) = U_{n-1}\left(\frac{1}{2}\right) - U_{n-2}\left(\frac{1}{2}\right)$$

and

$$(2.17) \quad w_n(2, 1; 1, 1) = v_n(1, 1) = 2T_n\left(\frac{1}{2}\right) = U_{n-1}\left(\frac{1}{2}\right) - 2U_{n-2}\left(\frac{1}{2}\right) .$$

Generally,

$$(2.18) \quad w_n(a, b; 2x, 1) = bU_{n-1}(x) - aU_{n-2}(x) .$$

By means of the  $w_n$ -notation, relationships among Tschebyscheff polynomials may be conveniently expressed. Recalling the known result [8], for instance, that

$$(2.19) \quad T_n(x) = 2xT_{n-1}(x) - T_{n-2}(x)$$

we may, writing for brevity,

$$(2.20) \quad w_n = w_n(2, 2x; 2x, 1) ,$$

express it in the form

$$(2.21) \quad \omega_n = 2x\omega_{n-1} - \omega_{n-2} .$$

Equations (2.4), (2.10) and (2.11) enable us to express every formula in the theory of our second-order recurrences as a corresponding formula involving trigonometrical functions. [Observe that  $q = 1$  invalidates any specialized application to the sequences  $\{h_n\}$ ,  $\{f_n\}$  and  $\{l_n\}$ , for all of which  $q = -1$ .]

Corresponding to the fundamental formula  $w_{n+r}w_{n-r} - w_n^2 = eq^{n-r}u_{r-1}^2$  ((4.5) in [5]), for instance, we have

$$(2.22) \quad a^2 \{ \cos(n+r)\theta \cos(n-r)\theta - \cos^2 n\theta \} \\ + (2-a)^2 \cot^2 \theta \{ \sin(n+r)\theta \sin(n-r)\theta - \sin^2 n\theta \} = e \frac{\sin^2 r\theta}{\sin^2 \theta}$$

where  $e$  is given by (2.7). For  $\{u_n\}$  and  $\{v_n\}$ , we obtain

$$(2.23) \quad \sin(n+r+1)\theta \sin(n-r+1)\theta - \sin^2(n+1)\theta = -\sin^2 r\theta$$

and

$$(2.24) \quad \cos(n+r)\theta \cos(n-r)\theta - \cos^2 n\theta = -\sin^2 r\theta,$$

in which  $e$  is given by (2.8) and (2.9), respectively. Both results (2.23) and (2.24) are easy to verify. The particular result  $w_n^2 + eu_{n-1}^2 = aw_{2n}$  ((4.6) [5]) derived by setting  $r = m$  implies the identity

$$\cos 2n\theta - \cos^2 n\theta = -\sin^2 n\theta$$

in (2.24).

Other trigonometrical identities are not hard to detect, but it is interesting to discover just how they are disguised. As further examples, we note that

$$pw_{n+2} - (p^2 - q)w_{n+1} + q^2w_{n-1} = 0$$

((3.3) [5]), and

$$\frac{w_{n+r} + q^r w_{n-r}}{w_n} = v_r$$

((3.16) [5]) lead to, respectively,

$$(2.25) \quad \left\{ \begin{array}{l} 2 \cos \theta \sin(n+3)\theta - (4 \cos^2 \theta - 1) \sin(n+2)\theta + \sin n\theta = 0 \\ 2 \cos \theta \cos(n+2)\theta - (4 \cos^2 \theta - 1) \cos(n+1)\theta + \cos(n-1)\theta = 0 \end{array} \right.$$

$$(2.26) \quad \left\{ \begin{array}{l} 2 \cos \theta \sin(n+3)\theta - (4 \cos^2 \theta - 1) \sin(n+2)\theta + \sin n\theta = 0 \\ 2 \cos \theta \cos(n+2)\theta - (4 \cos^2 \theta - 1) \cos(n+1)\theta + \cos(n-1)\theta = 0 \end{array} \right.$$

and

$$(2.27) \quad \left\{ \begin{array}{l} \frac{\sin(n+r+1)\theta + \sin(n-r+1)\theta}{\sin(n+1)\theta} = 2 \cos r\theta \\ \frac{\cos(n+r)\theta + \cos(n-r)\theta}{\cos n\theta} = 2 \cos r\theta \end{array} \right.$$

$$(2.28) \quad \left\{ \begin{array}{l} \frac{\sin(n+r+1)\theta + \sin(n-r+1)\theta}{\sin(n+1)\theta} = 2 \cos r\theta \\ \frac{\cos(n+r)\theta + \cos(n-r)\theta}{\cos n\theta} = 2 \cos r\theta \end{array} \right.$$

where, in each pair of identities, the first refers to  $\{u_n\}$  and the second to  $\{v_n\}$ . A formula also worth investigation is

$$aw_{m+n} + (b - pa)w_{m+n-1} = w_m w_n - qw_{m-1} w_{n-1}$$

((4.1) [5]). Furthermore, the summation formula (3.4) [5] indicates expressions for

$$\sum_{k=0}^{n-1} \cos k\theta$$

and

$$\sum_{k=0}^{n-1} \sin (k + 1)\theta$$

Similar remarks apply to the formulae for sums of squares and cubes.

Instead of (2.1)-(2.3), we may put

$$(2.29) \quad y = \cosh \phi = \cos i\phi$$

$$(2.30) \quad p = 2y, \quad q = 1$$

so that

$$(2.31) \quad d = 2 \sinh \phi = -2i \sin i\phi$$

and hence derive a set of parallel results for hyperbolic functions.

Apart from the Carlitz [3] reference quoted earlier, other sources of information regarding the relationships among Tschebyscheff polynomials and Fibonacci-type sequences are, say, Buschman [1] and Gould [4].

### 3. COMBINATORIAL FUNCTIONS

From (1.1), we have, using the combinatorial function  $L_n(x)$  used in Riordan [8],

$$(3.1) \quad u_n(1, -x) = \sum_{k=0}^{\lfloor n/2 \rfloor} \binom{n-k}{k} x^k = L_{n-1}(x).$$

Then, by the second half of the expression (2.14) [5],

$$(3.2) \quad \begin{aligned} w_n(a, b; 1, -x) &= b \sum_{k=0}^{\lfloor \frac{n-1}{2} \rfloor} \binom{n-1-k}{k} x^k + ax \sum_{k=0}^{\lfloor \frac{n-2}{2} \rfloor} \binom{n-2-k}{k} x^k \\ &= bL_{n-2}(x) + axL_{n-3}(x) \\ &= b \left\{ \frac{(1+g)^n - (1-g)^n}{2^n g^2} \right\} + ax \left\{ \frac{(1+g)^{n-1} - (1-g)^{n-1}}{2^{n-1} g^2} \right\} \end{aligned}$$

where, for brevity,  $g = (1+4x)^{\frac{1}{2}}$ .

More particularly, notice that

$$(3.3) \quad w_n(1, 1; 1, -x) = u_n(1, -x)$$

affords an alternative expression for the known recurrence relation [8].

$$(3.4) \quad L_{n-1}(x) = L_{n-2}(x) + xL_{n-3}(x) \quad [L_0 = 1, L_1 = 1+x]$$

while

$$(3.5) \quad w_{2n}(2, 1; 1, -x) = v_{2n}(1, -x)$$

is an alternative expression for the combinatorial function [8]

$$(3.6) \quad M_n(x) = L_{2n-1}(x) + xL_{2n-3}(x) \quad (n > 1).$$

Of course,

$$(3.7) \quad L_{n-1}(1) = f_n$$

$$(3.8) \quad M_n(1) = I_{2n}.$$

#### 4. OTHER FUNCTIONS

Besides these combinatorial functions and Tschebyscheff functions (themselves involving trigonometrical and hyperbolic functions), other functions are related to the Fibonacci-type recurrences. In this respect, a recent article by Byrd [2] is worth emphasizing, particularly as, it seems, his work offers possibilities for generalization. In this article, Byrd considers the expansion of analytical functions in a certain set of polynomials which can be associated with Fibonacci numbers. Bessel functions and modified Bessel functions are involved in the process.

Throughout, we have assumed that  $p^2 \neq 4q$ . The degenerate case  $p^2 = 4q$  has been discussed by Carlitz [3], who relates it to the Eulerian polynomial, and, briefly, by the author [5].

#### REFERENCES

1. R. Buschman, "Fibonacci Numbers, Chebychev Polynomials, Generalizations and Difference Equations," The Fibonacci Quarterly, Vol. 1, No. 4, 1963, pp. 1-7.
2. P. Byrd, "Expansion of Analytic Functions in Polynomials Associated with Fibonacci Numbers," The Fibonacci Quarterly, Vol. 1, No. 1, 1963, p. 16.
3. L. Carlitz, "Generating Functions for Powers of Certain Sequences of Numbers," Duke Math. J., Vol. 29, No. 4, 1962, pp. 521-538.
4. H. Gould, "A New Series Transform with Applications to Bessel, Legendre and Tschebyscheff Polynomials," Duke Math. J., Vol. 31, No. 2, 1964, pp. 325-334.
5. A. Horadam, "Basic Properties of a Certain Generalized Sequence of Numbers," The Fibonacci Quarterly, Vol. 3, No. 3, p. 161.
6. A. Horadam, "Special Properties of the Sequence  $\{w_n(a, b; p, q)\}$ ," The Fibonacci Quarterly, Vol. 5, No. 5, pp. 424-434.
7. E. Lucas, Théorie des Nombres, Paris, 1961, Chapter 18.
8. J. Riordan, An Introduction to Combinatorial Analysis, New York, 1958.

\*\*\*\*\*