

The Sequent Calculus of Skew Monoidal Categories

Tarmo Uustalu¹

*Reykjavik University, Reykjavik, Iceland
Tallinn University of Technology, Tallinn, Estonia*

Niccolò Veltri²

IT University of Copenhagen, Copenhagen, Denmark

Noam Zeilberger³

University of Birmingham, Birmingham, United Kingdom

Abstract

Szlachányi’s skew monoidal categories are a well-motivated variation of monoidal categories in which the unitors and associator are not required to be natural isomorphisms, but merely natural transformations in a particular direction. We present a sequent calculus for skew monoidal categories, building on the recent formulation by one of the authors of a sequent calculus for the Tamari order (skew semigroup categories). In this calculus, antecedents consist of a stoup (an optional formula) followed by a context (a list of formulae), and the connectives tensor and unit behave like in non-commutative multiplicative linear logic (the logic of monoidal categories) except that the left rules may only be applied in stoup position. We show the admissibility of two forms of the cut rule (stoup cut and context cut), and prove the calculus sound and complete with respect to existence of maps in the free skew monoidal category. We then introduce an equivalence relation on sequent calculus derivations and prove that there is a one-to-one correspondence between equivalence classes of derivations and maps in the free skew monoidal category. Finally, we identify a subcalculus of focused derivations, and establish that it contains exactly one canonical representative from each equivalence class. As an end result, we obtain simple algorithms both for deciding equality of maps in the free skew monoidal category and for enumerating any homset without duplicates, in particular, for deciding whether there is a map. We have formalized this development in the dependently typed programming language Agda.

Keywords: skew monoidal categories, substructural logics, sequent calculus, nonstandard sequent forms, cut admissibility, focusing, Agda

1 Introduction

Skew monoidal categories of Szlachányi [16] are a variation of monoidal categories [13,3]. In a skew monoidal category, the unitors and associator are not required to be natural isomorphisms, but only natural transformations in a particular direction. Szlachányi’s original motivation was from quantum structures. In a different context, the first author of this paper ran into skew monoidal categories studying a generalization of monads to functors between different categories—relative monads [1].

Skew monoidal categories are a fascinating structure, very simply defined but with remarkably subtle properties. Szlachányi’s paper was immediately noticed by Street, Lack and colleagues who have by now published a whole series of papers about them [9,10,6,4,5, ...]. Mac Lane’s original coherence theorem for

¹ Email: tarmo@ru.is

² Email: nive@itu.dk

³ Email: n.zeilberger@cs.bham.ac.uk

monoidal categories is often summarized as “all diagrams commute”, but this is no longer true in the skew monoidal case: it is not the case that there is at most one map between any two objects in the free skew monoidal category on a set of generators (not even for one generator); also it is not easy to give a simple necessary and sufficient condition for the existence of such a map. Curiously, there *is* at most one map between any two objects in the free skew *semigroup* category, so multiple maps originate from the presence of the unit. Uustalu [17] showed that there is at most one map between an object and an object in normal form, and exactly one map between an object and that object’s normal form. Lack and Street [10] gave a concrete description of the free skew monoidal category on one generating object \mathbf{Fsk} , and proved that it admits a faithful structure-preserving functor to Δ_{\perp} , the (strictly associative) skew-monoidal category of finite non-empty ordinals and first-element-and-order-preserving functions. This analysis was further elaborated by Bourke and Lack [4] with a more explicit description of the morphisms of \mathbf{Fsk} , although they still took as given the classical *Tamari order* [15], that is, the partial order on fully-bracketed words induced by a non-invertible associative law (or what can be equivalently seen as the free skew semigroup category on one generator).

In this paper, we present a sequent calculus for skew monoidal categories, building on a recent proof-theoretic analysis of the Tamari order by the third author [18]. There, it was observed that the Tamari order is precisely captured by a sequent calculus very similar to Lambek’s original calculus [11] for (what is nowadays referred to as) non-commutative intuitionistic linear logic, but with tensor as the only logical connective, and with the tensor left rule restricted to only apply to the first formula in the antecedent. The sequent calculus of [18] admits a strong form of cut elimination known as *focusing completeness* (after Andreoli [2]), with the consequence that valid entailments in the Tamari order are in one-to-one correspondence with focused sequent calculus derivations.

The situation becomes significantly more subtle with the addition of a unit. As we will explain, sequents now need to have an explicit “stoup”, corresponding to a distinguished first position in the antecedent which can either be empty or contain a formula. In particular, the left rules for unit and tensor can only be applied to a formula inside the stoup. We will develop the metatheory of this sequent calculus, and see that the presence of the stoup is crucial for adequacy with respect to skew monoidal categories.

After establishing the admissibility of two cut rules (stoup cut and context cut), we prove that the sequent calculus is sound and complete with respect to the free skew monoidal category in the sense that morphisms can be mapped to derivations and vice versa. We then impose a certain notion of equivalence on sequent calculus derivations in order to prove that these two mappings are inverses of each other (i.e., that derivations in the sequent calculus and maps in the free skew monoidal category are in bijection). Finally, we identify the subcalculus of focused derivations, and show that every equivalence class of derivations contains exactly one focused derivation. This means that the focused sequent calculus characterizes the free skew monoidal category in a particularly appealing fashion. Moreover, as the focused representation of a derivation can be easily computed, we get a simple algorithm for deciding equality of morphisms in the free skew monoidal category: two morphisms are equal if and only if they correspond to the same focused derivation. Also, since focused derivations can be systematically searched for, we get an algorithm for enumerating (without duplications) any homset, in particular, for deciding whether a homset is inhabited.

Bourke and Lack [5] have recently related skew monoidal categories to what they call skew multicategories and, in particular, established a 2-equivalence between skew monoidal categories and left representable skew multicategories. In terms of our sequent calculus, their “tight maps” (resp. “loose maps”) correspond to derivations of sequents where the stoup is non-empty (resp. empty). Likewise, their notion of “left” representability (which refines the more standard notion of representability for multicategories [12, Ch. 3]) is precisely analogous to the stoup restriction on left rules. A proper analysis of our sequent calculus from this perspective is left to future work, although we include some additional remarks about how the sequent calculus defines a generalized multicategory for a particular Cartesian monad with some superstructure.

The paper is organized as follows. In Section 2, we introduce skew monoidal categories and the free skew monoidal category as a deductive calculus. In Section 3, we introduce our sequent calculus, show that it admits the appropriate cut rules and satisfies a number of equations reminiscent of the equations of a multicategory. In Section 4, we present our soundness and completeness proofs of the sequent calculus wrt. the categorical calculus, introduce our notion of equality of derivations for the sequent calculus and prove that the derivations in the two calculi are in a bijection. We develop focusing in Section 5. In Section 6, we conclude and outline some avenues of future work. In Appendix A, we show that the sequent calculus is a generalized multicategory. In Appendix B, we discuss a partial order one can introduce on the categorical calculus derivations and the corresponding partial order on the sequent calculus derivations.

We have formalized the development of Sections 2-5 in the dependently typed programming language Agda.

Our Agda formalization is available at <http://cs.ioc.ee/~niccolo/skewmonseqcalc/>. The formalization uses Agda 2.5.3 and Agda standard library 0.14.

2 Skew monoidal categories

A *skew monoidal category* [16] is a category \mathcal{C} together with a distinguished object $\mathbb{1}$, a functor $\otimes : \mathcal{C} \times \mathcal{C} \rightarrow \mathcal{C}$ and three natural transformations

$$\lambda_A : \mathbb{1} \otimes A \rightarrow A \quad \rho_A : A \rightarrow A \otimes \mathbb{1} \quad \alpha_{A,B,C} : (A \otimes B) \otimes C \rightarrow A \otimes (B \otimes C)$$

satisfying the laws

$$\begin{array}{l} \text{(a)} \quad \begin{array}{c} \mathbb{1} \otimes \mathbb{1} \\ \rho_{\mathbb{1}} \swarrow \quad \searrow \lambda_{\mathbb{1}} \\ \mathbb{1} \end{array} \\ \text{(b)} \quad \begin{array}{ccc} (A \otimes \mathbb{1}) \otimes B & \xrightarrow{\alpha_{A,\mathbb{1},B}} & A \otimes (\mathbb{1} \otimes B) \\ \rho_{A \otimes B} \uparrow & & \downarrow A \otimes \lambda_B \\ A \otimes B & \xlongequal{\quad} & A \otimes B \end{array} \\ \text{(c)} \quad \begin{array}{ccc} (\mathbb{1} \otimes A) \otimes B & \xrightarrow{\alpha_{\mathbb{1},A,B}} & \mathbb{1} \otimes (A \otimes B) \\ \lambda_{A \otimes B} \searrow & & \swarrow \lambda_{A \otimes B} \\ & A \otimes B & \end{array} \\ \text{(d)} \quad \begin{array}{ccc} (A \otimes B) \otimes \mathbb{1} & \xrightarrow{\alpha_{A,B,\mathbb{1}}} & A \otimes (B \otimes \mathbb{1}) \\ \rho_{A \otimes B} \swarrow & & \searrow A \otimes \rho_B \\ & A \otimes B & \end{array} \\ \text{(e)} \quad \begin{array}{ccc} (A \otimes (B \otimes C)) \otimes D & \xrightarrow{\alpha_{A,B \otimes C,D}} & A \otimes ((B \otimes C) \otimes D) \\ \alpha_{A,B,C \otimes D} \uparrow & & \downarrow A \otimes \alpha_{B,C,D} \\ ((A \otimes B) \otimes C) \otimes D & \xrightarrow{\alpha_{A \otimes B,C,D}} (A \otimes B) \otimes (C \otimes D) & \xrightarrow{\alpha_{A,B,C \otimes D}} A \otimes (B \otimes (C \otimes D)) \end{array} \end{array}$$

Notice that (a)–(e) are directed versions of the original Mac Lane axioms [13]. Later, Kelly [8] observed that, when λ , ρ and α are natural isomorphisms, laws (a), (c), and (d) can be derived from (b) and (e). For skew monoidal categories, this is not the case.

Skew monoidal categories arise more often than one would perhaps first think, see [16,9,6,17]. The following are some examples from [17].

Example 2.1 A simple example of a skew monoidal category results from skewing a numerical addition monoid.

View the partial order (\mathbb{N}, \leq) of natural numbers as a thin category. Fix some natural number n and define $\mathbb{1} = n$ and $x \otimes y = (x \dot{-} n) + y$ where $\dot{-}$ is “truncating subtraction”. We have $\lambda_x : (n \dot{-} n) + x = 0 + x = x$, $\rho_x : x \leq x \max n = (x \dot{-} n) + n$, $\alpha_{x,y,z} : ((x \dot{-} n) + y) \dot{-} n + z \leq (x \dot{-} n) + (y \dot{-} n) + z$ (by a small case analysis).

Example 2.2 The category **Ptd** of pointed sets and point-preserving functions has the following skew monoidal structure.

Take $\mathbb{1} = (1, *)$ and $(X, p) \otimes (Y, q) = (X + Y, \text{inl } p)$ (notice the “skew” in choosing the point). We define $\lambda_X : (1, *) \otimes (X, p) = (1 + X, \text{inl } *) \rightarrow (X, p)$ by $\lambda_X(\text{inl } *) = p$, $\lambda_X(\text{inr } x) = x$ (this is not injective). We let $\rho_X : (X, p) \rightarrow (X + 1, \text{inl } p) = (X, p) \otimes (1, *)$ by $\rho_X x = \text{inl } x$ (this is not surjective). Finally we let $\alpha_{X,Y,Z} : ((X, p) \otimes (Y, q)) \otimes (Z, r) = ((X + Y) + Z, \text{inl } (\text{inl } p)) \rightarrow (X + (Y + Z), \text{inl } p) = (X, p) \otimes ((Y, q) \otimes (Z, r))$ be the obvious isomorphism.

(We note that **Ptd** has coproducts too: $(X, p) + (Y, q) = ((X + Y)/\sim, [\text{inl } p])$ where \sim is the equivalence relation on $X + Y$ induced by $\text{inl } p \sim \text{inr } q$.)

Example 2.3 Suppose given a monoidal category $(\mathcal{C}, \mathbb{1}, \otimes)$ together with a lax monoidal comonad $(D, \mathbf{e}, \mathbf{m})$ on \mathcal{C} . The category \mathcal{C} has a skew monoidal structure with $\mathbb{1}^D = \mathbb{1}$, $A \otimes^D B = A \otimes D B$. The unitors and associator are the following:

$$\lambda_A^D = \mathbb{1} \otimes D A \xrightarrow{\lambda_{DA}} D A \xrightarrow{\epsilon_A} A$$

$$\rho_A^D = A \xrightarrow{\rho_A} A \otimes \mathbb{1} \xrightarrow{A \otimes \mathbf{e}} A \otimes D \mathbb{1}$$

$$\alpha_{A,B,C}^D = (A \otimes D B) \otimes D C \xrightarrow{(A \otimes D B) \otimes \delta} (A \otimes D B) \otimes D (D C) \xrightarrow{\alpha_{A,DB,DC}} A \otimes (D B \otimes D (D C)) \xrightarrow{A \otimes \mathbf{m}_{B,C}} A \otimes D (B \otimes D C)$$

A similar skew monoidal category is also obtained from any oplax monoidal monad.

Example 2.4 Consider two categories \mathcal{J} and \mathcal{C} with a functor $J : \mathcal{J} \rightarrow \mathcal{C}$, and assume that the left Kan extension $\text{Lan}_J F : \mathcal{C} \rightarrow \mathcal{C}$ exists for every $F : \mathcal{J} \rightarrow \mathcal{C}$. Then the functor category $[\mathcal{J}, \mathcal{C}]$ has a skew monoidal structure given by $\mathbb{1} = J$, $F \otimes G = \text{Lan}_J F \circ G$. The unitors and associator are the canonical natural transformations $\lambda_F : \text{Lan}_J J \circ F \rightarrow F$, $\rho_F : F \rightarrow \text{Lan}_J F \circ J$, $\alpha_{F,G,H} : \text{Lan}_J (\text{Lan}_J F \circ G) \circ H \rightarrow \text{Lan}_J F \circ \text{Lan}_J G \circ H$. This category becomes properly monoidal under certain conditions on J : ρ is an isomorphism if J is fully-faithful, and λ is an isomorphism if J is dense. (This is the example from our relative monads work [1]. Relative monads on J are skew monoids in the skew monoidal category $[\mathcal{J}, \mathcal{C}]$.)

As our aim is to analyze the relationship of skew monoidal categories to a sequent calculus with the methods of structural proof theory, we will find it convenient to have an explicit description of the free skew monoidal category $\mathbf{Fsk}(\text{At})$ over a set At (whose elements we view as propositional letters, also called atoms) as a deductive calculus.

The objects are given by the set \mathbf{Fma} (of formulae). A formula is either an element X of At (an atomic formula); $\mathbb{1}$; or $A \otimes B$ where A, B are formulae.

The maps between two objects A and B are derivations of singleton-antecedent, singleton-succedent sequents $A \Rightarrow B$ constructed with the following inference rules

$$\begin{array}{c} \frac{}{A \Rightarrow A} \text{id} \quad \frac{A \Rightarrow B \quad B \Rightarrow C}{A \Rightarrow C} \text{comp} \quad \frac{A \Rightarrow C \quad B \Rightarrow D}{A \otimes B \Rightarrow C \otimes D} \otimes \\ \frac{}{\mathbb{1} \otimes A \Rightarrow A} \lambda \quad \frac{}{A \Rightarrow A \otimes \mathbb{1}} \rho \quad \frac{}{(A \otimes B) \otimes C \Rightarrow A \otimes (B \otimes C)} \alpha \end{array}$$

identified up to the least congruence given by the equations on Figure 1. In addition to the laws (a)–(e) above, these equations state that id and comp satisfy the laws of a category, that \otimes is functorial, and that λ , ρ and α are natural transformations. In the term notation for derivations, we write $g \circ f$ for $\text{comp } f g$ to agree with the standard categorical notation.

A necessary condition for the existence of a map $A \Rightarrow B$ is that A and B have the same underlying frontier of atoms (where the frontier ∂A of a formula A is defined by $\partial X = X$; $\partial \mathbb{1} = \cdot$; $\partial(A \otimes B) = \partial A, \partial B$). However, this is not sufficient. For example, there are no maps $X \Rightarrow \mathbb{1} \otimes X$, $X \otimes \mathbb{1} \Rightarrow X$ or $X \otimes (Y \otimes Z) \Rightarrow (X \otimes Y) \otimes Z$ (typing the inverses of the unitors and the associator in the monoidal case). Moreover, it is possible to have more than one map with the same domain and codomain. The prototypical examples are $\rho_{\mathbb{1}} \circ \lambda_{\mathbb{1}} \neq \text{id}_{\mathbb{1} \otimes \mathbb{1}} : \mathbb{1} \otimes \mathbb{1} \Rightarrow \mathbb{1} \otimes \mathbb{1}$, $\text{id}_{(X \otimes \mathbb{1}) \otimes Y} \neq (\rho_X \otimes \lambda_Y) \circ \alpha_{X, \mathbb{1}, Y}$ and $\text{id}_{X \otimes (\mathbb{1} \otimes Y)} \neq \alpha_{X, \mathbb{1}, Y} \circ (\rho_X \otimes \lambda_Y)$ (the alternative versions of axioms (a), (b) in the monoidal case).

3 Sequent Calculus

We now introduce a sequent calculus for skew monoidal categories inspired by the sequent calculus for the Tamari order [18].

Sequents are of the form $S \mid \Gamma \vdash A$, where the succedent A is a single formula and the antecedent is a pair of a *stoup* S and a *context* Γ . A stoup S is either nothing (written $-$) or a single formula, while a context Γ is a (possibly empty) list of formulae.

Derivations are constructed with the following inference rules:

$$\begin{array}{c} \frac{}{A \mid \vdash A} \text{ax} \quad \frac{A \mid \Gamma \vdash C}{- \mid A, \Gamma \vdash C} \text{shift} \\ \frac{- \mid \Gamma \vdash C}{\mathbb{1} \mid \Gamma \vdash C} \text{IL} \quad \frac{}{- \mid \vdash \mathbb{1}} \text{IR} \\ \frac{A \mid B, \Gamma \vdash C}{A \otimes B \mid \Gamma \vdash C} \otimes\text{L} \quad \frac{S \mid \Gamma \vdash A \quad - \mid \Delta \vdash B}{S \mid \Gamma, \Delta \vdash A \otimes B} \otimes\text{R} \end{array}$$

There are no primitive cut rules in this calculus, but we will shortly see that two cut rules are admissible. (In Section 5, we will describe a further restriction to *focused* derivations.)

Although these rules look very similar to the rules of the $\mathbb{1}, \otimes$ fragment of non-commutative intuitionistic linear logic (the sequent calculus of monoidal categories)—in particular, there is no left exchange rule, weakening or contraction—, there are two crucial differences. First, the left logical rules are restricted to apply only at the leftmost end of the antecedent, to the formula within the stoup. This restriction was present in the sequent

calculus for the Tamari order [18]. Second, and this is a new aspect, the stoup is allowed to be empty, permitting a distinction between antecedents of the form $A \mid \Gamma$ (with A inside the stoup) and antecedents of the form $- \mid A, \Gamma$ (with A outside the stoup). These ingredients are crucial for obtaining the correspondence with skew monoidal categories.

Let us demonstrate these restrictions in action on a few examples. We will state the exact relationship of skew monoidal categories and our sequent calculus later, but one of the results will be that there is a map $A \Rightarrow B$ in the free skew monoidal category iff the sequent $A \mid \vdash B$ is derivable (see Theorems 4.4 and 4.5).

In the sequent calculus of monoidal categories, one can build the derivation

$$\frac{\frac{\frac{\overline{X \vdash X} \text{ ax} \quad \overline{Y \vdash Y} \text{ ax}}{X, Y \vdash X \otimes Y} \otimes R \quad \overline{Z \vdash Z} \text{ ax}}{X, Y, Z \vdash (X \otimes Y) \otimes Z} \otimes R}{\frac{X, Y \otimes Z \vdash (X \otimes Y) \otimes Z} \otimes L}{X \otimes (Y \otimes Z) \vdash (X \otimes Y) \otimes Z} \otimes L}$$

corresponding to $\alpha_{X,Y,Z}^{-1} : X \otimes (Y \otimes Z) \Rightarrow (X \otimes Y) \otimes Z$. Notice that the second application of the $\otimes L$ rule (counting from the bottom) is to the formula second from the left in the antecedent. In the skew monoidal calculus, the corresponding sequent has no derivation; attempts to build one fail:

$$\frac{\frac{X \mid Y \otimes Z \vdash X \otimes Y \quad - \mid \vdash Z}{X \mid Y \otimes Z \vdash (X \otimes Y) \otimes Z} \otimes R}{X \otimes (Y \otimes Z) \mid \vdash (X \otimes Y) \otimes Z} \otimes L \quad \frac{\frac{X \mid \vdash X \otimes Y \quad - \mid Y \otimes Z \vdash Z}{X \mid Y \otimes Z \vdash (X \otimes Y) \otimes Z} \otimes R}{X \otimes (Y \otimes Z) \mid \vdash (X \otimes Y) \otimes Z} \otimes L$$

(Thanks to cut-freeness, the sequent calculus admits a root-first proof search strategy deciding derivability. We will say more about this in Section 5.)

In the same vein, one can build a derivation $X \otimes I \vdash X$ in the sequent calculus of monoidal categories (corresponding to $\rho_X^{-1} : X \otimes I \Rightarrow X$), but this relies on applicability of IL to the second formula in an antecedent, and there is no derivation of $X \otimes I \mid \vdash X$ in the skew monoidal sequent calculus.

Finally, corresponding to $\lambda_X^{-1} : X \Rightarrow I \otimes X$, one has the following derivation in the sequent calculus of monoidal categories:

$$\frac{\overline{\vdash I} \text{ IR} \quad \overline{X \vdash X} \text{ ax}}{X \vdash I \otimes X} \otimes R$$

But there is no such derivation in the skew monoidal sequent calculus:

$$\frac{X \mid \vdash I \quad - \mid \vdash X}{X \mid \vdash I \otimes X} \otimes R$$

This time the reason is that, while the context can be split freely in an $\otimes R$ inference, the stoup formula must go to the first premise.

At the same time, derivations corresponding to $\lambda_X, \rho_X, \alpha_{X,Y,X}$ can be smoothly constructed in our calculus despite the restrictions. They are needed and appear in the proof of Theorem 4.5 below.

As a more involved example, here is a derivation corresponding to the (incidentally) unique map $(X \otimes (I \otimes$

$$\frac{\frac{}{A \mid \vdash A} \text{ax} \quad \frac{\vdots g}{A \mid \Delta \vdash C} \text{scut}}{A \mid \Delta \vdash C} \text{scut} =_{\text{df}} \frac{\vdots g}{A \mid \Delta \vdash C}$$

- Case $f =_{\text{df}} \text{shift } f'$ for some $f' : A' \mid \Gamma' \vdash A$. In particular, $S = -$ and $\Gamma = A', \Gamma'$. We define:

$$\frac{\frac{\frac{\vdots f'}{A' \mid \Gamma' \vdash A} \text{shift} \quad \frac{\vdots g}{A \mid \Delta \vdash C} \text{scut}}{- \mid A', \Gamma' \vdash A} \text{scut}}{- \mid A', \Gamma', \Delta \vdash C} \text{scut} =_{\text{df}} \frac{\frac{\vdots f'}{A' \mid \Gamma' \vdash A} \quad \frac{\vdots g}{A \mid \Delta \vdash C} \text{scut}}{- \mid A', \Gamma', \Delta \vdash C} \text{shift}$$

- Case $f =_{\text{df}} \text{ll } f'$ for some $f' : - \mid \Gamma \vdash A$. In particular, $S = \text{l}$. We define:

$$\frac{\frac{\frac{\vdots f'}{- \mid \Gamma \vdash A} \text{ll} \quad \frac{\vdots g}{A \mid \Delta \vdash C} \text{scut}}{\text{l} \mid \Gamma, \Delta \vdash C} \text{scut}}{\text{l} \mid \Gamma, \Delta \vdash C} \text{scut} =_{\text{df}} \frac{\frac{\vdots f'}{- \mid \Gamma \vdash A} \quad \frac{\vdots g}{A \mid \Delta \vdash C} \text{scut}}{\text{l} \mid \Gamma, \Delta \vdash C} \text{ll}$$

- Case $f =_{\text{df}} \otimes \text{L } f'$ for some $f' : B \mid D, \Gamma \vdash A$. In particular, $S = B \otimes D$. We define:

$$\frac{\frac{\frac{\vdots f'}{B \mid D, \Gamma \vdash A} \otimes \text{L} \quad \frac{\vdots g}{A \mid \Delta \vdash C} \text{scut}}{B \otimes D \mid \Gamma, \Delta \vdash C} \text{scut}}{B \otimes D \mid \Gamma, \Delta \vdash C} \text{scut} =_{\text{df}} \frac{\frac{\vdots f'}{B \mid D, \Gamma \vdash A} \quad \frac{\vdots g}{A \mid \Delta \vdash C} \text{scut}}{B \otimes D \mid \Gamma, \Delta \vdash C} \otimes \text{L}$$

- Case $f =_{\text{df}} \text{IR}$. In particular, $S = -$, Γ is empty and $A = \text{l}$.

$$\frac{\frac{}{- \mid \vdash \text{l}} \text{IR} \quad \frac{\vdots g}{\text{l} \mid \Delta \vdash C} \text{scut}}{- \mid \Delta \vdash C} \text{scut} =_{\text{df}} \frac{\vdots g}{- \mid \Delta \vdash C} \text{ll}^{-1}$$

- Case $f =_{\text{df}} \otimes \text{R } f_1 f_2$ for some $f_1 : S \mid \Gamma_1 \vdash A_1$ and $f_2 : - \mid \Gamma_2 \vdash A_2$. In particular, $\Gamma = \Gamma_1, \Gamma_2$ and $A = A_1 \otimes A_2$. At this point, we proceed by induction on the derivation $g : A_1 \otimes A_2 \mid \Delta \vdash C$. Notice that g can neither be of the form shift nor IR since its stoup is non-empty. Moreover, g can neither be of the form ll since its stoup is not equal to l . Therefore, we only have to check the three remaining cases.

- Case $g =_{\text{df}} \text{ax}$. In particular, Δ is empty and $C = A_1 \otimes A_2$. We define:

$$\frac{\frac{\frac{\vdots f_1}{S \mid \Gamma_1 \vdash A_1} \quad \frac{\vdots f_2}{- \mid \Gamma_2 \vdash A_2}}{S \mid \Gamma_1, \Gamma_2 \vdash A_1 \otimes A_2} \otimes \text{R} \quad \frac{\vdots g}{A_1 \otimes A_2 \mid \vdash A_1 \otimes A_2} \text{ax}}{S \mid \Gamma_1, \Gamma_2 \vdash A_1 \otimes A_2} \text{scut} =_{\text{df}} \frac{\frac{\vdots f_1}{S \mid \Gamma_1 \vdash A_1} \quad \frac{\vdots f_2}{- \mid \Gamma_2 \vdash A_2}}{S \mid \Gamma_1, \Gamma_2 \vdash A_1 \otimes A_2} \otimes \text{R}$$

- Case $g =_{\text{df}} \otimes \text{R } g_1 g_2$ for some $g_1 : A_1 \otimes A_2 \mid \Delta_1 \vdash C_1$ and $g_2 : - \mid \Delta_2 \vdash C_2$. In particular, $\Delta = \Delta_1, \Delta_2$ and $C = C_1 \otimes C_2$. We define:

$$\frac{\frac{\frac{\frac{\vdots f_1}{S \mid \Gamma_1 \vdash A_1} \quad \frac{\vdots f_2}{- \mid \Gamma_2 \vdash A_2}}{S \mid \Gamma_1, \Gamma_2 \vdash A_1 \otimes A_2} \otimes \text{R} \quad \frac{\frac{\frac{\vdots g_1}{A_1 \otimes A_2 \mid \Delta_1 \vdash C_1} \quad \frac{\vdots g_2}{- \mid \Delta_2 \vdash C_2}}{A_1 \otimes A_2 \mid \Delta_1, \Delta_2 \vdash C_1 \otimes C_2} \otimes \text{R}}{S \mid \Gamma_1, \Gamma_2, \Delta_1, \Delta_2 \vdash C_1 \otimes C_2} \text{scut}}{S \mid \Gamma_1, \Gamma_2, \Delta_1, \Delta_2 \vdash C_1 \otimes C_2} \text{scut}} =_{\text{df}} \frac{\frac{\frac{\vdots f_1}{S \mid \Gamma_1 \vdash A_1} \quad \frac{\vdots f_2}{- \mid \Gamma_2 \vdash A_2}}{S \mid \Gamma_1, \Gamma_2 \vdash A_1 \otimes A_2} \otimes \text{R} \quad \frac{\frac{\vdots g_1}{A_1 \otimes A_2 \mid \Delta_1 \vdash C_1} \text{scut}}{S \mid \Gamma_1, \Gamma_2, \Delta_1 \vdash C_1} \text{scut} \quad \frac{\vdots g_2}{- \mid \Delta_2 \vdash C_2} \otimes \text{R}}{S \mid \Gamma_1, \Gamma_2, \Delta_1, \Delta_2 \vdash C_1 \otimes C_2} \otimes \text{R}$$

- Case $g =_{\text{df}} \otimes L g'$ for some $g' : A_1 \mid A_2, \Delta \vdash C$. We define:

$$\frac{\frac{\frac{\vdots f_1}{S \mid \Gamma_1 \vdash A_1} \quad \frac{\vdots f_2}{- \mid \Gamma_2 \vdash A_2}}{S \mid \Gamma_1, \Gamma_2 \vdash A_1 \otimes A_2} \otimes R \quad \frac{\frac{\vdots g'}{A_1 \mid A_2, \Delta \vdash C}}{A_1 \otimes A_2 \mid \Delta \vdash C} \otimes L}{S \mid \Gamma_1, \Gamma_2, \Delta \vdash C} \text{scut}}{S \mid \Gamma_1, \Gamma_2, \Delta \vdash C} \text{scut} =_{\text{df}} \frac{\frac{\frac{\vdots f_2}{- \mid \Gamma_2 \vdash A_2} \quad \frac{\frac{\vdots f_1}{S \mid \Gamma_1 \vdash A_1} \quad \frac{\vdots g'}{A_1 \mid A_2, \Delta \vdash C}}{S \mid \Gamma_1, A_2, \Delta \vdash C} \text{scut}}{S \mid \Gamma_1, \Gamma_2, \Delta \vdash C} \text{ccut}}{S \mid \Gamma_1, \Gamma_2, \Delta \vdash C} \text{ccut}}$$

We continue with the proof of ccut . Let $f : - \mid \Gamma \vdash A$ and $g : S \mid \Delta_0, A, \Delta_1 \vdash C$ be two derivations. The proof proceeds by induction on g . Notice that the g can neither be of the form ax nor IR , since the context of its endsequent is non-empty. Therefore, we only have to check the four remaining cases.

- Case $g =_{\text{df}} \text{shift } g'$, for some $g' : A' \mid \Delta' \vdash C$. In particular, $S = -$ and $\Delta_0, A, \Delta_1 = A', \Delta'$. We proceed by checking if the context Δ_0 is empty or not.
 - If Δ_0 is empty, then $A' = A$ and $\Delta' = \Delta_1$. We define:

$$\frac{\frac{\frac{\vdots f}{- \mid \Gamma \vdash A} \quad \frac{\frac{\vdots g'}{A \mid \Delta_1 \vdash C}}{- \mid A, \Delta_1 \vdash C} \text{shift}}{- \mid \Gamma, \Delta_1 \vdash C} \text{ccut}}{- \mid \Gamma, \Delta_1 \vdash C} \text{ccut} =_{\text{df}} \frac{\frac{\vdots f}{- \mid \Gamma \vdash A} \quad \frac{\vdots g'}{A \mid \Delta_1 \vdash C}}{- \mid \Gamma, \Delta_1 \vdash C} \text{scut}}$$

- If $\Delta_0 =_{\text{df}} A'', \Delta'_0$, then $A'' = A'$ and $\Delta' = \Delta'_0, A, \Delta_1$. We define:

$$\frac{\frac{\frac{\vdots f}{- \mid \Gamma \vdash A} \quad \frac{\frac{\frac{\vdots g'}{A' \mid \Delta'_0, A, \Delta_1 \vdash C}}{- \mid A', \Delta'_0, A, \Delta_1 \vdash C} \text{shift}}{- \mid A', \Delta'_0, \Gamma, \Delta_1 \vdash C} \text{ccut}}{- \mid A', \Delta'_0, \Gamma, \Delta_1 \vdash C} \text{ccut} =_{\text{df}} \frac{\frac{\frac{\vdots f}{- \mid \Gamma \vdash A} \quad \frac{\vdots g'}{A' \mid \Delta'_0, A, \Delta_1 \vdash C}}{A' \mid \Delta'_0, \Gamma, \Delta_1 \vdash C} \text{ccut}}{- \mid A', \Delta'_0, \Gamma, \Delta_1 \vdash C} \text{shift}}$$

- Case $g =_{\text{df}} \text{IL } g'$ for some $g' : - \mid \Delta_0, A, \Delta_1 \vdash C$. In particular, $S = \text{I}$. We define:

$$\frac{\frac{\frac{\vdots f}{- \mid \Gamma \vdash A} \quad \frac{\frac{\frac{\vdots g'}{- \mid \Delta_0, A, \Delta_1 \vdash C}}{\text{I} \mid \Delta_0, A, \Delta_1 \vdash C} \text{IL}}{\text{I} \mid \Delta_0, \Gamma, \Delta_1 \vdash C} \text{ccut}}{\text{I} \mid \Delta_0, \Gamma, \Delta_1 \vdash C} \text{ccut} =_{\text{df}} \frac{\frac{\frac{\vdots f}{- \mid \Gamma \vdash A} \quad \frac{\vdots g'}{- \mid \Delta_0, A, \Delta_1 \vdash C}}{- \mid \Delta_0, \Gamma, \Delta_1 \vdash C} \text{ccut}}{\text{I} \mid \Delta_0, \Gamma, \Delta_1 \vdash C} \text{IL}}$$

- Case $g =_{\text{df}} \otimes L g'$ for some $g' : B \otimes D \mid \Delta_0, A, \Delta_1 \vdash C$. In particular, $S = B \otimes D$. We define:

$$\frac{\frac{\frac{\vdots f}{- \mid \Gamma \vdash A} \quad \frac{\frac{\frac{\vdots g'}{B \mid D, \Delta_0, A, \Delta_1 \vdash C}}{B \otimes D \mid \Delta_0, A, \Delta_1 \vdash C} \otimes L}}{B \otimes D \mid \Delta_0, \Gamma, \Delta_1 \vdash C} \text{ccut} =_{\text{df}} \frac{\frac{\frac{\vdots f}{- \mid \Gamma \vdash A} \quad \frac{\vdots g'}{B \mid D, \Delta_0, A, \Delta_1 \vdash C}}{B \mid D, \Delta_0, \Gamma, \Delta_1 \vdash C} \text{ccut}}{B \otimes D \mid \Delta_0, \Gamma, \Delta_1 \vdash C} \otimes L}$$

- Case $g =_{\text{df}} \otimes R g_1 g_2$ for some $g_1 : S \mid \Lambda_1 \vdash C_1$ and $g_2 : - \mid \Lambda_2 \vdash C_2$. In particular, $C = C_1 \otimes C_2$ and $\Delta_0, A, \Delta_1 = \Lambda_1, \Lambda_2$. We proceed by checking if the formula A occurs in Λ_1 or in Λ_2 .
 - If A occurs in Λ_1 , we have $\Lambda_1 = \Delta_0, A, \Delta'_1$ and $\Delta_1 = \Delta'_1, \Lambda_2$. We define:

$$\frac{\frac{\frac{\frac{\vdots f}{- \mid \Gamma \vdash A} \quad \frac{\frac{\frac{\vdots g_1}{S \mid \Delta_0, A, \Delta'_1 \vdash C_1} \quad \frac{\vdots g_2}{- \mid \Lambda_2 \vdash C_2}}{S \mid \Delta_0, A, \Delta'_1, \Lambda_2 \vdash C_1 \otimes C_2} \otimes R}}{S \mid \Delta_0, \Gamma, \Delta'_1, \Lambda_2 \vdash C_1 \otimes C_2} \text{ccut} =_{\text{df}} \frac{\frac{\frac{\vdots f}{- \mid \Gamma \vdash A} \quad \frac{\frac{\vdots g_1}{S \mid \Delta_0, A, \Delta'_1 \vdash C_1}}{S \mid \Delta_0, \Gamma, \Delta'_1 \vdash C_1} \text{ccut}}{S \mid \Delta_0, \Gamma, \Delta'_1, \Lambda_2 \vdash C_1 \otimes C_2} \otimes R \quad \frac{\vdots g_2}{- \mid \Lambda_2 \vdash C_2} \otimes R}}$$

- If A occurs in Λ_2 , we have $\Lambda_2 = \Delta'_0, A, \Delta_1$ and $\Delta_0 = \Lambda_1, \Delta'_0$. We define:

$$\frac{-|\Gamma \vdash A \quad \frac{\frac{\frac{\vdots f}{-|\Gamma \vdash A} \quad \frac{\frac{\vdots g_1}{S \mid \Lambda_1 \vdash C_1} \quad -|\Delta'_0, A, \Delta_1 \vdash C_2}{S \mid \Lambda_1, \Delta'_0, A, \Delta_1 \vdash C_1 \otimes C_2} \otimes R}{S \mid \Lambda_1, \Delta'_0, \Gamma, \Delta_1 \vdash C_1 \otimes C_2} \text{ccut}}{S \mid \Lambda_1, \Delta'_0, \Gamma, \Delta_1 \vdash C_1 \otimes C_2} \otimes R}{S \mid \Lambda_1, \Delta'_0, \Gamma, \Delta_1 \vdash C_1 \otimes C_2} \text{ccut} =_{\text{df}} \frac{\frac{\frac{\vdots g_1}{S \mid \Lambda_1 \vdash C_1} \quad -|\Gamma \vdash A \quad -|\Delta'_0, A, \Delta_1 \vdash C_2}{-|\Delta'_0, \Gamma, \Delta_1 \vdash C_2} \otimes R}{S \mid \Lambda_1, \Delta'_0, \Gamma, \Delta_1 \vdash C_1 \otimes C_2} \otimes R}{S \mid \Lambda_1, \Delta'_0, \Gamma, \Delta_1 \vdash C_1 \otimes C_2} \text{ccut} \quad \square$$

The cut rules obey a number of equations reminiscent of the unit and associativity laws of multicategories. More precisely, our sequent calculus can be seen as a particular generalized multicategory in the sense of [12, Ch. 4]. We give the details of this connection in Section A.

Proposition 3.4 *The equations in Figure 2 hold.*

This proposition is used in the proofs of Theorems 4.14 and 4.19.

4 Adequacy

In this section, we show the connection between the categorical calculus and the sequent calculus. We start by showing how to interpret antecedents as formulae. This is performed in several steps, by first specifying how to interpret stoups and contexts.

Stoups are interpreted as formulae in \mathbf{Fma} :

$$\llbracket - \rrbracket =_{\text{df}} \text{l} \quad \llbracket A \rrbracket =_{\text{df}} A.$$

A context is interpreted as an endomap on \mathbf{Fma} . Given a context Γ , we write $C \llbracket \Gamma \rrbracket$ for the application of the interpretation of Γ to a formula C . The interpretation of contexts is the right action of contexts on formulae induced by \otimes :

$$C \llbracket \rrbracket =_{\text{df}} C, \quad C \llbracket A, \Gamma \rrbracket =_{\text{df}} (C \otimes A) \llbracket \Gamma \rrbracket$$

Using the interpretation of stoups and contexts, we define the interpretation of antecedents as formulae in \mathbf{Fma} :

$$\llbracket S \mid \Gamma \rrbracket =_{\text{df}} \llbracket S \rrbracket \llbracket \Gamma \rrbracket$$

Explicitly, the interpretation of antecedents works as follows. Let $\Gamma = A_1, \dots, A_n$. Given a formula A , we have $\llbracket A \mid \Gamma \rrbracket = (\dots((A \otimes A_1) \otimes A_2) \dots) \otimes A_n$. We also have $\llbracket - \mid \Gamma \rrbracket = (\dots((\text{l} \otimes A_1) \otimes A_2) \dots) \otimes A_n$, i.e., $\llbracket - \mid \Gamma \rrbracket = \llbracket \text{l} \mid \Gamma \rrbracket$.

In other words, depending on whether a sequent has an empty or non-empty stoup, the antecedent has to be interpreted with an l as the leftmost factor in the big tensor product or not. This is an important observation and the reason why stoups are needed in the first place: to give a correct interpretation of sequents. In fact, suppose we formulated the rules of the sequent calculus conflating the stoup and the context together into a flat antecedent. We would still interpret antecedents as big tensor products, but we could in principle choose to add an l in the leftmost position or not. In both cases, the interpretation of sequents would be unsound. In fact, if the interpretation of an antecedent A_1, \dots, A_n were $(\dots(A_1 \otimes A_2) \otimes \dots) \otimes A_n$, then the derivable sequent $X \vdash \text{l} \otimes X$ where X is an atom would be interpreted as $X \Rightarrow \text{l} \otimes X$, which cannot be derived in the categorical calculus. On the other hand, if the interpretation of an antecedent A_1, \dots, A_n were $(\dots((\text{l} \otimes A_1) \otimes A_2) \dots) \otimes A_n$, then the derivable sequent $X \otimes Y \vdash (\text{l} \otimes X) \otimes Y$ where X and Y are atoms would be interpreted as $\text{l} \otimes (X \otimes Y) \Rightarrow (\text{l} \otimes X) \otimes Y$, which is again not derivable in the categorical calculus.

We now show that the sequent calculus is sound. This relies on three lemmata involving the interpretation of antecedents.

Lemma 4.1 *For any derivation $f : A \Rightarrow B$ and context Γ , there is a derivation $\llbracket f \mid \Gamma \rrbracket : \llbracket A \mid \Gamma \rrbracket \Rightarrow \llbracket B \mid \Gamma \rrbracket$.*

Proof. We proceed by induction on Γ . If Γ is empty, then we take $\llbracket f \mid \rrbracket =_{\text{df}} f$. If $\Gamma = C, \Gamma'$, then we take $\llbracket f \mid C, \Gamma' \rrbracket =_{\text{df}} \llbracket f \otimes \text{id}_C \mid \Gamma' \rrbracket$. \square

Lemma 4.2 *For any formulae A, B and context Γ , there is a derivation $\psi_{A,B,\Gamma} : \llbracket A \otimes B \mid \Gamma \rrbracket \Rightarrow A \otimes \llbracket B \mid \Gamma \rrbracket$.*

Proof. We proceed by induction on Γ . If Γ is empty, then we take $\psi_{A,B,(\)} =_{\text{df}} \text{id}_{A \otimes B}$. If $\Gamma = C, \Gamma'$, then we take $\psi_{A,B,(C,\Gamma')} =_{\text{df}} \psi_{A,B \otimes C, \Gamma'} \circ \llbracket \alpha \mid \Gamma' \rrbracket$. \square

Lemma 4.3 *For any stoup S and contexts Γ, Δ , there is a derivation $\varphi_{S,\Gamma,\Delta} : \llbracket S \mid \Gamma, \Delta \rrbracket \Rightarrow \llbracket S \mid \Gamma \rrbracket \otimes \llbracket - \mid \Delta \rrbracket$.*

$$\begin{array}{c}
 \begin{array}{c} \vdots \text{ ax} \\ \vdots f \\ A \vdash A \quad A \mid \Delta \vdash C \\ \hline A \mid \Delta \vdash C \quad \text{scut} \end{array} \\
 = \\
 \begin{array}{c} \text{ax} \\ \frac{A \mid \vdash A}{- \mid A \vdash A} \quad \begin{array}{c} \vdots f \\ S \mid \Delta_0, A, \Delta_1 \vdash C \end{array} \\ \hline S \mid \Delta_0, A, \Delta_1 \vdash C \quad \text{ccut} \end{array} \\
 = \\
 \begin{array}{c} \vdots f \\ \frac{S \mid \Gamma \vdash A \quad \overline{A \mid \vdash A} \quad \text{ax}}{S \mid \Gamma \vdash A} \quad \text{scut} \end{array} \\
 = \\
 \begin{array}{c} \vdots g \quad \vdots h \\ \frac{\vdots f \quad A \mid \Delta \vdash B \quad B \mid \lambda \vdash C}{S \mid \Gamma \vdash A \quad A \mid \Delta, \lambda \vdash C} \quad \text{scut} \\ \hline S \mid \Gamma, \Delta, \lambda \vdash C \quad \text{scut} \end{array} \\
 = \\
 \begin{array}{c} \vdots f \quad \vdots g \quad \vdots h \\ \frac{\vdots f \quad S \mid \Delta_0, A, \Delta_1 \vdash B \quad B \mid \lambda \vdash C}{- \mid \Gamma \vdash A} \quad \text{scut} \\ \hline S \mid \Delta_0, A, \Delta_1, \lambda \vdash C \quad \text{ccut} \end{array} \\
 = \\
 \begin{array}{c} \vdots g \quad \vdots h \\ \frac{\vdots f \quad - \mid \Delta_0, A, \Delta_1 \vdash B \quad S \mid \lambda_0, B, \lambda_1 \vdash C}{- \mid \Gamma \vdash A} \quad \text{ccut} \\ \hline S \mid \lambda_0, \Delta_0, \Gamma, \Delta_1, \lambda_1 \vdash C \quad \text{ccut} \end{array} \\
 = \\
 \begin{array}{c} \vdots f_1 \quad \vdots f_2 \quad \vdots g \\ \frac{S \mid \Gamma_1 \vdash A_1 \quad - \mid \Gamma_2 \vdash A_2 \quad A_1 \mid \Delta_1, A_2, \Delta_2 \vdash C}{S \mid \Gamma_1, \Delta_1, \Gamma_2, \Delta_2 \vdash C} \quad \text{ccut} \\ \hline S \mid \Gamma_1 \vdash A_1 \quad - \mid \Gamma_2 \vdash A_2 \quad A_1 \mid \Delta_1, A_2, \Delta_2 \vdash C \quad \text{ccut} \end{array} \\
 = \\
 \begin{array}{c} \vdots f_1 \quad \vdots f_2 \quad \vdots g \\ \frac{S \mid \Gamma_1 \vdash A_1 \quad - \mid \Gamma_2 \vdash A_2 \quad S \mid \Delta_0, A_1, \Delta_1, A_2, \Delta_2 \vdash C}{- \mid \Gamma_1 \vdash A_1} \quad \text{ccut} \\ \hline S \mid \Delta_0, \Gamma_1, \Delta_1, \Gamma_2, \Delta_2 \vdash C \quad \text{ccut} \end{array} \\
 \end{array}$$

Fig. 2. Generalized multicategory equations

Proof. It is sufficient to construct $\varphi'_{A,\Gamma,\Delta} : A \langle \Gamma, \Delta \rangle \Rightarrow A \langle \Gamma \rangle \otimes \llbracket - \mid \Delta \rrbracket$ for $A : \mathbf{Fma}$, and define $\varphi_{S,\Gamma,\Delta} =_{\text{df}} \varphi'_{\llbracket S \mid \Gamma, \Delta \rrbracket}$. We proceed by induction on Γ . If Γ is empty, then we have to construct $\varphi'_{A,(),\Delta} : \llbracket A \mid \Delta \rrbracket \Rightarrow A \otimes \llbracket - \mid \Delta \rrbracket$. We take $\varphi'_{A,(),\Delta} =_{\text{df}} \psi_{A,!,\Delta} \circ \llbracket \rho \mid \Delta \rrbracket$. If $\Gamma = C, \Gamma'$, then we take $\varphi'_{A,(C,\Gamma'),\Delta} =_{\text{df}} \varphi'_{A \otimes C, \Gamma', \Delta}$. \square

Theorem 4.4 (Soundness) *For any derivation $f : S \mid \Gamma \vdash C$, there is a derivation $\text{sound } f : \llbracket S \mid \Gamma \rrbracket \Rightarrow C$. As a special case, for $f : A \mid \vdash C$, we have a derivation $\text{sound } f : A \Rightarrow C$.*

Proof. The proof proceeds by induction on f .

- Case $f =_{\text{df}} \text{ax}$. In particular, we have $S = C$ and Γ is empty. We define:

$$\text{sound} \left(\frac{}{C \mid \vdash C} \text{ax} \right) =_{\text{df}} \frac{\overline{C \Rightarrow C} \text{ id}}{\llbracket C \mid \rrbracket \Rightarrow C}$$

- Case $f =_{\text{df}} \text{shift } f'$, for some $f' : A \mid \Gamma' \vdash C$. In particular, $S = -$ and $\Gamma = A, \Gamma'$. We define:

$$\text{sound} \left(\frac{\frac{\vdots f'}{A \mid \Gamma' \vdash C} \text{shift}}{- \mid A, \Gamma' \vdash C} \right) =_{\text{df}} \frac{\frac{\llbracket \text{l} \otimes A \mid \Gamma' \rrbracket \Rightarrow A \mid \Gamma'}{\llbracket \text{l} \otimes A \mid \Gamma' \rrbracket \Rightarrow C} \llbracket \lambda \mid \Gamma' \rrbracket \quad \frac{\vdots \text{sound } f'}{A \mid \Gamma' \Rightarrow C} \text{comp}}{\llbracket - \mid A, \Gamma' \rrbracket \Rightarrow C}$$

- Case $f =_{\text{df}} \text{ll } f'$, for some $f' : - \mid \Gamma \vdash C$. In particular, $S = \text{l}$. We define:

$$\text{sound} \left(\frac{\frac{\vdots f'}{- \mid \Gamma \vdash C} \text{ll}}{\text{l} \mid \Gamma \vdash C} \right) =_{\text{df}} \frac{\frac{\vdots \text{sound } f'}{- \mid \Gamma \Rightarrow C}}{\llbracket \text{l} \mid \Gamma \rrbracket \Rightarrow C}$$

- Case $f =_{\text{df}} \otimes \text{L } f'$, for some $f' : A \mid B, \Gamma \vdash C$. In particular, $S = A \otimes B$. We define:

$$\text{sound} \left(\frac{\frac{\frac{\vdots f'}{A \mid B, \Gamma \vdash C} \otimes \text{L}}{A \otimes B \mid \Gamma \vdash C}}{\otimes \text{L}} \right) =_{\text{df}} \frac{\frac{\frac{\vdots \text{sound } f'}{A \mid B, \Gamma \Rightarrow C}}{\llbracket A \mid B, \Gamma \rrbracket \Rightarrow C}}{\llbracket A \otimes B \mid \Gamma \rrbracket \Rightarrow C}$$

- Case $f =_{\text{df}} \text{lr}$. In particular, $S = -$, $A = \text{l}$ and Γ is empty.

$$\text{sound} \left(\frac{}{- \mid \vdash \text{l}} \text{lr} \right) =_{\text{df}} \frac{\overline{\text{l} \Rightarrow \text{l}} \text{ id}}{\llbracket - \mid \rrbracket \Rightarrow \text{l}}$$

- Case $f =_{\text{df}} \otimes \text{R } f_1 f_2$, for some $f_1 : S \mid \Gamma_1 \vdash C_1$ and $f_2 : - \mid \Gamma_2 \vdash C_2$. In particular $\Gamma = \Gamma_1, \Gamma_2$ and $C = C_1 \otimes C_2$. We define:

$$\text{sound} \left(\frac{\frac{\frac{\frac{\vdots f_1}{S \mid \Gamma_1 \vdash C_1} \quad \frac{\vdots f_2}{- \mid \Gamma_2 \vdash C_2}}{S \mid \Gamma_1, \Gamma_2 \vdash C_1 \otimes C_2} \otimes \text{R}}{\otimes \text{R}} \right) =_{\text{df}} \frac{\frac{\llbracket S \mid \Gamma_1, \Gamma_2 \rrbracket \Rightarrow \llbracket S \mid \Gamma_1 \rrbracket \otimes \llbracket - \mid \Gamma_2 \rrbracket \quad \varphi_{S, \Gamma_1, \Gamma_2} \quad \frac{\frac{\vdots \text{sound } f_1}{\llbracket S \mid \Gamma_1 \rrbracket \Rightarrow C_1} \quad \frac{\vdots \text{sound } f_2}{\llbracket - \mid \Gamma_2 \rrbracket \Rightarrow C_2}}{\llbracket S \mid \Gamma_1 \rrbracket \otimes \llbracket - \mid \Gamma_2 \rrbracket \Rightarrow C_1 \otimes C_2} \otimes \text{comp}}{\llbracket S \mid \Gamma_1, \Gamma_2 \rrbracket \Rightarrow C_1 \otimes C_2}$$

\square

Next, we show that the sequent calculus is complete. Crucially, this relies on admissibility of scut , which we had to prove admissible together with ccut . (We chose to not have primitive cut rules in our sequent calculus, but completeness depends on their availability. Alternatively, one could take scut and ccut as primitive rules, and give a separate proof of cut elimination. This would slightly simplify the level of conceptual dependency, although it amounts to essentially the same amount of work in the end.)

Theorem 4.5 (Completeness) *For any derivation $f : A \Rightarrow C$, there is a derivation $\text{cmplt } f : A \mid \vdash C$.*

Proof. Just like the proof of soundness, the proof is by induction on the given derivation f .

- Case $f = \text{id}_A$. In particular, $A = C$. We define:

$$\text{cmplt} \left(\frac{}{A \Rightarrow A} \text{id} \right) =_{\text{df}} \frac{}{A \mid \vdash A} \text{ax}$$

- Case $f = h \circ g$ where $g : A \Rightarrow B$ and $h : B \Rightarrow C$. This is the case where we need `scut`. We define:

$$\text{cmplt} \left(\frac{\frac{\frac{\vdots}{A \Rightarrow B} g \quad \frac{\vdots}{B \Rightarrow C} h}{A \Rightarrow C} \text{comp}}{} \right) =_{\text{df}} \frac{\frac{\frac{\vdots}{A \mid \vdash B} \text{cmplt } g \quad \frac{\vdots}{B \mid \vdash C} \text{cmplt } h}{A \mid \vdash C} \text{scut}}{} \text{scut}$$

- Case $f = f_1 \otimes f_2$ where $f_1 : A_1 \Rightarrow C_1$ and $f_2 : A_2 \Rightarrow C_2$. In particular, $A = A_1 \otimes A_2$ and $C = C_1 \otimes C_2$. We define:

$$\text{cmplt} \left(\frac{\frac{\frac{\vdots}{A_1 \Rightarrow C_1} f_1 \quad \frac{\vdots}{A_2 \Rightarrow C_2} f_2}{A_1 \otimes A_2 \Rightarrow C_1 \otimes C_2} \otimes}{} \right) =_{\text{df}} \frac{\frac{\frac{\frac{\vdots}{A_1 \mid \vdash C_1} \text{cmplt } f_1 \quad \frac{\frac{\vdots}{A_2 \mid \vdash C_2} \text{cmplt } f_2}{- \mid A_2 \vdash C_2} \text{shift}}{A_1 \mid A_2 \vdash C_1 \otimes C_2} \otimes \text{R}}{A_1 \otimes A_2 \mid \vdash C_1 \otimes C_2} \otimes \text{L}}{} \otimes \text{L}$$

- Case $f = \lambda_C$. In particular, $A = \mathbb{1} \otimes C$. We define:

$$\text{cmplt} \left(\frac{}{\mathbb{1} \otimes C \Rightarrow C} \lambda \right) =_{\text{df}} \frac{\frac{\frac{\frac{}{C \mid \vdash C} \text{ax}}{- \mid C \vdash C} \text{shift}}{\mathbb{1} \mid C \vdash C} \text{IL}}{\mathbb{1} \otimes C \mid \vdash C} \otimes \text{L}}{} \otimes \text{L}$$

- Case $f = \rho_A$. In particular, $C = A \otimes \mathbb{1}$. We define:

$$\text{cmplt} \left(\frac{}{A \Rightarrow A \otimes \mathbb{1}} \rho \right) =_{\text{df}} \frac{\frac{\frac{}{A \mid \vdash A} \text{ax}}{- \mid \vdash \mathbb{1}} \text{IR}}{A \mid \vdash A \otimes \mathbb{1}} \otimes \text{R}}{} \otimes \text{R}$$

- Case $f = \alpha_{A', B, C'}$. In particular, $A = (A' \otimes B) \otimes C'$ and $C = A' \otimes (B \otimes C')$. We define:

$$\text{cmplt} \left(\frac{}{(A' \otimes B) \otimes C' \Rightarrow A' \otimes (B \otimes C')} \alpha \right) =_{\text{df}} \frac{\frac{\frac{\frac{\frac{\frac{}{A' \mid \vdash A'} \text{ax}}{- \mid B, C' \vdash B \otimes C'} \text{shift}}{B \mid C' \vdash B \otimes C'} \otimes \text{R}}{B \mid \vdash B} \text{ax}}{- \mid C' \vdash C'} \text{shift}}{A' \mid B, C' \vdash A' \otimes (B \otimes C')} \otimes \text{R}}{A' \otimes B \mid C' \vdash A' \otimes (B \otimes C')} \otimes \text{L}}{(A' \otimes B) \otimes C' \mid \vdash A' \otimes (B \otimes C')} \otimes \text{L}$$

□

The function `sound` is a left inverse of `cmplt` up to \doteq . In order to prove this, we define two categorical rules, `ssubst` and `csubst`. These rules corresponds to the rules `scut` and `ccut` of the sequent calculus. This is made precise by showing that the function `sound` sends a derivation ending with an application of the rule `scut` (or `ccut`) into a derivation ending with an application of the rule `ssubst` (or `csubst`).

The definitions of `ssubst` and `csubst` rely on the following lemma involving the interpretation of antecedents.

Lemma 4.6 *For any stoup S and contexts Γ, Δ , there is a derivation $\theta_{S, \Gamma, \Delta} : \llbracket S \mid \Gamma, \Delta \rrbracket \Rightarrow \llbracket \llbracket S \mid \Gamma \rrbracket \mid \Delta \rrbracket$.*

Proof. It is sufficient to construct $\theta'_{A, \Gamma, \Delta} : A \langle \Gamma, \Delta \rangle \Rightarrow (A \langle \Gamma \rangle) \langle \Delta \rangle$ for $A : \text{Fma}$, and define $\theta_{S, \Gamma, \Delta} =_{\text{df}} \theta'_{\llbracket S \mid \Gamma, \Delta \rrbracket}$. We proceed by induction on Γ . If Γ is empty, we take $\theta'_{A, (\cdot), \Delta} =_{\text{df}} \text{id}$. If $\Gamma = C, \Gamma'$, then we take $\theta'_{A, (C, \Gamma'), \Delta} =_{\text{df}} \theta'_{A \otimes C, \Gamma, \Delta}$. □

Lemma 4.7 *The following rules are admissible.*

$$\frac{\frac{\llbracket S \mid \Gamma \rrbracket \Rightarrow A \quad \llbracket A \mid \Delta \rrbracket \Rightarrow C}{\llbracket S \mid \Gamma, \Delta \rrbracket \Rightarrow C} \text{ ssubst}}{\llbracket - \mid \Gamma \rrbracket \Rightarrow A \quad \llbracket S \mid \Delta_0, A, \Delta_1 \rrbracket \Rightarrow C} \text{ csbst} \quad \frac{\llbracket S \mid \Gamma \rrbracket \Rightarrow A \quad \llbracket A \mid \Delta \rrbracket \Rightarrow C}{\llbracket S \mid \Gamma, \Delta \rrbracket \Rightarrow C} \text{ ssubst}$$

Lemma 4.8 *The following equations hold:*

$$\text{sound} \left(\frac{\frac{\frac{\vdots f \quad \vdots g}{S \mid \Gamma \vdash A \quad A \mid \Delta \vdash C} \text{ scut}}{S \mid \Gamma, \Delta \vdash C}}{\vdots \text{ sound } f \quad \vdots \text{ sound } g} \right) \doteq \frac{\frac{\llbracket S \mid \Gamma \rrbracket \Rightarrow A \quad \llbracket A \mid \Delta \rrbracket \Rightarrow C}{\llbracket S \mid \Gamma, \Delta \rrbracket \Rightarrow C} \text{ ssubst}}{\llbracket - \mid \Gamma \rrbracket \Rightarrow A \quad \llbracket S \mid \Delta_0, A, \Delta_1 \rrbracket \Rightarrow C} \text{ csbst}$$

$$\text{sound} \left(\frac{\frac{\frac{\vdots f \quad \vdots g}{- \mid \Gamma \vdash A \quad S \mid \Delta_0, A, \Delta_1 \vdash C} \text{ ccut}}{S \mid \Delta_0, \Gamma, \Delta_1 \vdash C}}{\vdots \text{ sound } f \quad \vdots \text{ sound } g} \right) \doteq \frac{\frac{\llbracket - \mid \Gamma \rrbracket \Rightarrow A \quad \llbracket S \mid \Delta_0, A, \Delta_1 \rrbracket \Rightarrow C}{\llbracket S \mid \Delta_0, \Gamma, \Delta_1 \rrbracket \Rightarrow C} \text{ csbst}}$$

Theorem 4.9 *For all $f : A \Rightarrow C$, we have $\text{sound}(\text{cmplt } f) \doteq f$.*

Proof. By induction on f . □

On the other hand, cmplt is *not* a left inverse of sound up to simple equality of sequent derivations. For example, consider the derivation $\text{IL IR} : I \mid \vdash I$, since $\text{cmplt}(\text{sound}(\text{IL IR})) = \text{ax} \neq \text{IL IR}$. We therefore need to identify derivations in the sequent calculus up to the least congruence \doteq containing the following equations:

$$\frac{}{I \mid \vdash I} \text{ ax} \doteq \frac{\frac{}{- \mid I} \text{ IR}}{I \mid \vdash I} \text{ IL}$$

$$\frac{}{A \otimes B \mid \vdash A \otimes B} \text{ ax} \doteq \frac{\frac{\frac{}{B \mid \vdash B} \text{ ax}}{A \mid \vdash A} \text{ ax} \quad \frac{}{- \mid B \vdash B} \text{ shift}}{A \mid B \vdash A \otimes B} \text{ } \otimes \text{R}}{A \otimes B \mid \vdash A \otimes B} \otimes \text{L}$$

$$\frac{\frac{\frac{A' \mid \Gamma \vdash A}{- \mid A', \Gamma \vdash A} \text{ shift} \quad - \mid \Delta \vdash B}{- \mid A', \Gamma, \Delta \vdash A \otimes B} \otimes \text{R}}{\frac{- \mid \Gamma \vdash A}{I \mid \Gamma \vdash A} \text{ IL} \quad - \mid \Delta \vdash B}{I \mid \Gamma, \Delta \vdash A \otimes B} \otimes \text{R}} \otimes \text{R} \doteq \frac{\frac{A' \mid \Gamma \vdash A \quad - \mid \Delta \vdash B}{A' \mid \Gamma, \Delta \vdash A \otimes B} \otimes \text{R}}{- \mid A', \Gamma, \Delta \vdash A \otimes B} \text{ shift} \otimes \text{R}$$

$$\frac{\frac{- \mid \Gamma \vdash A}{I \mid \Gamma \vdash A} \text{ IL} \quad - \mid \Delta \vdash B}{I \mid \Gamma, \Delta \vdash A \otimes B} \otimes \text{R} \doteq \frac{- \mid \Gamma \vdash A \quad - \mid \Delta \vdash B}{- \mid \Gamma, \Delta \vdash A \otimes B} \otimes \text{R} \text{ IL}$$

$$\frac{\frac{A' \mid B', \Gamma \vdash A}{A' \otimes B' \mid \Gamma \vdash A} \otimes \text{L} \quad - \mid \Delta \vdash B}{A' \otimes B' \mid \Gamma, \Delta \vdash A \otimes B} \otimes \text{R} \doteq \frac{A' \mid B', \Gamma \vdash A \quad - \mid \Delta \vdash B}{A' \mid B', \Gamma, \Delta \vdash A \otimes B} \otimes \text{R} \otimes \text{L}$$

The inverted left rules IL^{-1} and $\otimes \text{L}^{-1}$ are compatible with the relation \doteq .

Lemma 4.10 (i) *For all $f, g : I \mid \Gamma \vdash C$, if $f \doteq g$, then $\text{IL}^{-1} f \doteq \text{IL}^{-1} g$.*

(ii) *For all $f, g : A \otimes B \mid \Gamma \vdash C$, if $f \doteq g$, then $\otimes \text{L}^{-1} f \doteq \otimes \text{L}^{-1} g$.*

We already showed in Lemma 3.2 that the rules IL^{-1} and $\otimes \text{L}^{-1}$ are left inverses of IL and $\otimes \text{L}$, respectively. They are also right inverses up to \doteq .

Lemma 4.11 *The following equations hold:*

$$\frac{\frac{\vdots f}{I \mid \Gamma \vdash C} \text{ IL}^{-1}}{\frac{- \mid \Gamma \vdash C}{I \mid \Gamma \vdash C} \text{ IL}} \doteq \frac{\vdots f}{I \mid \Gamma \vdash C}$$

$$\frac{\frac{\frac{\vdots f}{A \otimes B \mid \Gamma \vdash C} \otimes \text{L}^{-1}}{\frac{A \mid B, \Gamma \vdash C}{A \otimes B \mid \Gamma \vdash C} \otimes \text{L}} \doteq \frac{\vdots f}{A \otimes B \mid \Gamma \vdash C}$$

The function sound sends \doteq -related derivations into \doteq -related derivations.

Theorem 4.12 For all $f, g : S \mid \Gamma \vdash C$, $f \doteq g$ implies $\text{sound } f \doteq \text{sound } g$.

The function cmplt sends \doteq -related derivations into \doteq -related derivations. The proof of this fact relies on the rules scut and ccut being compatible with the relation \doteq . Moreover, it relies on the unit and associativity laws of Figure 2.

Lemma 4.13 (i) For all $f_1, f_2 : S \mid \Gamma \vdash A$ and $g_1, g_2 : A \mid \Delta \vdash C$, if $f_1 \doteq f_2$ and $g_1 \doteq g_2$, then $\text{scut } f_1 g_1 \doteq \text{scut } f_2 g_2$.

(ii) For all $f_1, f_2 : - \mid \Gamma \vdash A$ and $g_1, g_2 : S \mid \Delta_0, A, \Delta_1 \vdash C$, if $f_1 \doteq f_2$ and $g_1 \doteq g_2$, then $\text{ccut } f_1 g_1 \doteq \text{ccut } f_2 g_2$.

Theorem 4.14 For all $f, g : A \Rightarrow C$, $f \doteq g$ implies $\text{cmplt } f \doteq \text{cmplt } g$.

We showed in Theorem 4.9 that the function sound is a left inverse of cmplt up to \doteq . It is also the right inverse up to \doteq . To prove this, we first construct a rule that unpacks a stoup of the form $\llbracket S \mid \Gamma \rrbracket$ by iterating $\otimes L^{-1}$ and then also applying ll^{-1} when appropriate.

Lemma 4.15 The following rule is admissible:

$$\frac{\llbracket S \mid \Gamma \rrbracket \mid \vdash C}{S \mid \Gamma \vdash C} L^{-1}$$

Proof. We proceed by induction on Γ .

- Case Γ is empty. We define:

$$\frac{\begin{array}{c} \vdots f \\ A \mid \vdash C \end{array} L^{-1}}{A \mid \vdash C} L^{-1} =_{\text{df}} \frac{\begin{array}{c} \vdots f \\ A \mid \vdash C \end{array}}{A \mid \vdash C} \quad \frac{\begin{array}{c} \vdots f \\ \text{I} \mid \vdash C \end{array} L^{-1}}{- \mid \vdash C} L^{-1} =_{\text{df}} \frac{\begin{array}{c} \vdots f \\ \text{I} \mid \vdash C \end{array}}{- \mid \vdash C} \text{ll}^{-1}$$

- Case $\Gamma = B, \Gamma'$. We define:

$$\frac{\begin{array}{c} \vdots f \\ \llbracket A \otimes B \mid \Gamma' \rrbracket \mid \vdash C \end{array} L^{-1}}{A \mid B, \Gamma' \vdash C} L^{-1} =_{\text{df}} \frac{\begin{array}{c} \vdots f \\ \llbracket A \otimes B \mid \Gamma' \rrbracket \mid \vdash C \end{array} L^{-1}}{\frac{A \otimes B \mid \Gamma' \vdash C}{A \mid B, \Gamma' \vdash C} \otimes L^{-1}} L^{-1} \quad \frac{\begin{array}{c} \vdots f \\ \llbracket \text{I} \otimes B \mid \Gamma' \rrbracket \mid \vdash C \end{array} L^{-1}}{- \mid B, \Gamma' \vdash C} L^{-1} =_{\text{df}} \frac{\begin{array}{c} \vdots f \\ \llbracket \text{I} \otimes B \mid \Gamma' \rrbracket \mid \vdash C \end{array} L^{-1}}{\frac{\text{I} \otimes B \mid \Gamma' \vdash C}{\text{I} \mid B, \Gamma' \vdash C} \otimes L^{-1}} L^{-1} \quad \frac{\begin{array}{c} \vdots f \\ \llbracket \text{I} \otimes B \mid \Gamma' \rrbracket \mid \vdash C \end{array} L^{-1}}{- \mid B, \Gamma' \vdash C} \text{ll}^{-1}$$

□

The rule L^{-1} is compatible with the relation \doteq . This is a consequence of Lemma 4.10.

Lemma 4.16 For all $f, g : \llbracket S \mid \Gamma \rrbracket \mid \vdash C$, if $f \doteq g$, then $L^{-1} f \doteq L^{-1} g$.

Notice that the postcomposition of cmplt with sound sends a derivation of the sequent $S \mid \Gamma \vdash C$ into a derivation of the sequent $\llbracket S \mid \Gamma \rrbracket \mid \vdash C$. Therefore, for general S, Γ , a derivation $f : S \mid \Gamma \vdash C$ is incomparable with $\text{cmplt}(\text{sound } f)$. In order to fix this, we note that Lemma 4.15 strengthens our completeness result Theorem 4.5.

Corollary 4.17 (Strong completeness) For any derivation $f : \llbracket S \mid \Gamma \rrbracket \Rightarrow C$, there is a derivation $\text{strcmplt } f : S \mid \Gamma \vdash C$.

Proof. We define:

$$\text{strcmplt} \left(\frac{\begin{array}{c} \vdots f \\ \llbracket S \mid \Gamma \rrbracket \Rightarrow C \end{array}}{\llbracket S \mid \Gamma \rrbracket \Rightarrow C} \right) =_{\text{df}} \frac{\begin{array}{c} \vdots \text{cmplt } f \\ \llbracket S \mid \Gamma \rrbracket \mid \vdash C \end{array} L^{-1}}{S \mid \Gamma \vdash C} L^{-1}$$

□

Lemma 4.16 makes it an immediate consequence of Theorem 4.14 that the function strcmplt sends \doteq -related derivations into \doteq -related derivations.

Corollary 4.18 For all $f, g : \llbracket S \mid \Gamma \rrbracket \Rightarrow C$, $f \doteq g$ implies $\text{strcmplt } f \doteq \text{strcmplt } g$.

Now notice that, given $f : S \mid \Gamma \vdash C$, the derivation $\text{strcmplt}(\text{sound } f)$ also proves the sequent $S \mid \Gamma \vdash C$. We can prove that the two derivations are equal up to \doteq .

Theorem 4.19 *For all $f : S \mid \Gamma \vdash C$, we have $\text{strcmplt}(\text{sound } f) \doteq f$.*

Proof. By induction on f . □

Corollary 4.20 *For all $f : A \mid \vdash C$, we have $\text{cmplt}(\text{sound } f) \doteq f$.*

We showed in Theorem 4.9 that cmplt is a right inverse of sound up to \doteq . We can now prove that also the function strcmplt is a right inverse of sound up to \doteq . This is immediate from the following lemma.

Lemma 4.21 *The following equation holds:*

$$\text{sound} \left(\frac{\begin{array}{c} \vdots \\ f \\ \hline \llbracket S \mid \Gamma \rrbracket \mid \vdash C \\ \hline S \mid \Gamma \vdash C \end{array}}{\text{L}^{-1}} \right) \doteq \text{sound} \left(\begin{array}{c} \vdots \\ f \\ \hline \llbracket S \mid \Gamma \rrbracket \mid \vdash C \end{array} \right)$$

Corollary 4.22 *For all $f : \llbracket S \mid \Gamma \rrbracket \Rightarrow C$, we have $\text{sound}(\text{strcmplt } f) \doteq f$.*

Theorem 4.19 and Corollary 4.22 show that sound and strcmplt make a bijection between the derivations of $S \mid \Gamma \vdash C$ (considered up to \doteq) and the derivations of $\llbracket S \mid \Gamma \rrbracket \Rightarrow C$ (considered up to \doteq). (And Corollary 4.20 and Theorem 4.9 demonstrate that sound and cmplt form a bijection between the derivations of $A \mid \vdash C$ and the derivations of $A \Rightarrow C$.)

5 Focusing

If we consider the congruence relation \doteq on sequent calculus derivations as a term rewrite system (just by directing every equation to go from the left to the right), we can notice that it is weakly confluent and strongly normalizing, hence strongly confluent with unique normal forms. It turns out that these normal forms admit a simple direct description, corresponding to a natural “focused” [2] subsystem of the sequent calculus.

We present the focused subsystem as a sequent calculus with an additional *mode annotation* on sequents, which alternates between L (the “left mode”) and R (the “right mode”). In an L-sequent there is no restriction on the stoup, but in an R-sequent the stoup has to be *irreducible*, i.e., be empty or contain an atom (we write T to range over such irreducible stoups).

$$\begin{array}{c} \frac{A \mid \Gamma \vdash_L C}{- \mid A, \Gamma \vdash_L C} \text{ shift} \quad \frac{T \mid \Gamma \vdash_R C}{T \mid \Gamma \vdash_L C} \text{ msw} \quad \frac{}{X \mid \vdash_R X} \text{ ax} \\ \\ \frac{- \mid \Gamma \vdash_L C}{\mid \Gamma \vdash_L C} \text{ lL} \quad \frac{}{- \mid \vdash_R \mid} \text{ lR} \\ \\ \frac{A \mid B, \Gamma \vdash_L C}{A \otimes B \mid \Gamma \vdash_L C} \otimes_L \quad \frac{T \mid \Gamma \vdash_R A \quad - \mid \Delta \vdash_L B}{T \mid \Gamma, \Delta \vdash_R A \otimes B} \otimes_R \end{array}$$

(Note that, in the rules msw and \otimes_R , the stoup T has to be irreducible.)

As in Andreoli’s original formulation for linear logic [2], we can think of focusing as defining a root-first proof search strategy which attempts to build a derivation of a sequent bottom-up. Beginning in L-mode, the invertible rules lL and \otimes_L are applied eagerly to break down the formula in the stoup and transform it into a list of additional subformulae in the context. Once the stoup is empty, there is a choice to either apply the shift rule to shift the next formula into the stoup and repeat the inversion process, or else apply the msw rule (for “mode switch”) and move into R-mode. During R-mode, the non-invertible rule \otimes_R is applied as necessary to attempt to continue the derivation (moving back into L-mode for the right premise), while the rules lR and ax are applied to attempt to finish off the derivation.

The focused calculus is clearly sound, in the sense that, if one ignores the mode annotations, all of the above rules are either rules of the original sequent calculus or else (in the case of msw) have conclusion equal to their premise. In fact, the focused calculus is complete, and indeed optimal in the sense that focused derivations give unique representatives of each \doteq -equivalence class. This is established by the following results.

Proposition 5.1 *For any derivation $f : S \mid \Gamma \vdash_L C$, there is a derivation $\text{emb } f : S \mid \Gamma \vdash C$.*

Theorem 5.2 *For any derivation $f : S \mid \Gamma \vdash C$, there is a derivation $\text{focus } f : S \mid \Gamma \vdash_{\mathbb{L}} C$.*

Proof. By induction on f . □

Theorem 5.3 *For any $f, g : S \mid \Gamma \vdash_{\mathbb{L}} C$, $\text{focus}(\text{emb } f) = f$.*

Proof. By induction on f . □

Theorem 5.4 *For any $f, g : S \mid \Gamma \vdash C$, if $f \doteq g$, then $\text{focus } f = \text{focus } g$.*

Proof. By induction on the proof of $f \doteq g$. □

Theorem 5.5 *For any $f : S \mid \Gamma \vdash C$, we have $\text{emb}(\text{focus } f) \doteq f$.*

Proof. By induction on f . □

Corollary 5.6 *For any $f, g : S \mid \Gamma \vdash C$, if $\text{focus } f = \text{focus } g$, then $f \doteq g$.*

Putting these results together with Theorems 4.4, 4.5 on soundness and completeness and Theorems 4.9, 4.12, 4.14, we obtain a simple decision procedure for equality of maps in the free skew monoidal category.

Corollary 5.7 *For any $f, g : A \Rightarrow C$, we have $f \doteq g$ if and only if $\text{focus}(\text{cmplt } f) = \text{focus}(\text{cmplt } g)$.*

It was already observed that the cut-free sequent calculus of Section 3 can be used to decide existence of maps in the free skew monoidal category. On the other hand, the focused calculus yields a more efficient decision procedure, since there is less non-determinism. Indeed, it provides a simple algorithm for listing the elements of any homset in the free skew monoidal category, without duplicates.

Theorem 5.8 *For any S, Γ, C , one can compute a duplicate-free list $\text{focderivs}(S, \Gamma, C)$ of derivations of $S \mid \Gamma \vdash_{\mathbb{L}} C$ containing every such derivation. In particular, we can decide whether $S \mid \Gamma \vdash_{\mathbb{L}} C$ is derivable.*

Proof. As explained above, we can consider the focused calculus as a root-first search strategy. This search is guaranteed to terminate because, for any goal sequent $S \mid \Gamma \vdash_m C$ ($m \in \{\mathbb{L}, \mathbb{R}\}$), there are only finitely many possible instances of rules to apply, and the subgoals that they generate are always smaller relative to a well-founded order on sequents. (We can rank sequents by lexicographically ordered triples consisting of the number of occurrences of \mathbb{I} and \otimes , the information whether the stoup is empty or not, with singleton $<$ empty, and the mode, with $\mathbb{R} < \mathbb{L}$.) □

Corollary 5.9 *For any A, C , one can compute a duplicate-free list of derivations $A \Rightarrow C$ that contains, for any derivation $f : A \Rightarrow C$ exactly one derivation g such that $f \doteq g$, as $(\text{sound} \circ \text{emb})^*(\text{focderivs}(A, (), C))$.*

It is worth mentioning that there are some surprisingly elegant formulae for *counting* different families of maps in the free skew semigroup category [7], and so it may be interesting to apply the focused sequent calculus to pursue a similar quantitative analysis of maps in the free skew monoidal category (cf. [18]).

6 Conclusion and Future Work

In this paper, we studied the free skew monoidal category from a proof-theorist’s point-of-view. We considered three calculi, a categorical calculus, which embodies the definition of the free skew monoidal category, a sequent calculus and a focused version thereof, and proved their sets of derivations to be in bijections. Nicely, in the focused calculus, there is no need to quotient derivations by a congruence relation, equality of derivations is just (literal) equality. The focused calculus thus provides a very concrete description of the free skew monoidal category. We learned that, although simply defined, skew monoidal categories are remarkably subtle. We also learned that methods of proof theory, such as proof techniques of cut admissibility, focusing or use of non-standard sequent forms, are surprisingly well suited for exploring them.

We envisage a number of directions for future work.

We linked the categorical calculus to the focused sequent calculus via the unrestricted sequent calculus. That the focused calculus is complete with respect to the categorical calculus can also be proved directly. The resulting focusing completeness proof can be shown to be equal to the post-composition of the unrestricted completeness proof with focusing. Whether this leads to a shorter development overall remains to be checked.

We hinted at a connection to multicategories, but this needs to be worked out properly. Under reasonable suitable definitions, we would expect the focused sequent calculus to give an explicit construction of the free *left representable skew multicategory* [5] over a set of atoms.

In the free skew monoidal category, there can be multiple maps between the same two objects, i.e., multiple focused derivations of the same sequent. We believe that we can partially order the derivations in a canonical way both in the categorical calculus and in the sequent calculus. In particular, we can have a greatest element,

i.e., a preferred derivation for any derivable sequent, and have soundness and completeness preserve these partial orders. We say some words about this plan in Appendix B.

A natural direction for continuing this work would be to consider the sequent calculus of skew closed categories and/or skew monoidal closed categories (corresponding to restrictions of Lambek’s original calculus with only one implication, with or without the tensor product). Finally, given the connections between the Tamari order and the higher-dimensional polytopes known as *associahedra* [15], it may be interesting to develop sequent calculi for higher-dimensional skew monoidal or skew semigroup categories.

Acknowledgements

T.U. was partially supported by the Estonian Ministry of Education and Research institutional research grant no. IUT33-13. N.V. was supported by the ERDF funded Estonian national CoE project EXCITE and a research grant (13156) from VILLUM FONDEN.

References

- [1] T. Altenkirch, J. Chapman, T. Uustalu (2015): *Monads need not be endofunctors*. *Log. Methods in Comput. Sci.* 11(1), article 3, 2015. doi: [10.2168/lmcs-11\(1:3\)](https://doi.org/10.2168/lmcs-11(1:3))
- [2] J.-M. Andreoli (1992): *Logic programming with focusing proofs in linear logic*. *J. of Log. and Comput.* 2(3), pp. 297–347. doi: [10.1093/logcom/2.3.297](https://doi.org/10.1093/logcom/2.3.297)
- [3] J. Bénabou (1963), *Catégories avec multiplication*. *C. R. Acad. Sci. Paris* 256, pp. 1887–1890. Available at <http://gallica.bnf.fr/ark:/12148/bpt6k3208j/f1965.image>.
- [4] J. Bourke, S. Lack (2017): *Free skew monoidal categories*. arXiv preprint 1708.06087. Available at <https://arxiv.org/abs/1708.06087>.
- [5] J. Bourke, S. Lack (2017): *Skew monoidal categories and skew multicategories*. arXiv preprint 1708.06087. Available at <https://arxiv.org/abs/1708.06087>.
- [6] M. Buckley, R. Garner, S. Lack, R. Street (2014): *The Catalan simplicial set*. *Math. Proc. of Cambridge Philos. Soc.* 158(12), pp. 211–222. doi: [10.1017/s0305004114000498](https://doi.org/10.1017/s0305004114000498)
- [7] F. Chapoton (2006): *Sur le nombre d’intervalles dans les treillis de Tamari*. *Séminaire Lotharingien de Combinatoire* 55, article B55f. <https://www.mat.univie.ac.at/~slc/wpapers/s55chapoton.html>
- [8] G. M. Kelly (1964): *On MacLane’s conditions for coherence of natural associativities, commutativities, etc.* *J. of Alg.* 1(4), pp. 397–402, doi: [10.1016/0021-8693\(64\)90018-3](https://doi.org/10.1016/0021-8693(64)90018-3).
- [9] S. Lack, R. Street (2012): *Skew monoidales, skew warpings and quantum categories*. *Theor. and Appl. of Categ.* 26, pp. 385–402. Available at <http://www.tac.mta.ca/tac/volumes/26/15/26-15abs.html>
- [10] S. Lack, R. Street (2014): *Triangulations, orientals, and skew monoidal categories*. *Adv. in Math.* 258, pp. 351–396, 2014. doi: [10.1016/j.aim.2014.03.003](https://doi.org/10.1016/j.aim.2014.03.003)
- [11] J. Lambek (1958): *The mathematics of sentence structure*. *Amer. Math. Monthly* 65(3), pp. 154–170. doi: [10.2307/2310058](https://doi.org/10.2307/2310058)
- [12] T. Leinster (2004): *Higher Operads, Higher Categories*, v. 298 of *London Math. Soc. Lect. Note Ser.* Cambridge Univ. Press. doi: [10.1017/cbo9780511525896](https://doi.org/10.1017/cbo9780511525896) (Preprint version available at <https://arxiv.org/abs/math/0305049>.)
- [13] S. Mac Lane (1963): *Natural associativity and commutativity*. *Rice Univ. Stud.* 49(4), pp. 28–46. Available at <http://hdl.handle.net/1911/62865>.
- [14] S. Mac Lane (1971): *Categories for the Working Mathematician*. Springer-Verlag. doi: [10.1007/978-1-4757-4721-8](https://doi.org/10.1007/978-1-4757-4721-8)
- [15] F. Müller-Hoissen, J.-M. Pallo, J. Stasheff (eds.) (2012): *Associahedra, Tamari Lattices and Related Structures: Tamari Memorial Festschrift*, v. 299 of *Progress in Mathematics*. Birkhäuser. doi: [10.1007/978-3-0348-0405-9](https://doi.org/10.1007/978-3-0348-0405-9)
- [16] K. Szlachányi (2012): *Skew-monoidal categories and bialgebroids*. *Adv. in Math.* 231(3–4), pp. 1694–1730, doi: [10.1016/j.aim.2012.06.027](https://doi.org/10.1016/j.aim.2012.06.027).
- [17] T. Uustalu (2014): *Coherence for skew-monoidal categories*. In P. Levy, N. Krishnaswami (eds.), *Proc. of 5th Wksh. on Mathematically Structured Programming, MSFP 2014*, v. 153 of *Electron. Proc. in Theor. Comput. Sci.*, pp. 68–77. Open Publishing Assoc. doi: [10.4204/eptcs.153.5](https://doi.org/10.4204/eptcs.153.5)
- [18] N. Zeilberger (2017): *A sequent calculus for a semi-associative law*. In D. Miller (ed.), *Proc. of 2nd Int. Conf. on Formal Structures for Computation and Deduction, FSCD 2017*, v. 84 of *Leibniz Int. Proc. in Inform.*, article 33. Dagstuhl Publishing. doi: [10.4230/lipics.fscd.2017.33](https://doi.org/10.4230/lipics.fscd.2017.33)

A Sequent Calculus as a Generalized Multicategory

The sequent calculus can be seen as a particular generalized multicategory for a Cartesian monad that we denote $(\cdot)^{?*}$. Its underlying functor is given by $X^{?*} =_{\text{df}} X^? \times X^*$ where $X^?$ is the free pointed set on X (obtained by adding an element to X) and X^* is the free monoid on X (the set of lists over X). For elements of $X^{?*}$, we use the notation we have been using for antecedents, i.e., we write $S \mid \Gamma$ for an element of TX , where S can either be nothing or an element of X and Γ is a list of elements of X . The unit $\eta_X^{?*} : X \rightarrow X^{?*}$ of $(\cdot)^{?*}$ is given by $\eta^{?*} A =_{\text{df}} A \mid \cdot$. The multiplication $\mu_X^{?*} : (X^{?*})^{?*} \rightarrow X^{?*}$ is given by:

$$\mu^{?*}(- \mid \Delta) =_{\text{df}} - \mid \mu^*([\cdot]^* \Delta) \qquad \mu^{?*}((S \mid \Gamma) \mid \Delta) =_{\text{df}} S \mid \Gamma, \mu^*([\cdot]^* \Delta)$$

where μ^* is the multiplication of the list monad $(\cdot)^*$, i.e., concatenation of lists, and $[\cdot]$ is a monad morphism typed $[\cdot]_X : X^{?*} \rightarrow X^*$ and given by $[- \mid \Gamma] =_{\text{df}} \Gamma$ and $[A \mid \Gamma] =_{\text{df}} A, \Gamma$.

The sequent calculus is an instance of a generalized multicategory for $(\cdot)^{?*}$. We call this generalized multicategory **SC**. An object of **SC** is a formula, i.e., an element of Fma . A map between $S \mid \Gamma : \text{Fma}^{?*}$ and $C : \text{Fma}$ is a derivations of the sequent $S \mid \Gamma \vdash C$.

The identity map on an object A is the derivation ax of the sequent $A \mid \vdash A$.

The composition of the generalized multicategory is given by the following two ‘‘multicut’’ rules, defined from shift , scut and ccut .

$$\frac{S \mid \Gamma \vdash B \quad S_1 \mid \Gamma_1 \vdash B_1 \quad \dots \quad S_n \mid \Gamma_n \vdash B_n \quad B \mid B_1, \dots, B_n \vdash C}{S \mid \Gamma, S_1, \Gamma_1, \dots, S_n, \Gamma_n \vdash C} \text{mcut}^j$$

$$\frac{S_1 \mid \Gamma_1 \vdash B_1 \quad \dots \quad S_n \mid \Gamma_n \vdash B_n \quad - \mid B_1, \dots, B_n \vdash C}{- \mid \Gamma, S_1, \Gamma_1, \dots, S_n, \Gamma_n \vdash C} \text{mcut}^n$$

In the conclusions of these rules, we are informal in our notation, instead of S_i, Γ_i , we should officially write $[S_i \mid \Gamma_i]$.

The construction is as follows. First, we define a generalized version of shift :

$$\frac{S \mid \Gamma \vdash C}{- \mid S, \Gamma \vdash C} \text{shift}^?$$

The definition is immediate:

$$\frac{\begin{array}{c} \vdots \\ f \\ A \mid \Gamma \vdash C \end{array}}{- \mid A, \Gamma \vdash C} \text{shift}^? =_{\text{df}} \frac{\begin{array}{c} \vdots \\ f \\ A \mid \Gamma \vdash C \end{array}}{- \mid A, \Gamma \vdash C} \text{shift} =_{\text{df}} \frac{\begin{array}{c} \vdots \\ f \\ - \mid \Gamma \vdash C \end{array}}{- \mid \Gamma \vdash C} \text{shift}^? =_{\text{df}} \frac{\begin{array}{c} \vdots \\ f \\ - \mid \Gamma \vdash C \end{array}}{- \mid \Gamma \vdash C}$$

Using $\text{shift}^?$, the rules mcut^j and mcut^n are defined as follows:

$$\begin{array}{c}
 \begin{array}{c} \vdots f \\ S | \Gamma \vdash B \end{array} \quad \begin{array}{c} \vdots f_1 \\ S_1 | \Gamma_1 \vdash B_1 \end{array} \quad \dots \quad \begin{array}{c} \vdots f_n \\ S_n | \Gamma_n \vdash B_n \end{array} \quad \begin{array}{c} \vdots g \\ B | B_1, \dots, B_n \vdash C \end{array} \\
 \hline
 S | \Gamma, S_1, \Gamma_1, \dots, S_n, \Gamma_n \vdash C \quad \text{mcut}^j
 \end{array}$$

$$\begin{array}{c}
 \begin{array}{c} \vdots f \\ S | \Gamma \vdash B \end{array} \\
 =_{\text{df}} \quad \frac{\begin{array}{c} \vdots f_1 \\ S_1 | \Gamma_1 \vdash B_1 \end{array} \quad \frac{\begin{array}{c} \vdots f_n \\ S_n | \Gamma_n \vdash B_n \end{array} \quad \frac{\begin{array}{c} \vdots g \\ B | B_1, \dots, B_n \vdash C \end{array}}{\begin{array}{c} - | S_n, \Gamma_n, \vdash B_n \end{array} \text{shift}^?} \quad \begin{array}{c} \vdots g \\ B | B_1, \dots, B_n \vdash C \end{array}}{\begin{array}{c} B | B_1, \dots, B_{n-1}, S_n, \Gamma_n \vdash C \end{array}} \text{ccut}}{\begin{array}{c} \vdots f \\ - | S_1, \Gamma_1, \vdash B_1 \end{array} \text{shift}^? \quad \begin{array}{c} \vdots \\ B | B_1, S_2, \Gamma_2, \dots, S_n, \Gamma_n \vdash C \end{array}} \text{ccut}} \\
 \hline
 S | \Gamma, S_1, \Gamma_1, \dots, S_n, \Gamma_n \vdash C \quad \text{scut}
 \end{array}$$

$$\begin{array}{c}
 \begin{array}{c} \vdots f_1 \\ S_1 | \Gamma_1 \vdash B_1 \end{array} \quad \dots \quad \begin{array}{c} \vdots f_n \\ S_n | \Gamma_n \vdash B_n \end{array} \quad \begin{array}{c} \vdots g \\ - | B_1, \dots, B_n \vdash C \end{array} \\
 \hline
 - | S_1, \Gamma_1, \dots, S_n, \Gamma_n \vdash C \quad \text{mcut}^n
 \end{array}$$

$$\begin{array}{c}
 \begin{array}{c} \vdots f_1 \\ S_1 | \Gamma_1 \vdash B_1 \end{array} \\
 =_{\text{df}} \quad \frac{\begin{array}{c} \vdots f_n \\ S_n | \Gamma_n \vdash B_n \end{array} \quad \frac{\begin{array}{c} \vdots g \\ - | B_1, \dots, B_n \vdash C \end{array}}{\begin{array}{c} - | S_n, \Gamma_n, \vdash B_n \end{array} \text{shift}^?} \quad \begin{array}{c} \vdots g \\ - | B_1, \dots, B_n \vdash C \end{array}}{\begin{array}{c} - | B_1, \dots, B_{n-1}, S_n, \Gamma_n \vdash C \end{array}} \text{ccut}}{\begin{array}{c} \vdots f_1 \\ - | S_1, \Gamma_1, \vdash B_1 \end{array} \text{shift}^? \quad \begin{array}{c} \vdots \\ - | B_1, S_2, \Gamma_2, \dots, S_n, \Gamma_n \vdash C \end{array}} \text{ccut}} \\
 \hline
 - | S_1, \Gamma_1, \dots, S_n, \Gamma_n \vdash C \quad \text{ccut}
 \end{array}$$

On the other hand, we could start with mcut^j and mcut^n and define scut and ccut as follows:

$$\begin{array}{c}
 \begin{array}{c} \vdots f \\ S | \Gamma \vdash B \end{array} \quad \begin{array}{c} \vdots g \\ B | B_1, \dots, B_n \vdash C \end{array} \\
 \hline
 S | \Gamma, B_1, \dots, B_n \vdash C \quad \text{scut}
 \end{array}
 =_{\text{df}}
 \begin{array}{c}
 \begin{array}{c} \vdots f \\ S | \Gamma \vdash B \end{array} \quad \frac{\begin{array}{c} \vdots g \\ B_1 | \vdash B_1 \end{array} \text{ax}}{\begin{array}{c} \vdots \\ B_1 | \vdash B_1 \end{array}} \quad \dots \quad \frac{\begin{array}{c} \vdots g \\ B_n | \vdash B_n \end{array} \text{ax}}{\begin{array}{c} \vdots \\ B_n | \vdash B_n \end{array}} \quad \begin{array}{c} \vdots g \\ B | B_1, \dots, B_n \vdash C \end{array} \\
 \hline
 S | \Gamma, B_1, \dots, B_n \vdash C \quad \text{mcut}^j
 \end{array}$$

$$\begin{array}{c}
 \begin{array}{c} \vdots f \\ - | \Gamma \vdash B \end{array} \quad \begin{array}{c} \vdots g \\ A | B_1, \dots, B_i, B, B_{i+1}, \dots, B_n \vdash C \end{array} \\
 \hline
 A | B_1, \dots, B_i, \Gamma, B_{i+1}, \dots, B_n \vdash C \quad \text{ccut}
 \end{array}$$

$$\begin{array}{c}
 \begin{array}{c} \vdots f \\ A | \vdash A \end{array} \text{ax} \quad \frac{\begin{array}{c} \vdots g \\ B_1 | \vdash B_1 \end{array} \text{ax}}{\begin{array}{c} \vdots \\ B_1 | \vdash B_1 \end{array}} \quad \dots \quad \frac{\begin{array}{c} \vdots f \\ - | \Gamma \vdash B \end{array}}{\begin{array}{c} \vdots \\ - | \Gamma \vdash B \end{array}} \quad \dots \quad \frac{\begin{array}{c} \vdots g \\ B_n | \vdash B_n \end{array} \text{ax}}{\begin{array}{c} \vdots \\ B_n | \vdash B_n \end{array}} \quad \begin{array}{c} \vdots g \\ A | B_1, \dots, B_i, B, B_{i+1}, \dots, B_n \vdash C \end{array} \\
 \hline
 A | B_1, \dots, B_i, \Gamma, B_{i+1}, \dots, B_n \vdash C \quad \text{mcut}^j
 \end{array}$$

$$\begin{array}{c}
 \begin{array}{c} \vdots f \\ - | \Gamma \vdash B \end{array} \quad \begin{array}{c} \vdots g \\ - | B_1, \dots, B_i, B, B_{i+1}, \dots, B_n \vdash C \end{array} \\
 \hline
 - | B_1, \dots, B_i, \Gamma, B_{i+1}, \dots, B_n \vdash C \quad \text{ccut}
 \end{array}$$

$$\begin{array}{c}
 \begin{array}{c} \vdots f \\ B_1 | \vdash B_1 \end{array} \text{ax} \quad \dots \quad \frac{\begin{array}{c} \vdots f \\ - | \Gamma \vdash B \end{array}}{\begin{array}{c} \vdots \\ - | \Gamma \vdash B \end{array}} \quad \dots \quad \frac{\begin{array}{c} \vdots g \\ B_n | \vdash B_n \end{array} \text{ax}}{\begin{array}{c} \vdots \\ B_n | \vdash B_n \end{array}} \quad \begin{array}{c} \vdots g \\ - | B_1, \dots, B_i, B, B_{i+1}, \dots, B_n \vdash C \end{array} \\
 \hline
 - | B_1, \dots, B_i, \Gamma, B_{i+1}, \dots, B_n \vdash C \quad \text{mcut}^n
 \end{array}$$

There is also another multicategory in the picture, this one standard, i.e., based on the list Cartesian monad $(\cdot)^*$. We call it \mathbf{SC}^n . An object of \mathbf{SC}^n is again a formula, i.e., an element of \mathbf{Fma}^* . A map between $\Gamma : \mathbf{Fma}^*$ and $C : \mathbf{Fma}^*$ is a derivation of the sequent $- | \Gamma \vdash C$. The identity on A is the derivation $\text{shift ax} : - | A \vdash A$. Composition is given by the restriction of mcut^n to the case where the stoups S_1, \dots, S_n are all $-$.

$\text{shift}^?$ defines an identity-on-objects functor from \mathbf{SC} to \mathbf{SC}^n wrt. the monad morphism $[\cdot]$ between the corresponding Cartesian monads.

B Not All Derivations Are Equal

As we discussed, the focused sequent calculus is almost deterministic for root-first proof search. Specifically, there are exactly two types of choice situations. First, in the L-mode, when the stoup is empty and the context contains at least one formula, both the rules `shift` and `msw` can be applied and sometimes both can lead to a derivation. Second, in the R-mode, when applying the \otimes R rule, one has to choose how to split the context and again sometimes several splits can lead to a derivation. In the first case, the choice is only interesting, if the succedent formula is of the form $A \otimes B$ and the `msw` inference is immediately followed by a \otimes R inference (if it is a propositional letter or `!`, then the attempt with `msw` is bound to fail). A small analysis shows that, if both avenues lead to a derivation, then, modulo \cong , the underlying choice is still between two different splittings of the context in an \otimes R inference.

Now, in the unfocused calculus one can show admissible the following rules where J is any formula constructed with `!` and \otimes only (a closed formula):

$$\frac{S \mid \Gamma, J \vdash C}{S \mid \Gamma \vdash C} \text{ b} \qquad \frac{- \mid \Delta \vdash C}{- \mid J, \Delta \vdash C} \#$$

This suggests the following inequation for derivations:

$$\frac{\frac{\vdots f}{S \mid \Gamma, J \vdash A} \quad \frac{\vdots g}{- \mid \Delta \vdash B}}{S \mid \Gamma, J, \Delta \vdash A \otimes B} \circ \geq \frac{\frac{\vdots f}{S \mid \Gamma \vdash A} \quad \frac{\vdots g}{- \mid \Delta \vdash B}}{\frac{S \mid \Gamma \vdash A}{- \mid J, \Delta \vdash B} \text{ b}} \# \geq \frac{S \mid \Gamma, J, \Delta \vdash A \otimes B}{S \mid \Gamma, J, \Delta \vdash A \otimes B}$$

The intuition here is that since the sequence of atoms in the antecedent and succedent of a derivable sequent (bracketing ignored, occurrences of `!` dropped) must coincide, the only actual non-determinism in choosing how to split an antecedent arises when there are formulae in the middle of the context that consist only of `!`'s. Such formulae can potentially be useful both in the first and the second half of the split. Whenever a formula J like this leads to a derivation if put into the first half, then we also have a derivation if we put J in the second half. The converse is not true. A derivation may exist for the placement of J into the second half while there is no derivation with J in the first half.

We conjecture that the partial order on sequent calculus derivations generated by this inequality corresponds to the partial order on categorical calculus derivations generated by the following inequations corresponding to alternative versions of Mac Lane's equations (a) and (b):

$$\begin{aligned} \frac{\frac{\overline{I \Rightarrow I \otimes I} \quad \rho \quad \overline{I \otimes I \Rightarrow I} \quad \lambda}{I \otimes I \Rightarrow I \otimes I} \text{ comp}}{\overline{I \otimes I \Rightarrow I \otimes I}} &\geq \overline{I \otimes I \Rightarrow I \otimes I} \text{ ax} \\ \overline{X \otimes (I \otimes Y) \Rightarrow X \otimes (I \otimes Y)} \text{ ax} &\geq \frac{\frac{\overline{X \Rightarrow X \otimes I} \quad \rho \quad \overline{I \otimes Y \Rightarrow Y} \quad \lambda}{X \otimes (I \otimes Y) \Rightarrow (X \otimes I) \otimes Y} \otimes \quad \frac{\overline{(X \otimes I) \otimes Y \Rightarrow X \otimes (I \otimes Y)}}{(X \otimes I) \otimes Y \Rightarrow X \otimes (I \otimes Y)} \alpha}{X \otimes (I \otimes Y) \Rightarrow X \otimes (I \otimes Y)} \text{ comp} \\ \overline{(X \otimes I) \otimes Y \Rightarrow (X \otimes I) \otimes Y} \text{ ax} &\geq \frac{\overline{(X \otimes I) \otimes Y \Rightarrow (X \otimes I) \otimes Y} \quad \alpha \quad \frac{\overline{X \Rightarrow X \otimes I} \quad \rho \quad \overline{I \otimes Y \Rightarrow Y} \quad \lambda}{X \otimes (I \otimes Y) \Rightarrow (X \otimes I) \otimes Y} \otimes}{(X \otimes I) \otimes Y \Rightarrow (X \otimes I) \otimes Y} \text{ comp} \end{aligned}$$

More precisely, we know that `cmplt` preserves \geq . We have proved some cases of `sound` preserving $\overset{\circ}{\geq}$. This is ongoing work.