

Optimal Two-Qubit Circuits for Universal Fault-Tolerant Quantum Computation

Andrew N. Glaudell,^{1,2,3,*} Neil J. Ross,⁴ and Jacob M. Taylor^{1,2}

¹ Institute for Advanced Computer Studies and Joint Center for Quantum Information and Computer Science,
University of Maryland, College Park, MD, USA

² Joint Quantum Institute, University of Maryland, College Park, MD, USA

³ Booz Allen Hamilton BASIQ Research Team, Washington, DC, USA

⁴ Department of Mathematics and Statistics, Dalhousie University, Halifax, NS, Canada

Abstract

We study two-qubit circuits over the Clifford+ CS gate set, which consists of the Clifford gates together with the controlled-phase gate $CS = \text{diag}(1, 1, 1, i)$. The Clifford+ CS gate set is universal for quantum computation and its elements can be implemented fault-tolerantly in most error-correcting schemes through magic state distillation. Since non-Clifford gates are typically more expensive to perform in a fault-tolerant manner, it is often desirable to construct circuits that use few CS gates. In the present paper, we introduce an efficient and optimal synthesis algorithm for two-qubit Clifford+ CS operators. Our algorithm inputs a Clifford+ CS operator U and outputs a Clifford+ CS circuit for U , which uses the least possible number of CS gates. Because the algorithm is deterministic, the circuit it associates to a Clifford+ CS operator can be viewed as a normal form for that operator. We give an explicit description of these normal forms and use this description to derive a worst-case lower bound of $5 \log_2(\frac{1}{\epsilon}) + O(1)$ on the number of CS gates required to ϵ -approximate elements of $SU(4)$. Our work leverages a wide variety of mathematical tools that may find further applications in the study of fault-tolerant quantum circuits.

1 Introduction

In the context of fault-tolerant quantum computing, operations from the Clifford group are relatively easy to perform and are therefore considered inexpensive. In contrast, operations that do not belong to the Clifford group are complicated to execute fault-tolerantly because they require resource intensive distillation protocols [29]. Since non-Clifford operations are necessary for universal quantum computing, it has become standard to use the number of non-Clifford gates in a circuit as a measure of its cost. This fault-tolerant perspective on the cost of circuits has profoundly impacted the field of quantum compiling and significant efforts have been devoted to minimizing the number of non-Clifford operations in circuits.

An important problem in quantum compiling is the problem of *exact synthesis*: given an operator U known to be exactly representable over some gate set G , find a circuit for U over G . An *exact synthesis algorithm* is a constructive solution to this problem. When the gate set G is an extension of the Clifford group, it is desirable that the exact synthesis algorithm for G be efficient and produce circuits that use as few non-Clifford gates as possible.

In the past few years, methods from algebraic number theory have been successfully applied to the exact synthesis problem associated to a variety of single-qubit [4, 6, 9, 22, 24, 30, 31] and single-qutrit [5, 15, 21, 28] gate sets. In many cases, the resulting exact synthesis algorithms efficiently produce circuits that are *optimal*, in the sense that they use the least possible number of non-Clifford gates. These powerful exact synthesis

*Corresponding author: Glaudell_Andrew@bah.com

methods were central in the development of good unitary approximation methods, which play a key role in the compilation of practical quantum programs [4, 6, 20, 23, 30, 31].

Exact synthesis algorithms also exist for various instantiations of the multi-qubit compiling problem, though each suffers shortcomings in some respect. Optimal algorithms for two-qubit circuits over continuous gate sets have been known for a number of years [34, 38]. Unfortunately, such gate sets are not well-suited for fault-tolerant quantum computing. Multi-qubit exact synthesis algorithms for universal and fault-tolerant gate sets were introduced more recently [1, 2, 13, 17, 19, 25, 26, 27]. While the algorithms of [1, 13] are far from optimal, the algorithms of [2, 17, 25, 27] synthesize provably optimal circuits by cleverly utilizing certain properties of fault-tolerant gate sets containing the Clifford group. However, the runtimes of these optimal synthesis algorithms are exponential in both qubit count and optimal circuit length. Powerful heuristics were introduced in [27] achieving polynomial scaling with optimal circuit length. Unfortunately, even this improved heuristic algorithm takes thousands of seconds to compute optimal two-qubit circuits of practical size (40 non-Clifford operations) on modest hardware.

Not only are these multi-qubit exact synthesis algorithms impractical in many cases, they also fail to shed much light on the *structure* of optimal circuits. In the single-qubit case, intimate knowledge of this structure for certain gate sets was developed by describing optimal circuits via regular expressions or, equivalently, automata [14]. Such descriptions are of theoretical interest, but also have practical consequences. In particular, for certain single-qubit gate sets these descriptions allowed researchers to derive a rigorous lower-bound on the number of non-Clifford gates required to approximate typical elements of $SU(2)$ [32]. Analogous statements about approximations of multi-qubit unitaries have eluded researchers thus far.

In the present paper, we introduce an efficient and optimal exact synthesis algorithm for a two-qubit gate set that is appropriate for universal and fault-tolerant quantum computing. We focus on two-qubit circuits over the Clifford+ CS gate set, which consists of the Clifford gates together with the non-Clifford controlled-phase gate $CS = \text{diag}(1, 1, 1, i)$. The CS gate has received recent attention as an alternative to the T -gate in methods for fault-tolerant quantum computing [3, 18] and due to its natural implementation as an entangling operation in certain superconducting qubit systems [8, 11, 12, 33] whose fidelity is approaching that of single-qubit gates [10, 37]. Our algorithm produces an optimal circuit in a number of arithmetic operations linear in the length of the optimal decomposition. This is unlike existing multi-qubit synthesis methods. Moreover, because our algorithm is deterministic, the circuit it associates to a Clifford+ CS operator can be viewed as a normal form for that operator. We give an explicit description of these normal forms in the language of automata and use this description to derive a worst-case lower bound of $5 \log_2(\frac{1}{\epsilon}) + O(1)$ on the number of CS gates required to ϵ -approximate elements of $SU(4)$. A Mathematica package implementing our algorithm is freely available on-line [16]. This code is very efficient, synthesizing optimal circuits of CS -count 10000 in 1.2 ± 0.1 seconds on modest hardware.

The paper is structured as follows. We first introduce a convenient set of generators in [Section 2](#). Then, in [Section 3](#), we describe the exceptional isomorphism $SU(4) \cong \text{Spin}(6)$. In [Section 4](#), we leverage this isomorphism to introduce an exact synthesis algorithm for Clifford+ CS operators. In [Section 5](#), we use the theory of automata to study the structure of the circuits produced by the exact synthesis algorithm. We take advantage of this structure in [Section 6](#) to establish a worst-case lower bound on the number of non-Clifford resources required to ϵ -approximate elements of $SU(4)$ using Clifford+ CS circuits. Finally, we conclude and discuss avenues for future work in [Section 7](#).

2 Generators

Throughout, we use \mathbb{N} , \mathbb{Z} , \mathbb{R} , and \mathbb{C} to denote the usual collection of numbers, \mathbb{Z}_p to denote the collection integers modulo p , and $\mathbb{Z}[i]$ to denote the collection of Gaussian integers (the complex numbers with integer real and imaginary parts). We write ρ for the canonical homomorphism $\mathbb{Z} \rightarrow \mathbb{Z}_2$ (if $n \in \mathbb{Z}$ then $\rho(n)$ is the parity of n). For two integers $n \leq m$, we write $[n, m]$ for the set $\{n, \dots, m\} \subseteq \mathbb{Z}$ and simply write $[m]$ for $[1, m]$. We view scalars and vectors as matrices so that any concept defined for matrices of arbitrary dimensions also applies to scalars and vectors. Finally, for readability, we use the symbol \cdot to denote the zero entries of a matrix.

The single-qubit *Pauli* gates X , Y , and Z are defined as

$$X = \begin{bmatrix} \cdot & 1 \\ 1 & \cdot \end{bmatrix}, \quad Y = \begin{bmatrix} \cdot & -i \\ i & \cdot \end{bmatrix}, \quad \text{and} \quad Z = \begin{bmatrix} 1 & \cdot \\ \cdot & -1 \end{bmatrix}.$$

These gates generate the *single-qubit Pauli group* $\{i^a P ; a \in \mathbb{Z}_4 \text{ and } P \in \{I, X, Y, Z\}\}$. The *two-qubit Pauli group*, which we denote by \mathcal{P} , is defined as $\mathcal{P} = \{i^a (P \otimes Q) ; a \in \mathbb{Z}_4 \text{ and } P, Q \in \{I, X, Y, Z\}\}$. The *Clifford* gates H , S , and CZ are defined as

$$H = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix}, \quad S = \begin{bmatrix} 1 & \cdot \\ \cdot & i \end{bmatrix}, \quad \text{and} \quad CZ = \begin{bmatrix} 1 & \cdot & \cdot & \cdot \\ \cdot & 1 & \cdot & \cdot \\ \cdot & \cdot & 1 & \cdot \\ \cdot & \cdot & \cdot & -1 \end{bmatrix}.$$

These gates are known as the *Hadamard* gate, the *phase* gate, and the *controlled-Z* gate, respectively. The *single-qubit Clifford group* is generated by H and S and contains the primitive 8-th root of unity $\omega = e^{\frac{i\pi}{4}}$. The *two-qubit Clifford group*, which we denote by \mathcal{C} , consists of the operators which can be represented by a two-qubit circuit over the gate set $\{H, S, CZ\}$. Equivalently, \mathcal{C} is generated by $H \otimes I$, $I \otimes H$, $S \otimes I$, $I \otimes S$, and CZ . Up to global phases, the Clifford groups are the normalizers of the Pauli groups.

Clifford gates are well-suited for fault-tolerant quantum computation but the Clifford group is not universal. One can obtain a universal group by extending \mathcal{C} with the *controlled-phase gate* CS defined as

$$CS = \begin{bmatrix} 1 & \cdot & \cdot & \cdot \\ \cdot & 1 & \cdot & \cdot \\ \cdot & \cdot & 1 & \cdot \\ \cdot & \cdot & \cdot & i \end{bmatrix}.$$

In what follows, we focus on the group \mathcal{G} of operators which can be represented by a two-qubit circuit over the universal gate set $\{H, S, CZ, CS\}$. Equivalently, \mathcal{G} is the group generated by $H \otimes I$, $I \otimes H$, $S \otimes I$, $I \otimes S$, CZ , and CS . We have $\mathcal{P} \subseteq \mathcal{C} \subseteq \mathcal{G}$. We sometimes refer to \mathcal{G} as the *Clifford+CS* group or *Clifford+controlled-phase* group. We know from [1] that \mathcal{G} is the group of 4×4 unitary matrices of the form

$$\frac{1}{\sqrt{2}^k} M \tag{1}$$

where $k \in \mathbb{N}$ and the entries of M belong to $\mathbb{Z}[i]$. In the fault-tolerant setting, the CS gate is considered vastly more expensive than any of the Clifford gates. As a result, the cost of a Clifford+ CS circuit is determined by its *CS-count*: the number of CS gates that appear in the circuit. Our goal is to find circuits for the elements of \mathcal{G} that are optimal in CS -count.

We start by introducing a generalization of the CS gate which will be helpful in describing the elements of \mathcal{G} .

Definition 2.1. Let P and Q be distinct elements of $\mathcal{P} \setminus \{I\}$ such that P and Q are Hermitian and $PQ = QP$. Then $R(P, Q)$ is defined as

$$R(P, Q) = \exp\left(\frac{i\pi}{2} \left(\frac{I-P}{2}\right) \left(\frac{I-Q}{2}\right)\right).$$

We have $R(Z \otimes I, I \otimes Z) = CS$. Moreover, since \mathcal{C} normalizes \mathcal{P} and $CR(P, Q)C^\dagger = R(CPC^\dagger, CQC^\dagger)$ for every $C \in \mathcal{C}$, we know that $R(P, Q) \in \mathcal{G}$ for every appropriate $P, Q \in \mathcal{P}$. We record some important properties of the $R(P, Q)$ gates in the lemma below. Because the proof of the lemma is tedious but relatively straightforward, it is given in [Appendix B](#).

$$\begin{array}{ccccc}
R(X \otimes I, I \otimes X) & R(Y \otimes I, I \otimes Y) & R(Z \otimes I, I \otimes Z) & R(Y \otimes I, I \otimes Z) & R(Z \otimes I, I \otimes Y) \\
R(Z \otimes I, I \otimes X) & R(X \otimes I, I \otimes Z) & R(X \otimes I, I \otimes Y) & R(Y \otimes I, I \otimes X) & R(X \otimes X, Y \otimes Y) \\
R(X \otimes X, Z \otimes Y) & R(Z \otimes X, Y \otimes Y) & R(Y \otimes X, X \otimes Y) & R(Z \otimes X, X \otimes Y) & R(Y \otimes X, Z \otimes Y)
\end{array}$$

Figure 1: The 15 elements of \mathcal{S} .

Lemma 2.2. *Let $C \in \mathcal{C}$ and let P, Q , and L be distinct elements of $\mathcal{P} \setminus \{I\}$. Assume that P, Q , and L are Hermitian and that $PQ = QP$, $PL = LP$, and $QL = -LQ$. Then the following relations hold:*

$$CR(P, Q)C^\dagger = R(CPC^\dagger, CQC^\dagger), \quad (2)$$

$$R(P, Q) = R(Q, P), \quad (3)$$

$$R(P, -PQ) = R(P, Q), \quad (4)$$

$$R(P, -Q) \in R(P, Q)\mathcal{C}, \quad (5)$$

$$R(P, Q)^2 \in \mathcal{C}, \text{ and} \quad (6)$$

$$R(P, L)R(P, Q) = R(P, Q)R(P, iQL). \quad (7)$$

We will use the $R(P, Q)$ gates of [Definition 2.1](#) to define normal forms for the elements of \mathcal{G} . The equivalences given by [Lemma 2.2](#) show that it is not necessary to use every $R(P, Q)$ gate and the following definition specifies the ones we will be using.

Definition 2.3. Let \mathcal{T}_1 and \mathcal{T}_2 be the subsets of $\mathcal{P} \times \mathcal{P}$ given below.

$$\mathcal{T}_1 = \{(P, Q) ; P \in \{X \otimes I, Y \otimes I, Z \otimes I\}, Q \in \{I \otimes X, I \otimes Y, I \otimes Z\}\}$$

$$\mathcal{T}_2 = \{(P, Q) ; P \in \{X \otimes X, Z \otimes X, Y \otimes X\}, Q \in \{Y \otimes Y, Z \otimes Y, X \otimes Y\}, \text{ and } PQ = QP\}.$$

The set \mathcal{S} is defined as $\mathcal{S} = \{R(P, Q) ; (P, Q) \in \mathcal{T}_1 \text{ or } (P, Q) \in \mathcal{T}_2\}$.

The set \mathcal{S} contains 15 elements which are explicitly listed in [Figure 1](#). It can be verified that all of the elements of \mathcal{S} are distinct, even up to right-multiplication by a Clifford gate. It will be helpful to consider the set \mathcal{S} ordered as in [Figure 1](#), which is to be read left-to-right and row-by-row. We then write \mathcal{S}_j to refer to the j -th element of \mathcal{S} . For example, \mathcal{S}_1 is in the top left of [Figure 1](#), \mathcal{S}_5 is in the top right, and \mathcal{S}_{15} is in the bottom right. The position of $R(P, Q)$ in this ordering roughly expresses the complexity of the Clifford circuit required to conjugate CS to $R(P, Q)$.

We close this section by showing that every element of \mathcal{G} can be expressed as a sequence of elements of \mathcal{S} followed by a single element of \mathcal{C} .

Lemma 2.4. *Let P and Q be distinct elements of $\mathcal{P} \setminus \{I\}$ such that P and Q are Hermitian and $PQ = QP$. Then there exists $P', Q' \in \mathcal{P}$ and $C \in \mathcal{C}$ such that $R(P', Q') \in \mathcal{S}$ and $R(P, Q) = R(P', Q')C$.*

Proof. Let $P = i^p(P_1 \otimes P_2)$ and $Q = i^q(Q_1 \otimes Q_2)$ with $P_1, P_2, Q_1, Q_2 \in \{I, X, Y, Z\}$. Since P and Q are Hermitian, p and q must be even. Moreover, by [Equations \(3\) and \(5\) of Lemma 2.2](#), we can assume without loss of generality that $p = q = 0$ so that $P = P_1 \otimes P_2$ and $Q = Q_1 \otimes Q_2$. Now, if one of P_1, P_2, Q_1 , or Q_2 is I , then we can use [Equations \(3\), \(4\) and \(5\) of Lemma 2.2](#) to rewrite $R(P, Q)$ as $R(P', Q')C$ with $C \in \mathcal{C}$ and $(P', Q') \in \mathcal{T}_1$ as in [Definition 2.3](#). If, instead, none of P_1, P_2, Q_1 , or Q_2 are I , then we can reason similarly to rewrite $R(P, Q)$ as $R(P', Q')C$ with $C \in \mathcal{C}$ and $(P', Q') \in \mathcal{T}_2$. \square

Proposition 2.5. *Let $V \in \mathcal{G}$. Then $V = R_1 \cdots R_n C$ where $C \in \mathcal{C}$ and $R_j \in \mathcal{S}$ for $j \in [n]$.*

Proof. Let $V \in \mathcal{G}$. Then V can be written as $V = C_1 \cdot CS \cdot C_2 \cdot CS \cdots C_n \cdot CS \cdot C_{n+1}$ where $C_j \in \mathcal{C}$ for $j \in [n+1]$. Since $CS = R(Z \otimes I, I \otimes Z)$ we have

$$V = C_1 \cdot R(Z \otimes I, I \otimes Z) \cdot C_2 \cdot R(Z \otimes I, I \otimes Z) \cdots C_n \cdot R(Z \otimes I, I \otimes Z) \cdot C_{n+1}. \quad (8)$$

Now, by Equation (2) of Lemma 2.2, $C_1 R(Z \otimes I, I \otimes Z) = C_1 R(Z \otimes I, I \otimes Z) C_1^\dagger C_1 = R(P, Q) C_1$ for some $P, Q \in \mathcal{P}$. We can then apply Lemma 2.4 to get

$$C_1 R(Z \otimes I, I \otimes Z) = R(P, Q) C_1 = R(P', Q') C C_1 = R(P', Q') C'$$

with $C' = C C_1 \in \mathcal{C}$ and $R(P', Q') \in \mathcal{S}$. Hence, setting $R_1 = R(P', Q')$ and $C'_2 = C' C_2$, Equation (8) becomes

$$V = R_1 \cdot C'_2 \cdot R(Z \otimes I, I \otimes Z) \cdot \dots \cdot C_n \cdot R(Z \otimes I, I \otimes Z) \cdot C_{n+1}$$

and we can proceed recursively to complete the proof. \square

3 The Isomorphism $SU(4) \cong Spin(6)$

In this section, we describe the exceptional isomorphism $SU(4) \cong Spin(6)$ which will allow us to rewrite two-qubit operators as elements of $SO(6)$. Consider some element U of $SU(4)$. Then U acts on \mathbb{C}^4 by left-multiplication. Moreover, this action is norm-preserving. Now let $\{e_j\}$ be the standard orthonormal basis of \mathbb{C}^4 . From this basis, we construct an alternative six-component basis using the *wedge product*.

Definition 3.1 (Wedge product). Let $a \wedge b$ be defined as the *wedge product* of a and b . Wedge products have the following properties given vectors $a, b, c \in \mathbb{C}^n$ and $\alpha, \beta \in \mathbb{C}$:

- Anticommutativity: $a \wedge b = -b \wedge a$.
- Associativity: $(a \wedge b) \wedge c = a \wedge (b \wedge c)$.
- Bilinearity: $(\alpha a + \beta b) \wedge c = \alpha(a \wedge c) + \beta(b \wedge c)$.

Note that the anticommutation of wedge products implies that $a \wedge a = 0$. We say that $v_1 \wedge \dots \wedge v_k \in \bigwedge^k \mathbb{C}^n$ for $v_j \in \mathbb{C}^n$. To compute the inner product of two wedge products $v_1 \wedge \dots \wedge v_k$ and $w_1 \wedge \dots \wedge w_k$, we compute

$$\langle v_1 \wedge \dots \wedge v_k, w_1 \wedge \dots \wedge w_k \rangle = \det(\langle v_q, w_r \rangle)$$

where $\langle v_q, w_r \rangle$ is the entry in the q -th row and r -th column of a $k \times k$ matrix.

Remark 3.2. The magnitude of a wedge product of n vectors can be thought of as the n dimensional volume of the parallelotope constructed from those vectors. The orientation of the wedge product defines the direction of circulation around that parallelotope by those vectors.

The wedge product of two vectors in \mathbb{C}^4 can be decomposed into a six-component basis as anticommutativity reduces the 16 potential wedge products of elements of $\{e_j\}$ to six. We choose this basis as

$$B = \{s_{-,12,34}, s_{+,12,34}, s_{-,23,14}, s_{+,24,13}, s_{-,24,13}, s_{+,23,14}\} \quad (9)$$

where

$$s_{\pm,ij,kl} = \frac{i^{\frac{1 \mp 1}{2}}}{\sqrt{2}} (e_i \wedge e_j \pm e_k \wedge e_l). \quad (10)$$

We note that B is an orthonormal basis and we assume that B is ordered as in Equation (9).

Definition 3.3. Let $U \in SU(4)$ and \overline{U} be its representation in the transformed basis. Let $v, w \in \mathbb{C}^4$ with $v \wedge w \in \bigwedge^2 \mathbb{C}^4$. Then the actions of U and \overline{U} are related by

$$\overline{U}(v \wedge w) = (Uv) \wedge (Uw).$$

To avoid confusion, we use an overline, as in \overline{O} , to denote the $SO(6)$ representation of an operator or set of operators O . We are now equipped to define the transformation from $SU(4)$ to $SO(6)$.

Definition 3.4. Let $U \in \text{SU}(4)$ and let $j, k \in [6]$. Then the entry in the j -th row and k -th column of the $\text{SO}(6)$ representation \bar{U} of U is

$$\bar{U}_{j,k} = \langle B_j, \bar{U} B_k \rangle \quad (11)$$

where B_j is the j -th element in the ordered basis B , the action of \bar{U} on B_k is defined by [Definitions 3.1](#) and [3.3](#), and the inner product is defined by [Definition 3.1](#).

As an illustration of the process specified in [Definition 3.4](#) we explicitly calculate the $\text{SO}(6)$ representation of a Clifford+ CS operator in [Appendix A](#). Moreover, we provide code to compute this isomorphism for any input with our Mathematica package [\[16\]](#).

Remark 3.5. The fact that this isomorphism yields special orthogonal operators is ultimately due to the fact that the Dynkin diagrams for the Lie algebras of $\text{SU}(4)$, $\text{Spin}(6)$, and $\text{SO}(6)$ are equivalent. However, this fact can be easily illustrated through the Euler decomposition of $\text{SU}(4)$ [\[36\]](#). Direct calculation of \bar{U} for the operator

$$U = \begin{bmatrix} 1 & \cdot & \cdot & \cdot \\ \cdot & 1 & \cdot & \cdot \\ \cdot & \cdot & \alpha & \cdot \\ \cdot & \cdot & \cdot & \alpha^* \end{bmatrix}$$

for $|\alpha| = 1$ and $\alpha = r + ic$ with $r, c \in \mathbb{R}$ yields

$$\bar{U} = \begin{bmatrix} 1 & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & 1 & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & r & \cdot & \cdot & c \\ \cdot & \cdot & \cdot & r & c & \cdot \\ \cdot & \cdot & \cdot & -c & r & \cdot \\ \cdot & \cdot & -c & \cdot & \cdot & r \end{bmatrix}$$

which is explicitly in $\text{SO}(6)$. Computation of the other 14 Euler angle rotations required for an $\text{SU}(4)$ parameterization yields similar matrices, likewise in $\text{SO}(6)$. Since $\text{SO}(6)$ is a group under multiplication, the isomorphism applied to any $U \in \text{SU}(4)$ yields $\bar{U} \in \text{SO}(6)$.

We close this section by explicitly calculating the $\text{SO}(6)$ representation of each of the generators of \mathcal{G} . We multiply the generators by overall phase factors to ensure that each operator has determinant one, and furthermore that single-qubit operators have determinant one on their single-qubit subspace. Later, when referring to gates or their $\text{SO}(6)$ representation, we omit overall phases for readability.

Proposition 3.6. *The image of the generators of \mathcal{C} in $\text{SO}(6)$ are*

$$\begin{aligned} \overline{(\omega^\dagger S) \otimes I} &= \begin{bmatrix} \cdot & -1 & \cdot & \cdot & \cdot & \cdot \\ 1 & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & 1 & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & 1 & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & 1 & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & 1 \end{bmatrix}, & \overline{I \otimes (\omega^\dagger S)} &= \begin{bmatrix} 1 & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & 1 & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & 1 & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & -1 & \cdot \\ \cdot & \cdot & \cdot & 1 & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & 1 \end{bmatrix}, \\ \overline{(iH) \otimes I} &= \begin{bmatrix} \cdot & \cdot & 1 & \cdot & \cdot & \cdot \\ \cdot & -1 & \cdot & \cdot & \cdot & \cdot \\ 1 & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & 1 & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & 1 & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & 1 \end{bmatrix}, & \overline{I \otimes (iH)} &= \begin{bmatrix} 1 & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & 1 & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & 1 & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & 1 \\ \cdot & \cdot & \cdot & \cdot & -1 & \cdot \\ \cdot & \cdot & \cdot & 1 & \cdot & \cdot \end{bmatrix}, \end{aligned}$$

$$\overline{\omega^\dagger CZ} = \begin{bmatrix} \cdot & -1 & \cdot & \cdot & \cdot & \cdot \\ 1 & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & -1 \\ \cdot & \cdot & \cdot & \cdot & -1 & \cdot \\ \cdot & \cdot & \cdot & 1 & \cdot & \cdot \\ \cdot & \cdot & 1 & \cdot & \cdot & \cdot \end{bmatrix}.$$

Proposition 3.7. *The elements of $\overline{\mathcal{S}}$ are given in [Figure 2](#).*

4 Exact Synthesis

In this section, we leverage the isomorphism $SU(4) \cong Spin(6)$ described in the previous section to find optimal decompositions for the elements of \mathcal{G} . We will be working extensively with the matrix group

$$\mathcal{H} = \left\{ \frac{1}{\sqrt{2^k}} M \in SO(6) ; k \in \mathbb{N}, M \in \mathbb{Z}^{6 \times 6} \right\}. \quad (12)$$

Note that $\mathcal{H} \subseteq SO(6)$. Our interest in \mathcal{H} stems from the following observation.

Proposition 4.1. *We have $\overline{\mathcal{G}} \subseteq \mathcal{H}$.*

Proof. The property holds for the generators of $\overline{\mathcal{G}}$ by [Propositions 3.6](#) and [3.7](#). \square

In the remainder of this section, we prove the converse of [Proposition 4.1](#) by defining an algorithm which inputs an element of \mathcal{H} and outputs a product of generators. We start by introducing a few notions that are useful in discussing the elements of \mathcal{H} .

Definition 4.2. Let $V \in \mathcal{H}$. We say that $\ell \in \mathbb{N}$ is a *denominator exponent* of V if $\sqrt{2}^\ell V \in \mathbb{Z}^{6 \times 6}$. The least such ℓ is the *least denominator exponent* of V , which we denote by $\text{lde}(V)$.

Lemma 4.3. *Let $U \in \mathcal{G}$ and suppose that $\text{lde}(\overline{U}) = k$. Then any Clifford+CS circuit for U has CS-count at least k .*

Proof. The only generators with a factor of $1/\sqrt{2}$ in their $SO(6)$ representation are the elements of \mathcal{S} . Thus, for a least denominator exponent of k there must be at least k of these operators, each of which requires a single CS gate. \square

Definition 4.4. Let $V \in \mathcal{H}$ and let ℓ be a denominator exponent of V . The ℓ -*residue* of V is the binary matrix $\rho_\ell(V) \in \mathbb{Z}_2^{6 \times 6}$ defined by

$$(\rho_\ell(V))_{i,j} = \rho((\sqrt{2}^\ell V)_{i,j})$$

where $\rho : \mathbb{Z} \rightarrow \mathbb{Z}_2$ is the canonical (parity) homomorphism.

The residue matrices introduced in [Definition 4.4](#) are important in the definition of the exact synthesis algorithm. Indeed, the ℓ -residue of a Clifford+CS operator U determines the element of \mathcal{S} to use in order to reduce the least denominator exponent of U (although not uniquely, as we discuss below). Similar residue matrices are used in the study of other fault-tolerant circuits [[1](#), [14](#)].

Recall that if A is a set, then a *partition* of A is a collection of disjoint nonempty subsets of A whose union is equal to A . The set of all partitions of a set A is denoted \mathcal{B}_A . Let p and p' be two partitions of A . If every element of p is a subset of an element of p' then we say that p' is *coarser* than p and that p is *finer* than p' .

Definition 4.5. Let $N \in \mathbb{Z}_2^{6 \times 6}$ be a binary matrix with rows r_1, \dots, r_6 and let $p = \{p_1, \dots, p_q\}$ be a partition of the set $[6]$. Then N has the *pattern* p if for any p_j in p and any $j_1, j_2 \in p_j$ we have $r_{j_1} = r_{j_2}$. In this case we also say that N has a $|p_1| \times \dots \times |p_q|$ *pattern*.

Definition 4.6. Let $V \in \mathcal{H}$ with $\text{lde}(V) = \ell$. We define the pattern map $\mathbf{p} : \mathcal{H} \rightarrow \mathcal{B}_{[6]}$ as the function which maps V to the pattern of $\rho_\ell(V)$. We say that $p = \mathbf{p}(V)$ is the pattern of V . If V_1 and V_2 are two elements of \mathcal{H} , we say that V_1 is *finer* than V_2 or that V_2 is *coarser* than V_1 if these statements hold for $\mathbf{p}(V_1)$ and $\mathbf{p}(V_2)$.

Remark 4.7. In a slight abuse of notation, we extend the pattern map to any valid representation of a Clifford+CS operator. Given a Clifford+CS operator with $\text{SU}(4)$ representation U which can be written as a word W over the generators and with $\text{SO}(6)$ representation \bar{U} , we set $\mathbf{p}(U) = \mathbf{p}(W) = \mathbf{p}(\bar{U})$. This extension is unambiguous after fixing our transformation from $\text{SU}(4)$ to $\text{SO}(6)$, as \mathbf{p} is insensitive to relative phase changes in U . We incorporate all relational notions described in [Definition 4.6](#) in this extension.

We now analyze the image in $\text{SO}(6)$ of certain subsets of \mathcal{G} . We start by showing that the image of the Clifford group \mathcal{C} is exactly the collection of elements of \mathcal{H} with least denominator 0. In other words, $\bar{\mathcal{C}}$ is the group of 6-dimensional signed permutation matrices.

Lemma 4.8. *Let $V \in \mathcal{H}$. Then $\text{lde}(V) = 0$ if and only if $V \in \bar{\mathcal{C}}$.*

Proof. The least denominator exponent of $\overline{H \otimes I}$, $\overline{I \otimes H}$, $\overline{S \otimes I}$, $\overline{I \otimes S}$, and \overline{CZ} is 0. Thus, if $U \in \mathcal{C}$ then $\text{lde}(\bar{U}) = 0$. For the converse, let C_1 and C_2 be the Clifford operators $(\omega^\dagger S) \otimes I$ and $(H \otimes H)(\omega^\dagger CZ)(Z \otimes Z)$, respectively. Then

$$\bar{C}_1 = \begin{bmatrix} \cdot & -1 & \cdot & \cdot & \cdot & \cdot \\ 1 & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & 1 & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & 1 & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & 1 & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & 1 \end{bmatrix} \quad \text{and} \quad \bar{C}_2 = \begin{bmatrix} \cdot & \cdot & \cdot & \cdot & \cdot & -1 \\ 1 & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & 1 & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & 1 & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & 1 & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & 1 & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & 1 \end{bmatrix}.$$

The operators \bar{C}_1 and \bar{C}_2 generate $\{V \in \mathcal{H} ; \text{lde}(V) = 0\}$. Hence, if $V \in \mathcal{H}$ and $\text{lde}(V) = 0$ then V can be expressed as a product of the image of Clifford gates. \square

Lemma 4.9. *Let $V \in \mathcal{H}$. Then $\text{lde}(V) = 1$ if and only if $V = \overline{RC}$ for some $R \in \mathcal{S}$ and some $C \in \mathcal{C}$. Furthermore, V has a $2 \times 2 \times 2$ pattern.*

Proof. The rows of V have unit norm and are pairwise orthogonal. Hence, up to a signed permutation of rows and columns, there is only one such matrix, e.g.,

$$\frac{1}{\sqrt{2}} \begin{bmatrix} 1 & -1 & \cdot & \cdot & \cdot & \cdot \\ 1 & 1 & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & 1 & -1 & \cdot & \cdot \\ \cdot & \cdot & 1 & 1 & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & 1 & -1 \\ \cdot & \cdot & \cdot & \cdot & 1 & 1 \end{bmatrix} = \bar{\mathcal{S}}_6. \quad (13)$$

By [Proposition 2.5](#) the proof is complete, since Clifford operators correspond to signed permutations by [Lemma 4.8](#). \square

Lemma 4.10. *Let $V \in \mathcal{H}$ with $\text{lde}(V) = k \geq 2$. Then V has either a $2 \times 2 \times 2$ or 2×4 pattern.*

Proof. Let $V \in \mathcal{H}$. Since V is orthogonal, we have $V^\dagger V = I$. Hence, $(\sqrt{2^k} V)^\dagger (\sqrt{2^k} V) = 2^k I$. Since $k \geq 2$, this implies that the inner product of any column of $\sqrt{2^k} V$ with itself is congruent to 0 modulo 4. Similarly, the inner product of two distinct columns of $\sqrt{2^k} V$ is congruent to 0 modulo 4. Letting, $M = \rho_k(V)$, we then have the column relations

$$\sum_l M_{lm}^2 = 0 \pmod{4} \quad (14)$$

$$\sum_l M_{lm} M_{ln} = 0 \pmod{2} \text{ for } m \neq n \quad (15)$$

as well as analogous row relations. For $x \in \mathbb{Z}$, $x^2 = 0 \pmod{4}$ if and only if $x = 0 \pmod{2}$. Hence, there must be exactly zero or four odd entries in every column (or row) of M by Equation (14). By Equation (15), we see that the inner product of any two distinct rows must be even. Up to a permutation of rows and columns, we can then deduce that M is one of the two matrices below, which completes the proof.

$$\begin{bmatrix} 1 & 1 & 1 & 1 & \cdot & \cdot \\ 1 & 1 & 1 & 1 & \cdot & \cdot \\ 1 & 1 & 1 & 1 & \cdot & \cdot \\ 1 & 1 & 1 & 1 & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \end{bmatrix} \quad \text{or} \quad \begin{bmatrix} 1 & 1 & 1 & 1 & \cdot & \cdot \\ 1 & 1 & 1 & 1 & \cdot & \cdot \\ 1 & 1 & \cdot & \cdot & 1 & 1 \\ 1 & 1 & \cdot & \cdot & 1 & 1 \\ \cdot & \cdot & 1 & 1 & 1 & 1 \\ \cdot & \cdot & 1 & 1 & 1 & 1 \end{bmatrix} \quad (16)$$

□

Corollary 4.11. *Let $V \in \mathcal{H}$ with $\text{lde}(V) = k \geq 1$. Then V has either a $2 \times 2 \times 2$ or 2×4 pattern.*

Lemma 4.12. *Let $V \in \mathcal{H}$ and assume that $\text{lde}(V) = k \geq 1$. If $\bar{R} \in \bar{\mathcal{S}}$ is finer than V , then $\text{lde}(\bar{R}^\top V) = k - 1$.*

Proof. For simplicity, we assume that $\mathfrak{p}(\bar{R}) = \{\{1, 2\}, \{3, 4\}, \{5, 6\}\}$. The cases in which $\mathfrak{p}(\bar{R})$ is another pattern are treated similarly. For $j \in [6]$, let r_j denote the rows of $\sqrt{2}^k V$. Since $\mathfrak{p}(V)$ is coarser than $\mathfrak{p}(\bar{R})$, we have $r_1 \equiv r_2$, $r_3 \equiv r_4$, $r_5 \equiv r_6$ modulo 2. This implies that $r_1 \pm r_2 \equiv r_3 \pm r_4 \equiv r_5 \pm r_6 \equiv 0$ modulo 2. Hence

$$\bar{R}^\top V = \frac{1}{\sqrt{2}^{k+1}} \begin{bmatrix} 1 & 1 & \cdot & \cdot & \cdot & \cdot \\ -1 & 1 & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & 1 & 1 & \cdot & \cdot \\ \cdot & \cdot & -1 & 1 & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & 1 & 1 \\ \cdot & \cdot & \cdot & \cdot & -1 & 1 \end{bmatrix} \begin{bmatrix} r_1 \\ r_2 \\ r_3 \\ r_4 \\ r_5 \\ r_6 \end{bmatrix} = \frac{1}{\sqrt{2}^{k+1}} \begin{bmatrix} r_1 - r_2 \\ r_1 + r_2 \\ r_3 - r_4 \\ r_3 + r_4 \\ r_5 - r_6 \\ r_5 + r_6 \end{bmatrix} = \frac{1}{\sqrt{2}^{k-1}} \begin{bmatrix} r'_1 \\ r'_2 \\ r'_3 \\ r'_4 \\ r'_5 \\ r'_6 \end{bmatrix}.$$

where each r'_j is a vector of integers. □

Lemma 4.13. *Let $V \in \mathcal{H}$ with $\text{lde}(V) \geq 1$. Then there exists $R \in \mathcal{S}$ such that $\text{lde}(\bar{R}^\top V) = \text{lde}(V) - 1$.*

Proof. By inspection of Figure 2 we see that for every $2 \times 2 \times 2$ pattern q there exists $R \in \mathcal{S}$ such that $\mathfrak{p}(\bar{R}) = q$. As a result, if $\mathfrak{p}(V)$ is a $2 \times 2 \times 2$ or a 2×4 pattern, then there exists $R \in \mathcal{S}$ such that \bar{R} has a pattern finer than $\mathfrak{p}(V)$. By Corollary 4.11, $\mathfrak{p}(V)$ is in fact a $2 \times 2 \times 2$ row-pattern or a 2×4 row-pattern and thus there exists $R \in \mathcal{S}$ such that \bar{R} is finer than V . We can then conclude by Lemma 4.12. □

Theorem 4.14. *We have $\bar{\mathcal{G}} = \mathcal{H}$.*

Proof. $\bar{\mathcal{G}} \subseteq \mathcal{H}$ by Proposition 4.1. We now show $\mathcal{H} \subseteq \bar{\mathcal{G}}$. Let $V \in \mathcal{H}$. We proceed by induction on the least denominator exponent of V . If $\text{lde}(V) = 0$ then, by Lemma 4.8, $V \in \bar{\mathcal{C}}$ and therefore $V \in \bar{\mathcal{G}}$. Now if $\text{lde}(V) > 0$, let R be the element of \mathcal{S} with the lowest index such that $\text{lde}(\bar{R}^\top V) = k - 1$. Such an element exists by Lemma 4.13. By the induction hypothesis we have $\bar{R}^\top V \in \bar{\mathcal{G}}$ which implies that $\bar{R}(\bar{R}^\top V) = V \in \bar{\mathcal{G}}$. □

The proof of Theorem 4.14 provides an algorithm to decompose an arbitrary element of $\bar{\mathcal{G}}$ into a product of elements of $\bar{\mathcal{S}}$, followed by an element of $\bar{\mathcal{C}}$. In the proof, there is freedom in choosing the element of $\bar{\mathcal{S}}$ used to reduce $\text{lde}(\bar{V})$. If there is more than one generator with a finer pattern than \bar{V} , we must make a choice. The ordering imposed on \mathcal{S} in Section 2 is used to make this choice in a uniform manner: we always choose the element of \mathcal{S} of lowest index. As a result, the exact synthesis algorithm becomes deterministic. The ambiguity in the choice of generator is a consequence of the relations given in Lemma 2.2. In particular, we have

$$R(P, L)R(P, Q) = R(P, Q)R(P, iQL) = R(P, iQL)R(P, L)$$

Generator	Associated Patterns Under First Finer Partition (FFP)
$R(X \otimes I, I \otimes X)$	$\{\{1, 4\}, \{2, 3\}, \{5, 6\}\}, \{\{1, 4\}, \{2, 3, 5, 6\}\}, \{\{2, 3\}, \{1, 4, 5, 6\}\}, \{\{5, 6\}, \{1, 2, 3, 4\}\}$
$R(Y \otimes I, I \otimes Y)$	$\{\{1, 3\}, \{2, 5\}, \{4, 6\}\}, \{\{1, 3\}, \{2, 4, 5, 6\}\}, \{\{2, 5\}, \{1, 3, 4, 6\}\}, \{\{4, 6\}, \{1, 2, 3, 5\}\}$
$R(Z \otimes I, I \otimes Z)$	$\{\{1, 2\}, \{3, 6\}, \{4, 5\}\}, \{\{1, 2\}, \{3, 4, 5, 6\}\}, \{\{3, 6\}, \{1, 2, 4, 5\}\}, \{\{4, 5\}, \{1, 2, 3, 6\}\}$
$R(Y \otimes I, I \otimes Z)$	$\{\{1, 3\}, \{2, 6\}, \{4, 5\}\}, \{\{2, 6\}, \{1, 3, 4, 5\}\}$
$R(Z \otimes I, I \otimes Y)$	$\{\{1, 2\}, \{3, 5\}, \{4, 6\}\}, \{\{3, 5\}, \{1, 2, 4, 6\}\}$
$R(Z \otimes I, I \otimes X)$	$\{\{1, 2\}, \{3, 4\}, \{5, 6\}\}, \{\{3, 4\}, \{1, 2, 5, 6\}\}$
$R(X \otimes I, I \otimes Z)$	$\{\{1, 6\}, \{2, 3\}, \{4, 5\}\}, \{\{1, 6\}, \{2, 3, 4, 5\}\}$
$R(X \otimes I, I \otimes Y)$	$\{\{1, 5\}, \{2, 3\}, \{4, 6\}\}, \{\{1, 5\}, \{2, 3, 4, 6\}\}$
$R(Y \otimes I, I \otimes X)$	$\{\{1, 3\}, \{2, 4\}, \{5, 6\}\}, \{\{2, 4\}, \{1, 3, 5, 6\}\}$
$R(X \otimes X, Y \otimes Y)$	$\{\{1, 4\}, \{2, 5\}, \{3, 6\}\}$
$R(X \otimes X, Z \otimes Y)$	$\{\{1, 4\}, \{2, 6\}, \{3, 5\}\}$
$R(Z \otimes X, Y \otimes Y)$	$\{\{1, 6\}, \{2, 5\}, \{3, 4\}\}$
$R(Y \otimes X, X \otimes Y)$	$\{\{1, 5\}, \{2, 4\}, \{3, 6\}\}$
$R(Z \otimes X, X \otimes Y)$	$\{\{1, 5\}, \{2, 6\}, \{3, 4\}\}$
$R(Y \otimes X, Z \otimes Y)$	$\{\{1, 6\}, \{2, 4\}, \{3, 5\}\}$

Table 1: The elements of \mathcal{S} and the explicit row patterns they are associated with under FFP.

and these three distinct sequences of generators denote the same operator. This is the source of the three-fold ambiguity in choosing a finer $2 \times 2 \times 2$ pattern for a given 2×4 pattern.

We will sometimes refer to the association between elements of \mathcal{S} and patterns used in the exact synthesis algorithm of [Theorem 4.14](#) as the *first finer partition* association, or FFP for short. The association is explicitly described [Table 1](#).

Theorem 4.15. *If U is a Clifford+CS operator such that $\text{lde}(\overline{U}) = k$, then U can be represented by a Clifford+CS circuit of CS-count k . This circuit is optimal in CS-count and can be constructed in $\mathcal{O}(k)$ arithmetic operations.*

Proof. Let U be as stated. If $k = 0$, then \overline{U} belongs to $\overline{\mathcal{C}}$ and U is therefore a Clifford. If $k > 0$, then as in [Theorem 4.14](#), there is a unique $R_k \in \mathcal{S}$ given by FFP such that $\text{lde}(\overline{R}_k^T \overline{U}) = k - 1$. By induction on the least denominator exponent, we have a deterministic synthesis algorithm to find a sequence such that

$$\overline{U} = \overline{R}_k \cdots \overline{R}_1 \cdot \overline{C}$$

which then implies that $U = R_k \cdots R_1 C$. Each of these k steps involves a constant number of basic arithmetic operations. This circuit has CS-count k , which is optimal by [Lemma 4.3](#). \square

Our Mathematica package [\[16\]](#) implements the algorithm referred to in [Theorem 4.15](#) as well as a significant amount of other tools for two-qubit Clifford + CS circuits. Testing of the performance of this algorithm on a modest device is presented in [Table 2](#).

5 Normal Forms

In the previous section, we introduced a synthesis algorithm for Clifford+CS operators. The algorithm takes as input a Clifford+CS matrix and outputs a circuit for the corresponding operator. The circuit produced by the synthesis algorithm is a word over the alphabet $\mathcal{S} \cup \mathcal{C}$. Because the algorithm is deterministic, the word it associates to each operator can be viewed as a normal form for that operator. In the present section, we use the language of automata to give a detailed description of the structure of these normal forms. We include the definitions of some basic concepts from the theory of automata for completeness. The reader looking for further details is encouraged to consult [\[35\]](#).

5.1 Automata


In what follows we sometimes refer to a finite set Σ as an *alphabet*. In such a context, the elements of Σ are referred to as *letters*, Σ^* denotes the set of *words* over Σ (which includes the empty word ε), and the subsets of Σ^* are called *languages over Σ* . If $w \in \Sigma^*$ is a word over the alphabet Σ , we write $|w|$ for the *length* of w . Finally, if L and L' are two languages over an alphabet Σ then their *concatenation* $L \circ L'$ is defined as $L \circ L' = \{ww' ; w \in L \text{ and } w' \in L'\}$.

Definition 5.1. A *nondeterministic finite automaton* is a 5-tuple $(\Sigma, Q, \text{In}, \text{Fin}, \delta)$ where Σ and Q are finite sets, In and Fin are subsets of Q , and $\delta : Q \times (\Sigma \cup \{\varepsilon\}) \rightarrow \mathcal{P}(Q)$ is a function whose codomain is the power set of Q . We call Σ the *alphabet*, Q the set of *states*, In and Fin the sets of *initial* and *final* states, and δ the *transition function*.

Remark 5.2. **Definition 5.1** is slightly non-standard. Indeed, automata are typically defined as having a single initial state, rather than a collection of them. One can then think of **Definition 5.1** as introducing a collection of automata: one for each element of In . Alternatively, **Definition 5.1** can also be recovered from the usual definition by assuming that every automaton in the sense of **Definition 5.1** in fact has a single initial state s_0 related to the elements of In by $\delta(s_0, \varepsilon) = \text{In}$. We chose to introduce automata as in **Definition 5.1** because this results in a slightly cleaner presentation.

It is common to define an automaton $A = (\Sigma, Q, \text{In}, \text{Fin}, \delta)$ by specifying a directed labelled graph called the *state graph* of A . The vertices of the graph are labelled by states and there is an edge labelled by a letter $w \in \Sigma$ between vertices labelled q and q' if $q' \in \delta(q, w)$. The initial and final states are distinguished using arrows and double lines, respectively. For brevity, parallel edges are drawn only once, with their labels separated by a comma.

Example 5.3. The state graph for a nondeterministic finite automaton $A = (\Sigma, Q, \delta, \text{In}, \text{Fin})$ is depicted below.



Here, $Q = \{q_0, q_1, q_2, q_3\}$, $\Sigma = \{0, 1\}$, the collection of initial states is $\text{In} = \{q_0\}$, the collection of final states is $\text{Fin} = \{q_3\}$, and we have, e.g., $\delta(q_0, 1) = \{q_0, q_1\}$.

An automaton $A = (\Sigma, Q, \text{In}, \text{Fin}, \delta)$ can be used to specify a language $\mathcal{L}(A) \subseteq \Sigma^*$. Intuitively, $\mathcal{L}(A)$ is the collection of all the words over Σ that specify a well-formed walk along the state graph of A . The following definition makes this intuition more precise.

<i>CS</i> -count	Mean Time (s)	Std. Dev. (s)
10	0.0138	0.0044
100	0.0281	0.0051
1000	0.1135	0.0091
10000	1.1883	0.0897

Table 2: Performance of the algorithm (in seconds) of **Theorem 4.15** as implemented in our Mathematica code [16]. Each run has constant overhead from computing the $\text{SO}(6)$ representation for each unitary. Deviations from linearity are due to arithmetic operations on increasingly large integers. Each mean and standard deviation is computed using a sample of 1000 runs with pseudorandomly generated operators known to have the given minimal *CS*-count. Times are measured using Mathematica’s in-built `AbsoluteTiming` function. Computations performed on a laptop with an Intel(R) Core(TM) i7 CPU running at 2.6 GHz with 6 cores and 16 GB of RAM running macOS Catalina version 10.15.7.

Definition 5.4. Let $A = (\Sigma, Q, \text{In}, \text{Fin}, \delta)$ be an automaton. Then A *accepts* a word $w = w_1 \cdots w_m \in \Sigma^*$ if there exists a sequence of states $s_0, s_1, \dots, s_m \in Q$ such that

1. $s_0 \in \text{In}$,
2. $s_{j+1} \in \delta(s_j, w_{j+1})$ for $j \in \{0, \dots, m-1\}$, and
3. $s_m \in \text{Fin}$.

The set of words accepted by A is called the language *recognized* by A and is denoted $\mathcal{L}(A)$.

Example 5.5. The alphabet for the automaton A given in [Example 5.3](#) is $\Sigma = \{0, 1\}$. The language recognized by A is $\mathcal{L}(A) = \{w \in \Sigma^* ; \text{the third rightmost letter of } w \text{ is } 1\}$.

If a language is recognized by some nondeterministic finite automata then that language is called *regular*. The collection of regular languages is closed under a variety of operations. In particular, regular languages are closed under concatenation.

Definition 5.6. Let $A = (\Sigma, Q, \text{In}, \text{Fin}, \delta)$ and $A' = (\Sigma, Q', \text{In}', \text{Fin}', \delta')$ be two automata. Then the *concatenation* of A and A' is the automaton $A \circ A' = (\Sigma, Q'', \text{In}, \text{Fin}', \delta'')$ where $Q'' = Q \sqcup Q'$ is the disjoint union of Q and Q' and

$$\delta''(q, s) = \begin{cases} \delta(q, s) & q \in Q \setminus \text{Fin}, \\ \delta(q, s) & q \in \text{Fin} \text{ and } s \neq \varepsilon, \\ \delta(q, s) \cup \text{In}' & q \in \text{Fin} \text{ and } s = \varepsilon, \text{ and} \\ \delta'(q, s) & q \in Q'. \end{cases}$$

Proposition 5.7. Let A and A' be automata recognizing languages L and L' , respectively. Then $A \circ A'$ recognizes $L \circ L'$.

An example of the concatenation of two automata is provided in [Figure 3](#) and [Example 5.11](#) based off of the automata defined in [Definitions 5.9](#) and [5.10](#) below.

5.2 The Structure of Normal Forms

We now consider the alphabet $\mathcal{S} \cup \mathcal{C}$ and describe the words over $\mathcal{S} \cup \mathcal{C}$ that are output by the synthesis algorithm of [Theorem 4.15](#).

Definition 5.8. Let $U \in \mathcal{G}$. The *normal form* of U is the unique word over $\mathcal{S} \cup \mathcal{C}$ output by the synthesis algorithm of [Theorem 4.15](#) on input U . We write \mathcal{N} for the collection of all normal forms.

To describe the elements of \mathcal{N} , we introduce several automata. It will be convenient for our purposes to enumerate the elements of \mathcal{C} . We therefore assume that a total ordering of the 92160 elements of \mathcal{C} is chosen and we write \mathcal{C}_j for the j -th element of \mathcal{C} .

Definition 5.9. Let $k = |\mathcal{C}|$ and $\Sigma = \mathcal{S} \cup \mathcal{C}$. The automaton \mathfrak{C} is defined as $\mathfrak{C} = (\Sigma, [0, k], \{0\}, [k], \delta_{\mathfrak{C}})$ where, for $s \in [0, k]$ and $\ell \in \Sigma$, we have

$$\delta_{\mathfrak{C}}(s, \ell) = \begin{cases} \{j\} & \text{if } s = 0 \text{ and } \ell = \mathcal{C}_j, \text{ and} \\ \emptyset & \text{otherwise.} \end{cases}$$

Definition 5.10. Let $\Sigma = \mathcal{S} \cup \mathcal{C}$. The automaton $\mathfrak{S}_{n,m}$ is defined as $\mathfrak{S}_{n,m} = (\Sigma, [m], [n, m], [m], \delta_{\mathfrak{S},m})$ where, for $s \in [m]$ and $\ell \in \Sigma$, we have

$$\delta_{\mathfrak{S},m}(s, \ell) = \begin{cases} \{t ; \mathfrak{p}(\overline{\mathcal{S}_s}) \cap \mathfrak{p}(\overline{\mathcal{S}_t}) = \emptyset\} & \text{if } \ell = \mathcal{S}_s \text{ and} \\ \emptyset & \text{otherwise.} \end{cases}$$




Figure 3: The automaton $\mathfrak{S}_{1,3} \circ \mathfrak{C}$. The set of states of this automata is $\{1, 2, 3, 0', 1', \dots, k'\}$, which is the disjoint union of the states $\{1, 2, 3\}$ of $\mathfrak{S}_{1,3}$ and the states $\{0, 1, \dots, k\}$ of \mathfrak{C} . The initial states are $\{1, 2, 3\}$, those of $\mathfrak{S}_{1,3}$, and the final states are $\{1', \dots, k'\}$, those of \mathfrak{C} . Because $\mathfrak{S}_{1,3}$ has $\text{Fin} = \{1, 2, 3\}$ and \mathfrak{C} has $\text{In} = \{0'\}$, the transition function δ of $\mathfrak{S}_{1,3} \circ \mathfrak{C}$ is such that $\delta(1, \varepsilon) = \delta(2, \varepsilon) = \delta(3, \varepsilon) = \{0'\}$. Otherwise, δ behaves like the transition function for $\mathfrak{S}_{1,3}$ on the subset of states $\{1, 2, 3\}$ and like the transition function for \mathfrak{C} on the subset of states $\{0', 1', \dots, k'\}$.

Example 5.11. To illustrate [Definitions 5.6, 5.9](#) and [5.10](#), the automaton $\mathfrak{S}_{1,3} \circ \mathfrak{C}$ is represented in [Figure 3](#). It can be verified that the words C_2 , $S_2S_1C_1$, and $S_3S_1S_2C_k$ are accepted by $\mathfrak{S}_{1,3} \circ \mathfrak{C}$ while the words $S_1S_1C_4$ and $S_3C_7S_1$ are not. Note in particular that if C_1 is the symbol for the identity, then S_3C_1 is distinct (as a word) from S_3 . The former is accepted by $\mathfrak{S}_{1,3} \circ \mathfrak{C}$ while the latter is not. Despite the state graph of $\mathfrak{S}_{1,3}$ being fully-connected, full-connectivity does not necessarily hold for state graphs of other $\mathfrak{S}_{n,m}$ automata.

We will use the automata introduced in [Definitions 5.9](#) and [5.10](#) to describe the elements of \mathcal{N} . Our goal is to show that

$$\mathcal{N} = \mathcal{L}(\mathfrak{S}_{1,3} \circ \mathfrak{S}_{4,9} \circ \mathfrak{S}_{10,15} \circ \mathfrak{C}) \tag{17}$$

We start by establishing a few propositions.

Proposition 5.12. *We have $\mathcal{L}(\mathfrak{C}) \subsetneq \mathcal{L}(\mathfrak{S}_{1,15} \circ \mathfrak{C}) \subsetneq \mathcal{L}(\mathfrak{S}_{1,9} \circ \mathfrak{S}_{10,15} \circ \mathfrak{C}) \subsetneq \mathcal{L}(\mathfrak{S}_{1,3} \circ \mathfrak{S}_{4,9} \circ \mathfrak{S}_{10,15} \circ \mathfrak{C})$, where \subsetneq denotes strict inclusion.*

Proof. By [Definitions 5.9](#) and [5.10](#). □

We emphasize that the inclusions in [Proposition 5.12](#) are strict. This implies that $\mathcal{L}(\mathfrak{S}_{1,3} \circ \mathfrak{S}_{4,9} \circ \mathfrak{S}_{10,15} \circ \mathfrak{C})$ can be written as the disjoint union of $\mathcal{L}(\mathfrak{C})$, $\mathcal{L}(\mathfrak{S}_{1,15} \circ \mathfrak{C})$, and $\mathcal{L}(\mathfrak{S}_{1,9} \circ \mathfrak{S}_{10,15} \circ \mathfrak{C})$. The lemmas below show that these languages correspond to disjoint subsets of \mathcal{N} and, in combination, suffice to prove [Equation \(17\)](#).

Lemma 5.13. *Let U be a word over $\mathcal{S} \cup \mathcal{C}$. Then $U \in \mathcal{L}(\mathfrak{C})$ if and only if $U \in \mathcal{N}$ and U has length 1, i.e. $U \in \mathcal{C}$.*

Proof. By [Definition 5.9](#) and [Theorem 4.15](#). □

Lemma 5.14. *Let U be a word over $\mathcal{S} \cup \mathcal{C}$. Then $U \in \mathcal{L}(\mathfrak{S}_{1,15} \circ \mathfrak{C}) \setminus \mathcal{L}(\mathfrak{C})$ if and only if $U \in \mathcal{N}$ and U has a $2 \times 2 \times 2$ pattern.*

Proof. First, note that $\mathcal{L}(\mathfrak{C})$ is the set of words of length 1 accepted by $\mathfrak{S}_{1,15} \circ \mathfrak{C}$. This means that $\mathcal{L}(\mathfrak{S}_{1,15} \circ \mathfrak{C}) \setminus \mathcal{L}(\mathfrak{C})$ consists of all the words of length $k \geq 2$ accepted by $\mathfrak{S}_{1,15} \circ \mathfrak{C}$. Furthermore, by [Lemma 4.8](#), there are no normal forms of length 1 which have a $2 \times 2 \times 2$ pattern. Thus, to prove our lemma it suffices to establish the following equality of sets

$$\{U \in \mathcal{L}(\mathfrak{S}_{1,15} \circ \mathfrak{C}) ; |U| = k\} = \{U \in \mathcal{N} ; |U| = k \text{ and } \mathfrak{p}(U) \text{ is a } 2 \times 2 \times 2 \text{ pattern}\} \quad (18)$$

for all $k \geq 2$. We proceed by induction on k .

- Note that, by definition of $\mathfrak{S}_{1,15} \circ \mathfrak{C}$, we have $\{U \in \mathcal{L}(\mathfrak{S}_{1,15} \circ \mathfrak{C}) ; |U| = 2\} = \mathcal{SC}$. Every element of \mathcal{SC} has a $2 \times 2 \times 2$ pattern by [Lemma 4.9](#). Moreover, for $U = SC$ with $S \in \mathcal{S}$ and $C \in \mathcal{C}$, $\mathfrak{p}(SC) = \mathfrak{p}(S)$. Thus, SC must also be the unique word produced by the synthesis algorithm on input U and hence $U \in \mathcal{N}$. This accounts for all words of length 2 in \mathcal{N} . Therefore [Equation \(18\)](#) holds when $k = 2$.
- Now suppose that [Equation \(18\)](#) holds for some $k \geq 2$. Let $U \in \mathcal{L}(\mathfrak{S}_{1,15} \circ \mathfrak{C})$ be a word of length k whose first letter is $S \in \mathcal{S}$. Then $U \in \mathcal{N}$ and $\mathfrak{p}(U) = \mathfrak{p}(S)$ is a $2 \times 2 \times 2$ pattern. Furthermore, the least denominator exponent of \bar{U} is $k - 1$. We will show that [Equation \(18\)](#) holds for $k + 1$ by establishing two inclusions. Because it will sometimes be convenient to refer to submatrices, if M is an $n \times n$ matrix and $x, y \subseteq [n]$, we write

$$M[x; y]$$

for the submatrix of M formed from the rows with index in x and the columns with index in y .

\subseteq : Suppose that $U' = S'U$ is a word of length $k + 1$ accepted by $\mathcal{L}(\mathfrak{S}_{1,15} \circ \mathfrak{C})$. Then by [Definition 5.10](#) we have $\mathfrak{p}(S') \cap \mathfrak{p}(S) = \emptyset$. Let $\{a, b\} \in \mathfrak{p}(S')$, and let r_a and r_b be the corresponding rows of the residue matrix of \bar{U} . Explicitly, we have

$$\rho_{k-1}(\bar{U})[\{a, b\}; [6]] = \begin{bmatrix} r_a \\ r_b \end{bmatrix}$$

with $r_a \neq r_b$ as $\{a, b\}$ is not a subset of any element of $\mathfrak{p}(U)$. Direct calculation of the rows of the residue matrix for \bar{U}' yields

$$\rho_k(\bar{U}')[\{a, b\}; [6]] = \begin{bmatrix} r_a + r_b \\ r_a + r_b \end{bmatrix}.$$

We conclude that $\{a, b\}$ is a subset of an element of $\mathfrak{p}(U')$. Furthermore, by [Lemma 4.10](#) and [Equation \(16\)](#) we see that, since $r_a + r_b \neq 0$, $\mathfrak{p}(U')$ cannot be a 2×4 pattern, and therefore $\{a, b\} \in \mathfrak{p}(U')$. As this holds for all $\{a, b\} \in \mathfrak{p}(S')$, we conclude that $\mathfrak{p}(S') = \mathfrak{p}(U')$. Thus, by the induction hypothesis, $S'U$ will be the word produced by the synthesis algorithm when applied to U' . Hence, $U' \in \mathcal{N}$ and $\mathfrak{p}(U')$ is a $2 \times 2 \times 2$ pattern.

\supseteq : Suppose that U' is a normal form of length $k + 1$ with a $2 \times 2 \times 2$ pattern. Write U' as $U' = S'V$ for some unknown normal form V . We then have $\mathfrak{p}(S') = \mathfrak{p}(U')$. Let $\{a, b\} \in \mathfrak{p}(S')$ and let the corresponding rows of the residue matrix of \bar{V} be r_a and r_b . Explicitly, we have

$$\rho_{k-1}(\bar{V})[\{a, b\}; [6]] = \begin{bmatrix} r_a \\ r_b \end{bmatrix}.$$

Direct calculation of the rows of the residue matrix for \bar{U}' yields

$$\rho_k(\bar{U}')[\{a, b\}; [6]] = \begin{bmatrix} r_a + r_b \\ r_a + r_b \end{bmatrix}.$$

Since $\mathfrak{p}(U')$ is not a 2×4 pattern, we conclude that $r_a + r_b \neq 0$ and thus that $r_a \neq r_b$. Therefore, there is no element of cardinality four in $\mathfrak{p}(V)$. Since $\text{lde}(V) > 0$, $\mathfrak{p}(V)$ must then be a $2 \times 2 \times 2$ pattern. Consequently, we have $V = U$ as defined above. Because $\{a, b\} \notin \mathfrak{p}(U) = \mathfrak{p}(S)$, we know $\mathfrak{p}(S') \cap \mathfrak{p}(S) = \emptyset$. Given that $S' = \mathcal{S}_{j'}$ and $S = \mathcal{S}_j$, we conclude that $j \in \delta_{\mathfrak{S}, 15}(j', S' = \mathcal{S}_{j'})$. Because $S = \mathcal{S}_j$ is the first letter of the word U , we know the initial state of U must be j . Therefore, by the induction hypothesis, $U' = S'U$ is accepted by $\mathfrak{S}_{1,15} \circ \mathfrak{C}$.

We have shown that [Equation \(18\)](#) holds for words of length $k + 1$ if it holds for words of length k . This completes the inductive step. \square

[Lemma 5.14](#) characterized the normal forms that have a $2 \times 2 \times 2$ pattern. The two lemmas below jointly characterize the normal forms that have a 2×4 pattern. Because their proofs are similar in spirit to that of [Lemma 5.14](#), they have been relegated to [Appendix C](#).

Lemma 5.15. *Let U be a word over $\mathcal{S} \cup \mathcal{C}$. Then $U \in \mathcal{L}(\mathfrak{S}_{1,9} \circ \mathfrak{S}_{10,15} \circ \mathfrak{C}) \setminus \mathcal{L}(\mathfrak{S}_{1,15} \circ \mathfrak{C})$ if and only if $U \in \mathcal{N}$ and U has a 2×4 pattern with $\mathfrak{p}(U) \cap \{\{x, y\} ; (x, y) \in [3] \times [4, 6]\} \neq \emptyset$.*

Lemma 5.16. *Let U be a word over $\mathcal{S} \cup \mathcal{C}$. Then $U \in \mathcal{L}(\mathfrak{S}_{1,3} \circ \mathfrak{S}_{4,9} \circ \mathfrak{S}_{10,15} \circ \mathfrak{C}) \setminus \mathcal{L}(\mathfrak{S}_{1,9} \circ \mathfrak{S}_{10,15} \circ \mathfrak{C})$ if and only if $U \in \mathcal{N}$ and U has a 2×4 pattern with $\mathfrak{p}(U) \cap \{\{x, y\} ; (x, y) \in ([3], [4, 6])\} = \emptyset$.*

Theorem 5.17. *Let U be a word over $\mathcal{S} \cup \mathcal{C}$. Then $U \in \mathcal{L}(\mathfrak{S}_{1,3} \circ \mathfrak{S}_{4,9} \circ \mathfrak{S}_{10,15} \circ \mathfrak{C})$ if and only if $U \in \mathcal{N}$.*

Proof. If $|U| = 1$ then the result follows from [Lemma 5.13](#). If $|U| > 1$, then U has a $2 \times 2 \times 2$ or a 2×4 pattern and the result follows from [Proposition 5.12](#) and [Lemmas 5.14](#), [5.15](#) and [5.16](#). \square

6 Lower Bounds

Recall that the *distance* between operators U and V is defined as $\|U - V\| = \sup\{\|Uv - Vv\| ; \|v\| = 1\}$. Because \mathcal{G} is universal, for every $\epsilon > 0$ and every element $U \in \text{SU}(4)$, there exists $V \in \mathcal{G}$ such that $\|U - V\| \leq \epsilon$. In such a case we say that V is an ϵ -approximation of U . We now take advantage of [Theorem 5.17](#) to count Clifford+CS operators and use these results to derive a worst-case lower bound on the CS-count of approximations.

Lemma 6.1. *Let $n \geq 1$. There are $86400(3 \cdot 8^n - 2 \cdot 4^n)$ Clifford+CS operators of CS-count exactly n .*

Proof. Each Clifford+CS operator is represented by a unique normal form and this representation is CS-optimal. Hence, to count the number of Clifford+CS operators of CS-count n , it suffices to count the normal forms of CS-count n . By [Theorem 5.17](#), and since Clifford operators have CS-count 0, a normal form of CS-count n is a word

$$w = w_1 w_2 w_3 w_4 \tag{19}$$

such that $w_1 \in \mathcal{L}(\mathfrak{S}_{1,3})$, $w_2 \in \mathcal{L}(\mathfrak{S}_{4,9})$, $w_3 \in \mathcal{L}(\mathfrak{S}_{10,15})$, $w_4 \in \mathcal{L}(\mathfrak{C})$ and the CS-counts of w_1 , w_2 , and w_3 sum to n . There are

$$(6 \cdot 8^{n-1} + 6 \cdot 4^{n-1} + 3 \cdot 2^{n-1}) \cdot |\mathcal{C}| \tag{20}$$

words of the form of [Equation \(19\)](#) such that exactly one of w_1 , w_2 , or w_3 is not ε . Similarly, there are

$$\left(\sum_{0 < l < n} 18 \cdot 2^{2n-3-l} + \sum_{0 < l < n} 18 \cdot 2^{3n-4-2l} + \sum_{0 < j < n} 36 \cdot 2^{3n-5-j} \right) \cdot |\mathcal{C}| \tag{21}$$

words of the form of [Equation \(19\)](#) such that exactly two of w_1 , w_2 , or w_3 are not ε . Finally, the number of words of the form of [Equation \(19\)](#) such that w_1 , w_2 , and w_3 are not ε is

$$\left(\sum_{0 < l < n-j} \sum_{0 < j < n} 108 \cdot 2^{3n-6-j-2l} \right) \cdot |\mathcal{C}| \tag{22}$$

Summing [Equations \(20\), \(21\) and \(22\)](#) and applying the geometric series formula then yields the desired result. \square

Corollary 6.2. *For $n \in \mathbb{N}$, there are $\frac{46080}{7}(45 \cdot 8^n - 35 \cdot 4^n + 4)$ distinct Clifford+CS operators of CS-count at most n .*

Proof. Recall that the Clifford+CS operators of CS-count 0 are exactly the Clifford operators and that $|\mathcal{C}| = 92160$. The result then follows from [Lemma 6.1](#) and the geometric series formula. \square

Proposition 6.3. *For every $\epsilon \in \mathbb{R}^{>0}$, there exists $U \in \text{SU}(4)$ such that any Clifford+CS ϵ -approximation of U has CS-count at least $5 \log_2(1/\epsilon) - 0.67$.*

Proof. By a volume counting argument. Each operator must occupy an ϵ -ball worth of volume in 15-dimensional $\text{SU}(4)$ space, and the sum of all these volumes must add to the total volume of $\text{SU}(4)$ which is $(\sqrt{2}\pi^9)/3$. The number of circuits up to CS-count n is taken from [Corollary 6.2](#) (we must divide the result by two to account for the absence of overall phase ω in the special unitary group) and a 15-dimensional ϵ -ball has a volume of

$$\frac{\pi^{\frac{15}{2}}}{\Gamma(\frac{15}{2} + 1)} \epsilon^{15}. \quad \square$$

Let U be an element of \mathcal{G} of determinant 1. By [Equation \(1\) of Section 2](#), U can be written as

$$U = \frac{1}{\sqrt{2}^k} M$$

where $k \in \mathbb{N}$ and the entries of M belong to $\mathbb{Z}[i]$. We can therefore talk about the least denominator exponent of the $\text{SU}(4)$ representation of U . We finish this section by relating the least denominator exponent of the $\text{SU}(4)$ representation of U and the CS-count of the normal form of U .

Proposition 6.4. *Let U be an element of \mathcal{G} of determinant 1, let k be the least denominator exponent of the $\text{SU}(4)$ representation of U , and let k' be the CS-count of the normal form of U . Then*

$$\frac{k-3}{2} \leq k' \leq 2k+2.$$

Proof. The CS-count of the normal form of U is equal to the least denominator exponent of the $\text{SO}(6)$ representation of U . [Equation \(11\)](#) then implies the upper bound for k' . Likewise, examination of [Theorem 5.17](#) reveals that the CS operators in the circuit for U must be separated from one another by a Clifford with a least denominator exponent of at most 2 in its unitary representation. Combining this with the fact that the largest least denominator exponent of an operator in \mathcal{C} is 3, we arrive at the lower bound for k' . \square

Remark 6.5. It was established in [\[31\]](#) that, for single-qubit Clifford+ T operators of determinant 1, there is a simple relation between the least denominator exponent of an operator and its T -count: if the least denominator exponent of the operator is k , then its T -count is $2k - 2$ or $2k$. Interestingly, this is not the case for Clifford+CS operators in $\text{SU}(4)$, as suggested by [Proposition 6.4](#). Clearly, the CS-count of an operator always scales linearly with the least denominator exponent of its unitary representation. For large k , computational experiments with our code [\[16\]](#) suggest that most operators are such that $k' \approx k$, though there are examples of operators with $k' \approx 2k$. One example of such an operator is $[R(X \otimes I, I \otimes Z)R(X \otimes I, I \otimes X)R(Z \otimes I, I \otimes X)R(Z \otimes I, I \otimes Z)]^m$ for $m \in \mathbb{N}$.

7 Conclusion

We described an exact synthesis algorithm for a fault-tolerant multi-qubit gate set which is simultaneously optimal, practically efficient, and explicitly characterizes all possible outputs. The algorithm establishes the existence of a unique normal form for two-qubit Clifford+ CS circuits. We showed that the normal form for an operator can be computed with a number of arithmetic operations linear in the gate-count of the output circuit. Finally, we used a volume counting argument to show that, in the typical case, ϵ -approximations of two-qubit unitaries will require a CS -count of at least $5 \log_2(1/\epsilon)$.

We hope that the techniques developed in the present work can be used to obtain optimal multi-qubit normal forms for other two-qubit gate sets, such as the two-qubit Clifford+ T gate set. Indeed, it can be shown that the $SO(6)$ representation of Clifford+ T operators are exactly the set of $SO(6)$ matrices with entries in the ring $\mathbb{Z}[1/\sqrt{2}]$. Further afield, the exceptional isomorphism for $SU(8)$ could potentially be leveraged to design good synthesis algorithms for three-qubit operators. Such algorithms would provide a powerful basis for more general quantum compilers.

An interesting avenue for future research is to investigate whether the techniques and results presented in this paper can be used in the context of *synthillation*. Quantum circuit synthesis and magic state distillation are often kept separate. But it was shown in [7] that performing synthesis and distillation simultaneously (synthillation) can lead to overall savings. The analysis presented in [7] uses T gates and T states. Leveraging higher-dimensional synthesis methods such as the ones presented here, along with distillation of CS states, could yield further savings.

8 Acknowledgements

AG was partially supported by the Princeton Center for Complex Materials, a MRSEC supported by NSF grant DMR 1420541. NJR was partially supported by the Natural Sciences and Engineering Research Council of Canada (NSERC), funding reference number RGPIN-2018-04064.

We would like to thank Matthew Amy, Xiaoning Bian, and Peter Selinger for helpful discussions. In addition, we would like to thank the anonymous reviewers whose comments greatly improved the paper.

9 Contributions

All authors researched, collated, and wrote this paper.

10 Competing Interests

The authors declare no competing interests.

11 Data Availability

The sets of various CS -count operators used to generate the algorithmic performance information in [Table 2](#) are available at [\[16\]](#).

References

- [1] M. Amy, A. N. Glaudell, and N. J. Ross. Number-theoretic characterizations of some restricted Clifford+ T circuits. *Quantum*, 4:252, Apr. 2020. Also available from [arXiv:1908.06076](#).

- [2] M. Amy, D. Maslov, M. Mosca, and M. Roetteler. A meet-in-the-middle algorithm for fast synthesis of depth-optimal quantum circuits. *IEEE Trans. Comput. Aided Des. Integr. Circuits Syst.*, 32(6):818–830, 2013.
- [3] M. Beverland, E. Campbell, M. Howard, and V. Kliuchnikov. Lower bounds on the non-Clifford resources for quantum computations. *Quantum Science and Technology*, 5(3):035009, 2020.
- [4] A. Blass, A. Bocharov, and Y. Gurevich. Optimal ancilla-free Pauli+ V circuits for axial rotations. *Journal of Mathematical Physics*, 56, 12 2014.
- [5] A. Bocharov, X. Cui, V. Kliuchnikov, and Z. Wang. Efficient topological compilation for a weakly integral anyonic model. *Physical Review A*, 93(1):012313, 2016. Also available from [arXiv:1504.03383](https://arxiv.org/abs/1504.03383).
- [6] A. Bocharov, Y. Gurevich, and K. M. Svore. Efficient decomposition of single-qubit gates into V basis circuits. *Physical Review A*, 88:012313, Jul 2013. Also available from [arXiv:1303.1411](https://arxiv.org/abs/1303.1411).
- [7] E. T. Campbell and M. Howard. Unified framework for magic state distillation and multiqubit gate synthesis with reduced resource cost. *Phys. Rev. A*, 95:022316, Feb 2017.
- [8] A. W. Cross et al. Scalable randomised benchmarking of non-Clifford gates. *npj Quantum Information*, 2(1):1–5, 2016.
- [9] S. Forest, D. Gosset, V. Kliuchnikov, and D. McKinnon. Exact synthesis of single-qubit unitaries over Clifford-cyclotomic gate sets. *Journal of Mathematical Physics*, 56(8):082201, 2015. Also available from [arXiv:1501.04944](https://arxiv.org/abs/1501.04944).
- [10] B. Foxen et al. Demonstrating a continuous set of two-qubit gates for near-term quantum algorithms. *Phys. Rev. Lett.*, 125:120504, Sep 2020.
- [11] S. Garion and A. W. Cross. Synthesis of CNOT-dihedral circuits with optimal number of two qubit gates. *Quantum*, 4:369, 2020.
- [12] S. Garion et al. Experimental implementation of non-Clifford interleaved randomized benchmarking with a controlled- S gate, 2020.
- [13] B. Giles and P. Selinger. Exact synthesis of multiqubit Clifford+ T circuits. *Physical Review A*, 87:032332, 2013. Also available from [arXiv:1212.0506](https://arxiv.org/abs/1212.0506).
- [14] B. Giles and P. Selinger. Remarks on Matsumoto and Amano’s normal form for single-qubit Clifford+ T operators. Also available from [arXiv:1312.6584](https://arxiv.org/abs/1312.6584), Dec. 2013.
- [15] A. N. Glaudell, N. J. Ross, and J. M. Taylor. Canonical forms for single-qutrit Clifford+ T operators. *Annals of Physics*, 406:54–70, 8/19/2019 2019. Also available from [arXiv:1803.05047](https://arxiv.org/abs/1803.05047).
- [16] A. N. Glaudell, N. J. Ross, and J. M. Taylor. GaussSynth, 2020. doi:10.5281/zenodo.4549819.
- [17] D. Gosset, V. Kliuchnikov, M. Mosca, and V. Russo. An algorithm for the T -count. *Quantum Info. Comput.*, 14(15–16):1261–1276, Nov. 2014.
- [18] J. Haah and M. B. Hastings. Codes and protocols for distilling T , controlled- S , and Toffoli gates. *Quantum*, 2:71, 2018.
- [19] L. E. Heyfron and E. T. Campbell. An efficient quantum compiler that reduces T count. *Quantum Science and Technology*, 4(1):015004, 2018. Also available from [arXiv:1712.01557](https://arxiv.org/abs/1712.01557).
- [20] V. Kliuchnikov, A. Bocharov, M. Roetteler, and J. Yard. A framework for approximating qubit unitaries. Preprint available from [arXiv:1510.03888](https://arxiv.org/abs/1510.03888), 2015.

- [21] V. Kliuchnikov, A. Bocharov, and K. M. Svore. Asymptotically optimal topological quantum compiling. *Physical review letters*, 112(14):140504, 2014. Also available from [arXiv:1310.4150](#).
- [22] V. Kliuchnikov, D. Maslov, and M. Mosca. Fast and efficient exact synthesis of single-qubit unitaries generated by Clifford and t gates. *Quantum Information & Computation*, 13(7–8):607–630, July 2013. Also available from [arXiv:1206.5236](#).
- [23] V. Kliuchnikov, D. Maslov, and M. Mosca. Practical approximation of single-qubit unitaries by single-qubit quantum Clifford and T circuits. *IEEE Transactions on Computers*, 65(1):161–172, Jan 2016. Also available from [arXiv:1212.6964](#).
- [24] V. Kliuchnikov and J. Yard. A framework for exact synthesis. Preprint available from [arXiv:1504.04350](#), 2015.
- [25] O. D. Matteo and M. Mosca. Parallelizing quantum circuit synthesis. *Quantum Science and Technology*, 1(1):015003, mar 2016.
- [26] G. Meuli, M. Soeken, and G. D. Micheli. SAT-based $\{\text{CNOT}, T\}$ quantum circuit synthesis. In *Proceedings of the 10th International Conference on Reversible Computation*, RC '17, pages 175–188, 2018.
- [27] M. Mosca and P. Mukhopadhyay. A polynomial time and space heuristic algorithm for T -count. Preprint available from [arXiv:2006.12440](#), 2020.
- [28] S. Prakash, A. Jain, B. Kapur, and S. Seth. Normal form for single-qutrit Clifford+ T operators and synthesis of single-qutrit gates. *Physical Review A*, 98:032304, Sep 2018. Also available from [arXiv:1803.03228](#).
- [29] B. W. Reichardt. Quantum universality from magic states distillation applied to CSS codes. *Quantum Information Processing*, 4(3):251–264, 2005. Also available from [arXiv:quant-ph/0411036](#).
- [30] N. J. Ross. Optimal ancilla-free Clifford+ V approximation of z -rotations. *Quantum Information & Computation*, 15(11–12):932–950, 2015. Also available from [arXiv:1409.4355](#).
- [31] N. J. Ross and P. Selinger. Optimal ancilla-free Clifford+ T approximation of z -rotations. *Quantum Information & Computation*, 16(11-12):901–953, 2016. Also available from [arXiv:1403.2975](#).
- [32] P. Selinger. Efficient clifford+ T approximation of single-qubit operators. *Quantum Info. Comput.*, 15(1–2):159–180, Jan. 2015.
- [33] S. Sheldon, E. Magesan, J. M. Chow, and J. M. Gambetta. Procedure for systematically tuning up cross-talk in the cross-resonance gate. *Phys. Rev. A*, 93:060302, Jun 2016.
- [34] V. V. Shende, I. L. Markov, and S. S. Bullock. Minimal universal two-qubit controlled-NOT-based circuits. *Physical Review A*, 69:062321, Jun 2004. Also available from [arXiv:quant-ph/0308033](#).
- [35] M. Sipser. *Introduction to the Theory of Computation*. International Thomson Publishing, 1st edition, 1996.
- [36] T. Tilma and E. Sudarshan. Generalized Euler angle parametrization for $\text{SU}(N)$. *Journal of Physics A: Mathematical and General*, 35(48):10467, 2002. Also available from [arXiv:math-ph/0205016](#).
- [37] Y. Xu et al. High-fidelity, high-scalability two-qubit gate scheme for superconducting qubits. *Phys. Rev. Lett.*, 125:240503, Dec 2020.
- [38] J. Zhang, J. Vala, S. Sastry, and K. B. Whaley. Geometric theory of nonlocal two-qubit operations. *Phys. Rev. A*, 67:042313, Apr 2003.

A Computing SO(6) Representations

Consider the unitary matrix

$$U = \frac{1}{4} \begin{bmatrix} 3+i & -1-i & -2 & 0 \\ 1-i & 3-i & 0 & -2 \\ 2 & 0 & 3-i & 1+i \\ 0 & 2 & -1+i & 3+i \end{bmatrix}.$$

Suppose we want to compute the entry in the third row and fourth column of it's equivalent SO(6) representative \bar{U} , i.e. $\bar{U}_{3,4}$. Note that we have $\det(U) = 1$, so $U \in \text{SU}(4)$ and we do not have to multiply by a phase before mapping U to \bar{U} .

We need

$$B_3 = s_{-,23,14} = \frac{i}{\sqrt{2}}(e_2 \wedge e_3 - e_1 \wedge e_4)$$

$$B_4 = s_{+,24,13} = \frac{1}{\sqrt{2}}(e_2 \wedge e_4 + e_3 \wedge e_1)$$

in order to calculate

$$\bar{U}_{3,4} = \langle B_3, \bar{U}B_4 \rangle.$$

Computing $\bar{U}B_4$ directly, we have

$$\begin{aligned} \bar{U}B_4 &= \frac{1}{\sqrt{2}} [\bar{U}(e_2 \wedge e_4) + \bar{U}(e_3 \wedge e_1)] \\ &= \frac{1}{\sqrt{2}} [(Ue_2) \wedge (Ue_4) + (Ue_3) \wedge (Ue_1)] \\ &= \frac{1}{16\sqrt{2}} \{ [(-1-i)e_1 + (3-i)e_2 + 2e_4] \wedge [-2e_2 + (1+i)e_3 + (3+i)e_4] \\ &\quad + [-2e_1 + (3-i)e_3 + (-1+i)e_4] \wedge [(3+i)e_1 + (1-i)e_2 + 2e_3] \} \\ &= \frac{1}{16\sqrt{2}} [(2+2i)(e_1 \wedge e_2) - 2i(e_1 \wedge e_3) - (2+4i)(e_1 \wedge e_4) + (-6+2i)(e_2 \wedge e_2) \\ &\quad + (4+2i)(e_2 \wedge e_3) + 10(e_2 \wedge e_4) - 4(e_4 \wedge e_2) + (2+2i)(e_4 \wedge e_3) \\ &\quad + (6+2i)(e_4 \wedge e_4) - (6+2i)(e_1 \wedge e_1) + (-2+2i)(e_1 \wedge e_2) - 4(e_1 \wedge e_3) \\ &\quad + 10(e_3 \wedge e_1) + (2-4i)(e_3 \wedge e_2) + (6-2i)(e_3 \wedge e_3) \\ &\quad + (-4+2i)(e_4 \wedge e_1) + 2i(e_4 \wedge e_2) + (-2+2i)(e_4 \wedge e_3)]. \end{aligned}$$

Using the anticommutation of the wedge product, we can simplify this expression as

$$\begin{aligned} \bar{U}B_4 &= \frac{1}{8\sqrt{2}} [2i(e_1 \wedge e_2) + (-7-i)(e_1 \wedge e_3) + (1-3i)(e_1 \wedge e_4) \\ &\quad + (1+3i)(e_2 \wedge e_3) + (7-i)(e_2 \wedge e_4) - 2i(e_3 \wedge e_4)]. \end{aligned}$$

Examining the rule for computing inner products of wedge products, we see that we have

$$\langle e_i \wedge e_j, e_k \wedge e_\ell \rangle = \langle e_i, e_k \rangle \langle e_j, e_\ell \rangle - \langle e_i, e_\ell \rangle \langle e_j, e_k \rangle = \delta_{i,k} \delta_{j,\ell} - \delta_{i,\ell} \delta_{j,k}$$

and using this property along with the linearity of the inner product, we compute

$$\begin{aligned}
\bar{U}_{3,4} &= \langle B_3, \bar{U}B_4 \rangle \\
&= -\frac{i}{16} [2i\langle e_2 \wedge e_3, e_1 \wedge e_2 \rangle - 2i\langle e_1 \wedge e_4, e_1 \wedge e_2 \rangle + (-7-i)\langle e_2 \wedge e_3, e_1 \wedge e_3 \rangle \\
&\quad -(-7-i)\langle e_1 \wedge e_4, e_1 \wedge e_3 \rangle + (1-3i)\langle e_2 \wedge e_3, e_1 \wedge e_4 \rangle - (1-3i)\langle e_1 \wedge e_4, e_1 \wedge e_4 \rangle \\
&\quad + (1+3i)\langle e_2 \wedge e_3, e_2 \wedge e_3 \rangle - (1+3i)\langle e_1 \wedge e_4, e_2 \wedge e_3 \rangle + (7-i)\langle e_2 \wedge e_3, e_2 \wedge e_4 \rangle \\
&\quad - (7-i)\langle e_1 \wedge e_4, e_2 \wedge e_4 \rangle - 2i\langle e_2 \wedge e_3, e_3 \wedge e_4 \rangle + 2i\langle e_1 \wedge e_4, e_3 \wedge e_4 \rangle] \\
&= -\frac{i}{16} [-(1-3i) + (1+3i)] \\
&= \frac{3}{8}.
\end{aligned}$$

Computing the full matrix, we have

$$\bar{U} = \frac{1}{8} \begin{bmatrix} 4 & 0 & 6 & 2 & 2 & -2 \\ 0 & 8 & 0 & 0 & 0 & 0 \\ -6 & 0 & 1 & 3 & 3 & -3 \\ 2 & 0 & -6 & 7 & -1 & 1 \\ 2 & 0 & -3 & -1 & 7 & 1 \\ -2 & 0 & 3 & 1 & 1 & 7 \end{bmatrix}.$$

B Proof of Lemma 2.2

This appendix contains a proof of Lemma 2.2, whose statement we reproduce below for completeness.

Lemma. *Let $C \in \mathcal{C}$ and let P , Q , and L be distinct elements of $\mathcal{P} \setminus \{I\}$. Assume that P , Q , and L are Hermitian and that $PQ = QP$, $PL = LP$, and $QL = -LQ$. Then the following relations hold:*

$$CR(P, Q)C^\dagger = R(CPC^\dagger, CQC^\dagger), \quad (23)$$

$$R(P, Q) = R(Q, P), \quad (24)$$

$$R(P, -PQ) = R(P, Q), \quad (25)$$

$$R(P, -Q) \in R(P, Q)\mathcal{C}, \quad (26)$$

$$R(P, Q)^2 \in \mathcal{C}, \text{ and} \quad (27)$$

$$R(P, L)R(P, Q) = R(P, Q)R(P, iQL). \quad (28)$$

Proof. Since \mathcal{C} is the normalizer of \mathcal{P} , CPC^\dagger and CQC^\dagger are Hermitian and commuting elements of $\mathcal{P} \setminus \{I\}$. Equation (23) then follows Definition 2.1. Equation (24) is a direct consequence of Definition 2.1. Now, we have $R(P, Q) = \exp((i\pi/8)(I - P - Q + PQ))$ and therefore

$$R(P, -PQ) = \exp((i\pi/8)(I - P - (-PQ) + (P)(-PQ))) = \exp((i\pi/8)(I - P - Q + PQ)) = R(P, Q),$$

which proves Equation (25). For Equation (26), first note that

$$\begin{aligned}
R(P, -Q) &= \exp((i\pi/8)(I - P - (-Q) + (P)(-Q))) = R(P, Q) \\
&= \exp((i\pi/8)(I - P - Q + PQ)) \exp((i\pi/4)Q) \exp((i\pi/4)(-PQ)) \\
&= R(P, Q) \exp((i\pi/4)Q) \exp((i\pi/4)(-PQ)).
\end{aligned} \quad (29)$$

We now show that the last two terms in Equation (29) are Clifford operators. For any $A, B \in \mathcal{P}$, either

$AB = BA$ or $AB = -BA$ holds. For Hermitian Pauli A and any $B \in \mathcal{P}$, we compute

$$\begin{aligned} \exp((i\pi/4)A)B \exp((i\pi/4)A)^\dagger &= \frac{I + iA}{\sqrt{2}} B \frac{I - iA}{\sqrt{2}} \\ &= A \frac{AB + BA}{2} + i \frac{AB - BA}{2} \\ &= \begin{cases} B \in \mathcal{P} & \text{if } AB = BA \\ iAB \in \mathcal{P} & \text{if } AB = -BA \end{cases} \end{aligned}$$

By the definition of the Clifford group and as $\exp((i\pi/4)I) = \exp(i\pi/4)I \in \mathcal{C}$, we therefore conclude

$$\exp((i\pi/4)A) \in \mathcal{C} \text{ for all } A \in \mathcal{P} \text{ with } A^\dagger = A. \quad (30)$$

As both Q and $-PQ$ are Hermitian Paulis, we conclude from [Equations \(29\)](#) and [\(30\)](#) that

$$R(P, -Q) = R(P, Q) \exp((i\pi/4)Q) \exp((i\pi/4)(-PQ)) \in R(P, Q)\mathcal{C}$$

which proves [Equation \(26\)](#). From [Equation \(30\)](#) we also conclude [Equation \(27\)](#) holds as $\{I, -P, -Q, PQ\}$ is a set of commuting Hermitian Paulis and thus

$$R(P, Q)^2 = \exp((i\pi/4)I) \exp((i\pi/4)(-P)) \exp((i\pi/4)(-Q)) \exp((i\pi/4)PQ) \in \mathcal{C}.$$

Finally, note that if $R(P, Q)$ is as in [Definition 2.1](#) then $(I - P)/2$ and $(I - Q)/2$ are idempotent. We can therefore explicitly compute the exponential to get

$$R(P, Q) = I + (i - 1) \left(\frac{I - P}{2} \right) \left(\frac{I - Q}{2} \right).$$

Using the above expression, together with the fact that $QL = -LQ$, yields

$$\begin{aligned} R(P, L)R(P, Q) &= I + (i - 1) \left(\frac{I - P}{2} \right) \left[\frac{3+i}{4}I - \frac{1+i}{4}L - \frac{1+i}{4}Q - \frac{1+i}{4}(-iLQ) \right] \\ &= I + (i - 1) \left(\frac{I - P}{2} \right) \left[\frac{3+i}{4}I - \frac{1+i}{4}(-iQiQL) - \frac{1+i}{4}Q - \frac{1+i}{4}(iQL) \right] \\ &= I + (i - 1) \left(\frac{I - P}{2} \right) \left[\frac{3+i}{4}I - \frac{1+i}{4}Q - \frac{1+i}{4}iQL - \frac{1+i}{4}(-iQiQL) \right] \\ &= R(P, Q)R(P, iQL), \end{aligned}$$

which proves [Equation \(28\)](#) and thereby completes our proof. \square

C Proofs of [Lemma 5.15](#) and [Lemma 5.16](#)

In this appendix, we provide proofs for [Lemma 5.15](#) and [Lemma 5.16](#). We first establish a useful lemma for notational purposes.

Lemma C.1. *Let $U = S_1 \cdots S_k$ be a word over $\mathcal{S} \cup \mathcal{C}$ such that $U \in \mathcal{L}(\mathfrak{S}_{1,15})$. Then $\mathfrak{p}(U^{-1}) = \mathfrak{p}(S_k)$.*

Proof. By [Lemma 5.14](#), we know \bar{U} has least denominator exponent k and a $2 \times 2 \times 2$ pattern, and by [Lemmas 4.9](#) and [4.10](#) we observe that \bar{U}^\top must likewise have a $2 \times 2 \times 2$ pattern. Therefore, if V is the normal form equivalent to U^{-1} , we have $V \in \mathcal{L}(\mathfrak{S}_{1,15} \circ \mathfrak{C})$ such that $\bar{V} = \bar{U}^\top$ has least denominator exponent k and $\mathfrak{p}(V) = \mathfrak{p}(U^{-1})$. By [Lemma 2.2](#), we know that $\bar{S}_k^2 \in \bar{\mathcal{C}}$ and so US_k has an $\text{SO}(6)$ representation with least denominator exponent $k - 1$. Then we have that $\bar{S}_k^\top \bar{V}$ has least denominator exponent $k - 1$, and as V has a $2 \times 2 \times 2$ pattern, S_k is the first letter of the normal form V . Thus we conclude that $\mathfrak{p}(S_k) = \mathfrak{p}(V) = \mathfrak{p}(U^{-1})$. \square

We use [Lemma C.1](#) to concisely describe the pattern of the rightmost letter of a word $U \in \mathcal{L}(\mathfrak{S}_{a,b}) \subseteq \mathcal{L}(\mathfrak{S}_{1,15})$ throughout the remaining lemmas.

Lemma. *Let U be a word over $\mathcal{S} \cup \mathcal{C}$. Then $U \in \mathcal{L}(\mathfrak{S}_{1,9} \circ \mathfrak{S}_{10,15} \circ \mathfrak{C}) \setminus \mathcal{L}(\mathfrak{S}_{1,15} \circ \mathfrak{C})$ if and only if $U \in \mathcal{N}$ and U has a 2×4 pattern with $\mathfrak{p}(U) \cap \{\{x, y\} ; (x, y) \in [3] \times [4, 6]\} \neq \emptyset$.*

Proof. By [Lemmas 5.13](#) and [5.14](#), $\mathcal{L}(\mathfrak{S}_{1,15} \circ \mathfrak{C})$ accounts for all Clifford and $2 \times 2 \times 2$ pattern normal forms. Any accepting word of $\mathfrak{S}_{1,9} \circ \mathfrak{S}_{10,15} \circ \mathfrak{C}$ which is not accepted by $\mathfrak{S}_{1,15} \circ \mathfrak{C}$ is of the form $U_1 U_2$ where $U_1 \in \mathcal{L}(\mathfrak{S}_{1,9}) \subsetneq \mathcal{L}(\mathfrak{S}_{1,15})$, $U_2 \in \mathcal{L}(\mathfrak{S}_{10,15} \circ \mathfrak{C}) \subsetneq \mathcal{L}(\mathfrak{S}_{1,15} \circ \mathfrak{C})$, and $\mathfrak{p}(U_1^{-1}) \cap \mathfrak{p}(U_2) \neq \emptyset$. We can then restate our lemma as follows:

$$\begin{aligned} & \{U_1 U_2 ; U_1 \in \mathcal{L}(\mathfrak{S}_{1,9}), U_2 \in \mathcal{L}(\mathfrak{S}_{10,15} \circ \mathfrak{C}), |U_1| = \ell, |U_2| = k - \ell, \text{ and } \mathfrak{p}(U_1^{-1}) \cap \mathfrak{p}(U_2) \neq \emptyset\} \\ & \qquad \qquad \qquad = \\ & \{U \in \mathcal{N} ; |U| = k \text{ and } \mathfrak{p}(U) \text{ is a } 2 \times 4 \text{ pattern s.t. } \mathfrak{p}(U) \cap \{\{x, y\} ; (x, y) \in [3] \times [4, 6]\} \neq \emptyset\} \end{aligned} \tag{31}$$

for all $k \geq 3$ and $1 \leq \ell \leq k - 2$. We begin with a useful result, and afterwards proceed by induction on k .

- Consider a length k accepting word U' of the above form such that $\ell = 1$. Then $U' = S'U$ where, given that the first letter of U is S , we have $S' \in \{\mathcal{S}_1, \dots, \mathcal{S}_9\}$, $S \in \{\mathcal{S}_{10}, \dots, \mathcal{S}_{15}\}$, and $U \in \mathcal{L}(\mathfrak{S}_{10,15} \circ \mathfrak{C})$. We know that we must have $\mathfrak{p}(S') \cap \mathfrak{p}(S) \neq \emptyset$, and by inspection the only way this is achieved is if $\mathfrak{p}(S') \cap \mathfrak{p}(S) = \{\{a, b\}\} \subseteq \{\{x, y\} ; (x, y) \in [3] \times [4, 6]\}$. As $\mathfrak{p}(S'^{-1}) = \mathfrak{p}(S')$ is not finer than $\mathfrak{p}(S) = \mathfrak{p}(U)$, we conclude that the least denominator exponent of \bar{U}' is $k - 1$ by [Lemmas 4.13](#) and [5.14](#). Furthermore, we have

$$\rho_{k-1}(\bar{U}')[\{a, b\} ; [6]] = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

and so U' must have a 2×4 pattern with $\{a, b\} \in \mathfrak{p}(U')$. This means that

$$\mathfrak{p}(U') \cap \{\{x, y\} ; (x, y) \in [3] \times [4, 6]\} \neq \emptyset$$

and $\mathfrak{p}(S')$ is finer than $\mathfrak{p}(U')$. For each such 2×4 pattern, there is one and only one element of $\{\mathcal{S}_1, \dots, \mathcal{S}_9\}$ which is a finer partition than $\mathfrak{p}(U')$, and therefore S' is the leftmost syllable of the normal form equivalent to U' under FFP. As $U \in \mathcal{N}$, we thus conclude that $U' \in \mathcal{N}$.

- We now show that accepting words of the above form with $k = 3$ (enforcing $\ell = 1$) constitute all such length 3 normal forms with the described pattern. Clearly, any $U \in \mathcal{N}$ with the aforementioned pattern must be of the form $U = S_1 S_2 C$ where $S_1 \in \{\mathcal{S}_1, \dots, \mathcal{S}_9\}$, $S_2 \in \mathcal{S}$, and $C \in \mathcal{C}$. Furthermore, we must have $\mathfrak{p}(S_1) \cap \mathfrak{p}(S_2) = \{\{a, b\}\}$ with $\{a, b\} \in \{\{x, y\} ; (x, y) \in [3] \times [4, 6]\}$ to produce the appropriate 2×4 pattern. By inspection, this implies that $S_2 \in \{\mathcal{S}_{10}, \dots, \mathcal{S}_{15}\}$, and so U is accepted by $\mathfrak{S}_{1,9} \circ \mathfrak{S}_{10,15} \circ \mathfrak{C}$ but not $\mathfrak{S}_{1,15} \circ \mathfrak{C}$ as required. Therefore, [Equation \(31\)](#) holds for $k = 3$.
- Now suppose that [Equation \(31\)](#) holds for some $k \geq 3$. We will show that [Equation \(31\)](#) holds for $k + 1$ by establishing two inclusions.

\subseteq : We have already proven this inclusion in the case of $\ell = 1$ for all k . We therefore need only consider the $\ell > 1$ case. Let $U \in \mathcal{L}(\mathfrak{S}_{1,9} \circ \mathfrak{S}_{10,15} \circ \mathfrak{C}) \setminus \mathcal{L}(\mathfrak{S}_{1,15} \circ \mathfrak{C})$ be a word of length k whose first letter is $S \in \{\mathcal{S}_1, \dots, \mathcal{S}_9\}$. Then $U \in \mathcal{N}$ and $\mathfrak{p}(U)$ is a 2×4 pattern such that $\mathfrak{p}(U) \cap \mathfrak{p}(S) = \{\{a, b\}\}$ with $a \in [3]$ and $b \in [4, 6]$. By inspection, we explicitly have $S = \{\{a, b\}, \{c, d\}, \{e, f\}\}$ with $\{a, c, d\} = [3]$ and $\{b, e, f\} = [4, 6]$. Furthermore, the least denominator exponent of \bar{U} is $k - 1$. Suppose that $U' = S'U$ is a word of length $k + 1$ accepted by $\mathfrak{S}_{1,9} \circ \mathfrak{S}_{10,15} \circ \mathfrak{C}$. Then by [Definition 5.10](#) we have $\mathfrak{p}(S') \cap \mathfrak{p}(S) = \emptyset$ with $S' \in \{\mathcal{S}_1, \dots, \mathcal{S}_9\}$. By inspection, this implies that S' must have the pattern $\mathfrak{p}(S') = \{\{a, c\}, \{b, e\}, \{d, f\}\}$ (up to $c \leftrightarrow d$ and $e \leftrightarrow f$) with $\{a, b, c, d, e, f\}$ as defined before. As $\mathfrak{p}(S'^{-1}) = \mathfrak{p}(S')$ is such that $\mathfrak{p}(S'^{-1}) \cap \mathfrak{p}(U) = \emptyset$, then $\mathfrak{p}(S'^{-1})$ is not finer than $\mathfrak{p}(U)$ and so the least denominator exponent of U' must be k . Consider

the set $\{d, f\}$ and let r_d and r_f be the corresponding rows of the residue matrix of \bar{U} . Explicitly, we have

$$\rho_{k-1}(\bar{U})[\{d, f\}; [6]] = \begin{bmatrix} r_d \\ r_f \end{bmatrix}$$

with $r_d = r_f$ as $\{d, f\}$ is a subset of the cardinality four element of $\mathfrak{p}(U)$. Directly calculating the rows of the residue matrix for \bar{U}' yields

$$\rho_k(\bar{U}')[\{d, f\}; [6]] = \begin{bmatrix} r_d + r_f \\ r_d + r_f \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

and we therefore conclude that $\{d, f\} \in \mathfrak{p}(U')$ and U' has a 2×4 pattern. As $\mathfrak{p}(S')$ is the lowest-indexed element of \mathcal{S} finer than $\mathfrak{p}(U')$, under FFP we conclude that S' is the leftmost syllable of the normal form equivalent to U' . Since $U \in \mathcal{N}$ by assumption, we therefore conclude $U' = S'U \in \mathcal{N}$, and have established that U' has a 2×4 pattern such that $\mathfrak{p}(U') \cap \{\{x, y\} ; (x, y) \in [3] \times [4, 6]\} \neq \emptyset$.

⊇: Suppose that U' is a normal form of length $k + 1$ with a 2×4 pattern such that $\mathfrak{p}(U') \cap \{\{x, y\} ; (x, y) \in [3] \times [4, 6]\} \neq \emptyset$. Write U' as $U' = S'U$ for some unknown normal form U . We then have $S' \in \{\mathcal{S}_1, \dots, \mathcal{S}_9\}$ and $\mathfrak{p}(S') = \{\{a, b\}, \{c, d\}, \{e, f\}\}$ such that $\{a, c, d\} = [3]$, $\{b, e, f\} = [4, 6]$, and $\mathfrak{p}(S') \cap \mathfrak{p}(U') = \{\{a, b\}\}$. For $\{i, j\} \in \mathfrak{p}(S')$, let the corresponding rows of the residue matrix of \bar{U} be r_i and r_j . Explicitly, we have

$$\rho_{k-1}(\bar{U})[\{i, j\}; [6]] = \begin{bmatrix} r_i \\ r_j \end{bmatrix}.$$

Direct calculation of the rows for the residue matrix of \bar{U}' yields

$$\rho_k(\bar{U}')[\{i, j\}; [6]] = \begin{bmatrix} r_i + r_j \\ r_i + r_j \end{bmatrix}.$$

As $r_a + r_b = 0$ per the pattern of U' , we conclude that $r_a = r_b$. For the sets $\{c, d\}$ and $\{e, f\}$ the corresponding rows in the residue matrix of \bar{U}' are nonzero, and so by similar reasoning we conclude that $r_c \neq r_d$ and $r_e \neq r_f$. This leaves two possibilities for the pattern of U : either $\mathfrak{p}(U)$ is a $2 \times 2 \times 2$ pattern with $\{a, b\} \in \mathfrak{p}(U)$ and $\mathfrak{p}(U) \neq \mathfrak{p}(S')$, or $\mathfrak{p}(U)$ is a 2×4 pattern such that $\mathfrak{p}(U) = \{\{d, f\}, \{a, b, c, e\}\}$ (up to $c \rightarrow d$ and $e \rightarrow f$), i.e that $\mathfrak{p}(U) \cap \{\{x, y\} ; (x, y) \in [3] \times [4, 6]\} \neq \emptyset$.

1. In the case that U has a $2 \times 2 \times 2$ pattern, then the leftmost letter S of U is such that $\mathfrak{p}(S) = \mathfrak{p}(U) \neq \mathfrak{p}(S')$ and $\mathfrak{p}(S) \cap \mathfrak{p}(S') = \{\{a, b\}\}$ with $\{a, b\} \in \{\{x, y\} ; (x, y) \in [3] \times [4, 6]\}$. By inspection, we see that the only possibility is that $S \in \{\mathcal{S}_{10}, \dots, \mathcal{S}_{15}\}$. Noting that $\mathfrak{p}(S) \cap \mathfrak{p}(S') \neq \emptyset$, and given that $S' = \mathcal{S}_{j'}$ and $S = \mathcal{S}_j$, we conclude that $j \notin \delta_{\mathfrak{S}, 15}(j', S' = \mathcal{S}_{j'})$. As U is accepted by $\mathfrak{S}_{10, 15} \circ \mathfrak{C}$ by [Lemma 5.14](#) and S' is accepted by $\mathfrak{S}_{1, 9}$, we conclude that $U' = S'U \in \mathcal{L}(\mathfrak{S}_{1, 9} \circ \mathfrak{S}_{10, 15} \circ \mathfrak{C}) \setminus \mathcal{L}(\mathfrak{S}_{1, 15} \circ \mathfrak{C})$.
2. In the case that U has a 2×4 pattern with $\mathfrak{p}(U) \cap \{\{x, y\} ; (x, y) \in [3] \times [4, 6]\} \neq \emptyset$, as the length of U is k we assume by the induction hypothesis that $U \in \mathcal{L}(\mathfrak{S}_{1, 9} \circ \mathfrak{S}_{10, 15} \circ \mathfrak{C}) \setminus \mathcal{L}(\mathfrak{S}_{1, 15} \circ \mathfrak{C})$. Under FFP, we note that the leftmost letter S of U must be an element of $\{\mathcal{S}_1, \dots, \mathcal{S}_9\}$ and have the pattern $\mathfrak{p}(S) = \{\{a, c\}, \{b, e\}, \{d, f\}\}$ given that $\mathfrak{p}(U) = \{\{d, f\}, \{a, b, c, e\}\}$ (again, up to $c \rightarrow d$ and $e \rightarrow f$). Thus, $\mathfrak{p}(S') \cap \mathfrak{p}(S) = \emptyset$. Letting $S' = \mathcal{S}_{j'}$ and $S = \mathcal{S}_j$, we conclude that $j \in \delta_{\mathfrak{S}, 9}(j', S' = \mathcal{S}_{j'})$. Because $S = \mathcal{S}_j$ is the first letter of the word U , we know that the initial state of U must be j . Therefore, by the induction hypothesis, $U' = S'U \in \mathcal{L}(\mathfrak{S}_{1, 9} \circ \mathfrak{S}_{10, 15} \circ \mathfrak{C}) \setminus \mathcal{L}(\mathfrak{S}_{1, 15} \circ \mathfrak{C})$.

We have exhausted all cases for U , and so we conclude that the leftward inclusion holds.

We have shown that [Equation \(31\)](#) holds for words of length $k + 1$ if it holds for words of length k . This completes the inductive step. \square

Lemma. *Let U be a word over $\mathcal{S} \cup \mathcal{C}$. Then $U \in \mathcal{L}(\mathfrak{S}_{1,3} \circ \mathfrak{S}_{4,9} \circ \mathfrak{S}_{10,15} \circ \mathfrak{C}) \setminus \mathcal{L}(\mathfrak{S}_{1,9} \circ \mathfrak{S}_{10,15} \circ \mathfrak{C})$ if and only if $U \in \mathcal{N}$ and U has a 2×4 pattern with $\mathfrak{p}(U) \cap \{\{x, y\} ; (x, y) \in [3] \times [4, 6]\} = \emptyset$.*

Proof. By [Lemmas 5.13, 5.14](#) and [5.15](#), $\mathcal{L}(\mathfrak{S}_{1,9} \circ \mathfrak{S}_{10,15} \circ \mathfrak{C})$ accounts for all Cliffords, $2 \times 2 \times 2$ pattern normal forms, and 2×4 pattern normal forms where the pattern contains an element of $\{\{x, y\} ; (x, y) \in [3] \times [4, 6]\}$. Any accepting word of $\mathfrak{S}_{1,3} \circ \mathfrak{S}_{4,9} \circ \mathfrak{S}_{10,15} \circ \mathfrak{C}$ which is not accepted by $\mathfrak{S}_{1,9} \circ \mathfrak{S}_{10,15} \circ \mathfrak{C}$ is of the form $U_1 U_2 U_3$ where $U_1 \in \mathcal{L}(\mathfrak{S}_{1,3}) \subsetneq \mathcal{L}(\mathfrak{S}_{1,15})$, $U_2 \in \mathcal{L}(\mathfrak{S}_{4,9}) \subsetneq \mathcal{L}(\mathfrak{S}_{1,15})$, $U_3 \in \mathcal{L}(\mathfrak{S}_{10,15} \circ \mathfrak{C}) \subsetneq \mathcal{L}(\mathfrak{S}_{1,15} \circ \mathfrak{C})$, and $\mathfrak{p}(U_1^{-1}) \cap \mathfrak{p}(U_2) \neq \emptyset$. We can then restate our lemma as follows:

$$\begin{aligned} & \{U_1 U_2 U_3 ; U_1 \in \mathcal{L}(\mathfrak{S}_{1,3}), U_2 \in \mathcal{L}(\mathfrak{S}_{4,9}), U_3 \in \mathcal{L}(\mathfrak{S}_{10,15} \circ \mathfrak{C}), |U_1| = \ell, \\ & \quad |U_2| = m, |U_3| = k - \ell - m, \text{ and } \mathfrak{p}(U_1^{-1}) \cap \mathfrak{p}(U_2) \neq \emptyset\} \\ & \quad = \\ & \{U \in \mathcal{N} ; |U| = k \text{ and } \mathfrak{p}(U) \text{ is a } 2 \times 4 \text{ pattern s.t. } \mathfrak{p}(U) \cap \{\{x, y\} ; (x, y) \in [3] \times [4, 6]\} = \emptyset\} \end{aligned} \tag{32}$$

for all $k \geq 3$, $1 \leq \ell \leq k - 2$, and $1 \leq m \leq k - 1 - \ell$. We begin with a useful result, and afterwards proceed by induction on k .

- Consider a length k accepting word U' of the above form such that $\ell = 1$. Then $U' = S'U$ where, given that the first letter of U is S , we have $S' \in \{\mathcal{S}_1, \dots, \mathcal{S}_3\}$, $S \in \{\mathcal{S}_4, \dots, \mathcal{S}_9\}$, and $U \in \mathcal{L}(\mathfrak{S}_{4,9} \circ \mathfrak{S}_{10,15} \circ \mathfrak{C})$. We know that we must have $\mathfrak{p}(S') \cap \mathfrak{p}(S) \neq \emptyset$, and by inspection the only way this is achieved is if $\mathfrak{p}(S') \cap \mathfrak{p}(S) = \{\{a, b\}\} \not\subseteq \{\{x, y\} ; (x, y) \in [3] \times [4, 6]\}$. As $\mathfrak{p}(S'^{-1}) = \mathfrak{p}(S')$ is such that $\mathfrak{p}(S'^{-1}) \cap \mathfrak{p}(U) = \emptyset$, we conclude that S'^{-1} is not finer than $\mathfrak{p}(U)$ and that the least denominator exponent of \bar{U}' is $k - 1$ by [Lemmas 4.13](#) and [5.14](#). In the case that U has a $2 \times 2 \times 2$ pattern, we know that $\mathfrak{p}(U) = \mathfrak{p}(S)$ and so $\{a, b\} \in \mathfrak{p}(U)$. In the case that U has a 2×4 pattern, we know that there exists $\{c, d\} \in \{\{x, y\} ; (x, y) \in [3] \times [4, 6]\}$ such that $\mathfrak{p}(S) \cap \mathfrak{p}(U) = \{\{c, d\}\}$. As $\{c, d\} \neq \{a, b\}$ and $\{\{a, b\}, \{c, d\}\} \subseteq \mathfrak{p}(S)$, we know that $\{a, b\}$ is a subset of the cardinality four element of $\mathfrak{p}(U)$. Therefore, in both cases we have

$$\rho_{k-1}(\bar{U}')[\{a, b\} ; [6]] = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

and so U' must have a 2×4 pattern with $\{a, b\} \in \mathfrak{p}(U')$. This means that

$$\mathfrak{p}(U') \cap \{\{x, y\} ; (x, y) \in [3] \times [4, 6]\} = \emptyset$$

and that $\mathfrak{p}(S')$ is finer than $\mathfrak{p}(U')$. For each such 2×4 pattern, there is one and only one element of $\{\mathcal{S}_1, \dots, \mathcal{S}_3\}$ which is a finer partition than $\mathfrak{p}(U')$, and therefore S' is the leftmost syllable of the normal form equivalent to U' under FFP. As $U \in \mathcal{N}$, we thus conclude that $U' \in \mathcal{N}$.

- We now show that accepting words of the above form with $k = 3$ (enforcing $\ell = 1$ and $m = 1$) constitute all such length 3 normal forms with the described pattern. Clearly, any $U \in \mathcal{N}$ with the aforementioned pattern must be of the form $U = S_1 S_2 C$ where $S_1 \in \{\mathcal{S}_1, \dots, \mathcal{S}_3\}$, $S_2 \in \mathcal{S}$, and $C \in \mathcal{C}$. Furthermore, we must have $\mathfrak{p}(S_1) \cap \mathfrak{p}(S_2) = \{\{a, b\}\}$ with $\{a, b\} \notin \{\{x, y\} ; (x, y) \in [3] \times [4, 6]\}$ to produce the appropriate 2×4 pattern. By inspection, this implies that $S_2 \in \{\mathcal{S}_4, \dots, \mathcal{S}_9\}$, and so U is accepted by $\mathfrak{S}_{1,3} \circ \mathfrak{S}_{4,9} \circ \mathfrak{S}_{10,15} \circ \mathfrak{C}$ but not by $\mathfrak{S}_{1,9} \circ \mathfrak{S}_{10,15} \circ \mathfrak{C}$ as required. Therefore [Equation \(32\)](#) holds for $k = 3$.
- Now suppose that [Equation \(32\)](#) holds for some $k \geq 3$. We will show that [Equation \(32\)](#) holds for $k + 1$ by establishing two inclusions.

\subseteq : We have already proven this inclusion in the case of $\ell = 1$ for all k . We therefore need only consider the $\ell > 1$ case. Let $U \in \mathcal{L}(\mathfrak{S}_{1,3} \circ \mathfrak{S}_{4,9} \circ \mathfrak{S}_{10,15} \circ \mathfrak{C}) \setminus \mathcal{L}(\mathfrak{S}_{1,9} \circ \mathfrak{S}_{10,15} \circ \mathfrak{C})$ be a word of length k whose first letter is $S \in \{\mathcal{S}_1, \dots, \mathcal{S}_3\}$. Then $U \in \mathcal{N}$ and $\mathfrak{p}(U)$ is a 2×4 pattern such that $\mathfrak{p}(U) \cap \mathfrak{p}(S) = \{\{a, b\}\}$ with either $a, b \in [3]$ or $a, b \in [4, 6]$. By inspection, we explicitly have $S = \{\{a, b\}, \{c, d\}, \{e, f\}\}$ with either $\{a, b, e\} = [3]$ or $\{a, b, e\} = [4, 6]$ and $\{c, d, f\} =$

$[6] \setminus \{a, b, e\}$. Furthermore, the least denominator exponent of \bar{U} is $k - 1$. Suppose that $U' = S'U$ is a word of length $k + 1$ accepted by $\mathfrak{S}_{1,3} \circ \mathfrak{S}_{4,9} \circ \mathfrak{S}_{10,15} \circ \mathfrak{C}$. Then by [Definition 5.10](#) we have $\mathfrak{p}(S') \cap \mathfrak{p}(S) = \emptyset$ with $S' \in \{\mathcal{S}_1, \dots, \mathcal{S}_3\}$. By inspection, this implies that S' must have the pattern $\mathfrak{p}(S') = \{\{a, e\}, \{b, d\}, \{c, f\}\}$ (up to $a \leftrightarrow b$ and $c \leftrightarrow d$) with $\{a, b, c, d, e, f\}$ as defined before. As $\mathfrak{p}(S'^{-1}) = \mathfrak{p}(S')$ is such that $\mathfrak{p}(S'^{-1}) \cap \mathfrak{p}(U) = \emptyset$, then $\mathfrak{p}(S'^{-1})$ is not finer than $\mathfrak{p}(U)$ and so the least denominator exponent of U' must be k . Consider the set $\{c, f\}$ and let r_c and r_f be the corresponding rows of the residue matrix of \bar{U} . Explicitly, we have

$$\rho_{k-1}(\bar{U})[\{c, f\}; [6]] = \begin{bmatrix} r_c \\ r_f \end{bmatrix}$$

with $r_c = r_f$ as $\{c, f\}$ is a subset of the cardinality four element of $\mathfrak{p}(U)$. Directly calculating the rows of the residue matrix for \bar{U}' yields

$$\rho_k(\bar{U}')[\{d, f\}; [6]] = \begin{bmatrix} r_c + r_f \\ r_c + r_f \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

and we therefore conclude that $\{c, f\} \in \mathfrak{p}(U')$ and U' has a 2×4 pattern. As $\mathfrak{p}(S')$ is the lowest-indexed element of \mathcal{S} finer than $\mathfrak{p}(U')$, under FFP we conclude that S' is the leftmost syllable of the normal form equivalent to U' . Since $U \in \mathcal{N}$ by assumption, we therefore conclude $U' = S'U \in \mathcal{N}$, and have established U' has a 2×4 pattern such that $\mathfrak{p}(U') \cap \{\{x, y\}; (x, y) \in [3] \times [4, 6]\} = \emptyset$.

- ⊇: Suppose that U' is a normal form of length $k + 1$ with a 2×4 pattern such that $\mathfrak{p}(U') \cap \{\{x, y\}; (x, y) \in [3] \times [4, 6]\} = \emptyset$. Write U' as $U' = S'U$ for some unknown normal form U . We then have $S' \in \{\mathcal{S}_1, \dots, \mathcal{S}_3\}$ and $\mathfrak{p}(S') = \{\{a, b\}, \{c, d\}, \{e, f\}\}$ such that either $\{a, b, e\} = [3]$ or $\{a, b, e\} = [4, 6]$, $\{c, d, f\} = [6] \setminus \{a, b, e\}$, $a \equiv c \pmod{3}$, $b \equiv d \pmod{3}$, $e \equiv f \pmod{3}$, and $\mathfrak{p}(S') \cap \mathfrak{p}(U') = \{\{a, b\}\}$. For $\{i, j\} \in \mathfrak{p}(S')$, let the corresponding rows of the residue matrix of \bar{U} be r_i and r_j . Explicitly, we have

$$\rho_{k-1}(\bar{U})[\{i, j\}; [6]] = \begin{bmatrix} r_i \\ r_j \end{bmatrix}.$$

Direct calculation of the rows for the residue matrix of \bar{U}' yields

$$\rho_k(\bar{U}')[\{i, j\}; [6]] = \begin{bmatrix} r_i + r_j \\ r_i + r_j \end{bmatrix}.$$

As $r_a + r_b = 0$ per the pattern of U' , we conclude that $r_a = r_b$. For the sets $\{c, d\}$ and $\{e, f\}$ the corresponding rows in the residue matrix of \bar{U}' are nonzero, and so by similar reasoning we conclude that $r_c \neq r_d$ and $r_e \neq r_f$. This leaves three possibilities for the pattern of U : either $\mathfrak{p}(U)$ is a $2 \times 2 \times 2$ pattern with $\{a, b\} \in \mathfrak{p}(U)$ and $\mathfrak{p}(U) \neq \mathfrak{p}(S')$, or $\mathfrak{p}(U)$ is a 2×4 pattern such that $\mathfrak{p}(U) = \{\{c, e\}, \{a, b, d, f\}\}$ (up to $c \leftrightarrow d$) so that $\mathfrak{p}(U) \cap \{\{x, y\}; (x, y) \in [3] \times [4, 6]\} \neq \emptyset$, or $\mathfrak{p}(U)$ is a 2×4 pattern such that $\mathfrak{p}(U) = \{\{c, f\}, \{a, b, d, e\}\}$ (up to $c \leftrightarrow d$) so that $\mathfrak{p}(U) \cap \{\{x, y\}; (x, y) \in [3] \times [4, 6]\} = \emptyset$

1. In the case that U has a $2 \times 2 \times 2$ pattern, then the leftmost letter S of U is such that $\mathfrak{p}(S) = \mathfrak{p}(U) \neq \mathfrak{p}(S')$ and $\mathfrak{p}(S) \cap \mathfrak{p}(S') = \{\{a, b\}\}$ with $\{a, b\} \notin \{\{x, y\}; (x, y) \in [3] \times [4, 6]\}$. By inspection, we see that the only possibility is that $S \in \{\mathcal{S}_4, \dots, \mathcal{S}_9\}$. Noting that $\mathfrak{p}(S) \cap \mathfrak{p}(S') \neq \emptyset$, and given that $S' = \mathcal{S}_{j'}$ and $S = \mathcal{S}_j$, we conclude that $j \notin \delta_{\mathfrak{S}, 9}(j', S' = \mathcal{S}_{j'})$. As U is accepted by $\mathfrak{S}_{4,9} \circ \mathfrak{S}_{10,15} \circ \mathfrak{C}$ by [Lemmas 5.14](#) and [5.15](#) and S' is accepted by $\mathfrak{S}_{1,3}$, we conclude that $U' = S'U \in \mathcal{L}(\mathfrak{S}_{1,3} \circ \mathfrak{S}_{4,9} \circ \mathfrak{S}_{10,15} \circ \mathfrak{C}) \setminus \mathcal{L}(\mathfrak{S}_{1,9} \circ \mathfrak{S}_{10,15} \circ \mathfrak{C})$.
2. In the case that U has a 2×4 pattern with $\mathfrak{p}(U) \cap \{\{x, y\}; (x, y) \in [3] \times [4, 6]\} \neq \emptyset$, by [Lemma 5.15](#) we know that the leftmost letter S of U is such that $\mathfrak{p}(S) \cap \mathfrak{p}(U) = \{c, e\}$ (up to $c \leftrightarrow d$). We note that $c \not\equiv d \not\equiv e \pmod{3}$, and under FFP must therefore have

$S \in \{\mathcal{S}_4, \dots, \mathcal{S}_9\}$. This implies that we have $\mathbf{p}(S) = \{\{a, b\}, \{c, e\}, \{d, f\}\}$ (up to $c \leftrightarrow d$), and we therefore conclude that $\mathbf{p}(S') \cap \mathbf{p}(S) \neq \emptyset$. Given $S' = \mathcal{S}_{j'}$ and $S = \mathcal{S}_j$, we conclude that $j \notin \delta_{\mathfrak{S},9}(j', S' = \mathcal{S}_{j'})$. As U is accepted by $\mathfrak{S}_{4,9} \circ \mathfrak{S}_{10,15} \circ \mathfrak{C}$ by [Lemma 5.15](#) and S' is accepted by $\mathfrak{S}_{1,3}$, we get $U' = S'U \in \mathcal{L}(\mathfrak{S}_{1,3} \circ \mathfrak{S}_{4,9} \circ \mathfrak{S}_{10,15} \circ \mathfrak{C}) \setminus \mathcal{L}(\mathfrak{S}_{1,9} \circ \mathfrak{S}_{10,15} \circ \mathfrak{C})$.

3. In the case that U has a 2×4 pattern with $\mathbf{p}(U) \cap \{\{x, y\} ; (x, y) \in [3] \times [4, 6]\} = \emptyset$, as the length of U is k we assume by the induction hypothesis that $U \in \mathcal{L}(\mathfrak{S}_{1,3} \circ \mathfrak{S}_{4,9} \circ \mathfrak{S}_{10,15} \circ \mathfrak{C}) \setminus \mathcal{L}(\mathfrak{S}_{1,9} \circ \mathfrak{S}_{10,15} \circ \mathfrak{C})$. Under FFP, we note that the leftmost letter S of U must be an element of $\{\mathcal{S}_1, \dots, \mathcal{S}_3\}$ and have the pattern $\mathbf{p}(S) = \{\{a, e\}, \{b, d\}, \{c, f\}\}$ given that $\mathbf{p}(U) = \{\{c, f\}, \{a, b, d, e\}\}$ (up to $(a, c) \leftrightarrow (b, d)$). Thus, $\mathbf{p}(S') \cap \mathbf{p}(S) = \emptyset$. Letting $S' = \mathcal{S}_{j'}$ and $S = \mathcal{S}_j$, we conclude that $j \in \delta_{\mathfrak{S},3}(j', S' = \mathcal{S}_{j'})$. Because $S = \mathcal{S}_j$ is the first letter of the word U , we know the initial state of U must be j . Therefore, by the induction hypothesis, $U' = S'U \in \mathcal{L}(\mathfrak{S}_{1,3} \circ \mathfrak{S}_{4,9} \circ \mathfrak{S}_{10,15} \circ \mathfrak{C}) \setminus \mathcal{L}(\mathfrak{S}_{1,9} \circ \mathfrak{S}_{10,15} \circ \mathfrak{C})$.

We have exhausted all cases for U , and so we conclude that the leftward inclusion holds.

We have shown that [Equation \(32\)](#) holds for words of length $k + 1$ if it holds for words of length k . This completes the inductive step. \square