



SIXTY-SIXTH ANNUAL  
WILLIAM LOWELL PUTNAM MATHEMATICAL COMPETITION

Saturday, December 3, 2005

Examination A


**Problem A1**

Show that every positive integer is a sum of one or more numbers of the form  $2^r 3^s$ , where  $r$  and  $s$  are nonnegative integers and no summand divides another.  
(For example,  $23 = 9 + 8 + 6$ .)

**Problem A2**

Let  $S = \{(a,b) \mid a = 1, 2, \dots, n, b = 1, 2, 3\}$ . A *rook tour* of  $S$  is a polygonal path made up of line segments connecting points  $p_1, p_2, \dots, p_{3n}$  in sequence such that (i)  $p_i \in S$ , (ii)  $p_i$  and  $p_{i+1}$  are a unit distance apart, for  $1 \leq i < 3n$ , (iii) for each  $p \in S$  there is a unique  $i$  such that  $p_i = p$ . How many rook tours are there that begin at  $(1,1)$  and end at  $(n,1)$ ?

(An example of such a rook tour for  $n = 5$  is depicted below.)



**Problem A3**

Let  $p(z)$  be a polynomial of degree  $n$ , all of whose zeros have absolute value 1 in the complex plane. Put  $g(z) = p(z)/z^{n/2}$ . Show that all zeros of  $g'(z) = 0$  have absolute value 1.



SIXTY-SIXTH ANNUAL  
WILLIAM LOWELL PUTNAM MATHEMATICAL COMPETITION  
Saturday, December 3, 2005

**Problem A4**

Let  $H$  be an  $n \times n$  matrix all of whose entries are  $\pm 1$  and whose rows are mutually orthogonal. Suppose  $H$  has an  $a \times b$  submatrix whose entries are all 1. Show that  $ab \leq n$ .

**Problem A5**

Evaluate  $\int_0^1 \frac{\ln(x+1)}{x^2+1} dx$ .

**Problem A6**

Let  $n$  be given,  $n \geq 4$ , and suppose that  $P_1, P_2, \dots, P_n$  are  $n$  randomly, independently and uniformly, chosen points on a circle. Consider the convex  $n$ -gon whose vertices are the  $P_i$ . What is the probability that at least one of the vertex angles of this polygon is acute?



SIXTY-SIXTH ANNUAL  
WILLIAM LOWELL PUTNAM MATHEMATICAL COMPETITION  
Saturday, December 3, 2005

Examination B

**Problem B1**

Find a nonzero polynomial  $P(x, y)$  such that  $P(\lfloor a \rfloor, \lfloor 2a \rfloor) = 0$  for all real numbers  $a$ .  
(Note:  $\lfloor v \rfloor$  is the greatest integer less than or equal to  $v$ .)

**Problem B2**

Find all positive integers  $n, k_1, \dots, k_n$  such that  $k_1 + \dots + k_n = 5n - 4$  and

$$\frac{1}{k_1} + \dots + \frac{1}{k_n} = 1.$$

**Problem B3**

Find all differentiable functions  $f: (0, \infty) \rightarrow (0, \infty)$  for which there is a positive real number  $a$  such that

$$f' \left( \frac{a}{x} \right) = \frac{x}{f(x)}$$

for all  $x > 0$ .

**Problem B4**

For positive integers  $m$  and  $n$ , let  $f(m, n)$  denote the number of  $n$ -tuples  $(x_1, x_2, \dots, x_n)$  of integers such that  $|x_1| + |x_2| + \dots + |x_n| \leq m$ . Show that  $f(m, n) = f(n, m)$ .



SIXTY-SIXTH ANNUAL  
WILLIAM LOWELL PUTNAM MATHEMATICAL COMPETITION  
Saturday, December 3, 2005

Examination B

**Problem B5**

Let  $P(x_1, \dots, x_n)$  denote a polynomial with real coefficients in the variables  $x_1, \dots, x_n$ , and suppose that

$$(a) \quad \left( \frac{\partial^2}{\partial x_1^2} + \dots + \frac{\partial^2}{\partial x_n^2} \right) P(x_1, \dots, x_n) = 0 \quad (\text{identically})$$

and that

$$(b) \quad x_1^2 + \dots + x_n^2 \text{ divides } P(x_1, \dots, x_n).$$

Show that  $P = 0$  identically.

**Problem B6**

Let  $S_n$  denote the set of all permutations of the numbers  $1, 2, \dots, n$ . For  $\pi \in S_n$ , let  $\sigma(\pi) = 1$  if  $\pi$  is an even permutation and  $\sigma(\pi) = -1$  if  $\pi$  is an odd permutation. Also, let  $v(\pi)$  denote the number of fixed points of  $\pi$ . Show that

$$\sum_{\pi \in S_n} \frac{\sigma(\pi)}{v(\pi) + 1} = (-1)^{n+1} \frac{n}{n+1}.$$