


**MAT 3321, COMPLEX ANALYSIS AND INTEGRAL TRANSFORMS,  
WINTER 2005**

**Answers to Homework 6  
13.3 #4,8,18; 13.4 #4,16**

**Problem 13.3 #4** We will integrate  $f(z) = z^2/(z^4 - 1)$  counterclockwise around the path  $C$  given by  $x^2 + 16y^2 = 4$ . Note that this is not a circle, but an ellipse with  $x$ -intercepts  $\pm 2$  and  $y$ -intercepts  $\pm 1/2$ , as shown on the right. Note that  $z^4 - 1 = (z^2 - 1)(z^2 + 1) = (z + 1)(z - 1)(z + i)(z - i)$ , thus the function  $f(z)$  has 4 singularities at  $1, -1, i$ , and  $-i$ . Of these, only  $1$  and  $-1$  lie inside  $C$ . By independence of path, the desired integral is equal to the sum of the integrals around  $C_1$  and  $C_2$  as shown in the figure. By Cauchy's integral formula, we have



$$\int_{C_1} \frac{z^2}{z^4 - 1} dz = \int_{C_1} \frac{z^2/(z - 1)(z^2 + 1)}{(z + 1)} dz = 2\pi i g(-1) = -\frac{\pi}{2}i,$$

where  $g(z) = z^2/(z - 1)(z^2 + 1)$ . Similarly,

$$\int_{C_2} \frac{z^2}{z^4 - 1} dz = \int_{C_2} \frac{z^2/(z + 1)(z^2 + 1)}{(z - 1)} dz = 2\pi i h(1) = \frac{\pi}{2}i,$$

where  $h(z) = z^2/(z + 1)(z^2 + 1)$ . Finally,


$$\int_C \frac{z^2}{z^4 - 1} dz = \int_{C_1} \frac{z^2}{z^4 - 1} dz + \int_{C_2} \frac{z^2}{z^4 - 1} dz = 0$$

**Problem 13.3 #8** We are integrating  $f(z) = \frac{z^3 \sin z}{3z - 1}$  counterclockwise around the unit circle. The function  $f(z)$  has a unique singularity at  $z = 1/3$ . We use Cauchy's integral formula:

$$\int_C \frac{z^3 \sin z}{3z - 1} dz = \int_C \frac{\frac{1}{3}z^3 \sin z}{z - \frac{1}{3}} dz = 2\pi i g\left(\frac{1}{3}\right) = 2\pi i \frac{1}{81} \sin \frac{1}{3} \approx 0.02538i,$$

where  $g(z) = \frac{1}{3}z^3 \sin z$ .

**Problem 13.3 #18** We are integrating  $f(z) = \frac{\text{Ln}(z + 1)}{z^2 + 1}$  along the path  $C$  which consists of  $|z - i| = 1.4$  (counterclockwise) and  $|z| = 0.2$  (clockwise), as shown on the right. The function  $f(z)$  has three singularities at  $z = \pm i$  and  $z = -1$ ; the latter singularity is because  $\text{Ln}(0)$  is undefined. Further, the function has a discontinuity along the negative  $x$ -axis starting from  $x = -1$ ; this is due to the discontinuity of  $\text{Ln}(z)$ . Fortunately, our paths of integration do not cross this discontinuity, so we can safely ignore it.



By Cauchy's integral formula, we have

$$\int_C e^{\frac{\text{Ln}(z + 1)}{z^2 + 1}} dz = \int_C e^{\frac{\text{Ln}(z + 1)/(z + i)}{z - i}} dz = 2\pi i g(i),$$

where  $g(z) = \text{Ln}(z + 1)/(z + i)$ . Therefore,  $g(i) = \text{Ln}(1 + i)/2i$ . Since  $1 + i = \sqrt{2}e^{\pi/4}$  in polar coordinates, we have

$$\text{Ln}(1 + i) = \frac{1}{2} \ln 2 + \frac{\pi}{4}i,$$

therefore

$$\int_C \frac{\text{Ln}(z + 1)}{z^2 + 1} dz = 2\pi i \frac{1}{2i} \left( \frac{1}{2} \ln 2 + \frac{\pi}{4}i \right) = \frac{\pi}{2} \ln 2 + \frac{\pi^2}{4}i \approx 1.0888 + 2.4674i$$

**Problem 13.4 #4** Using Cauchy's integral formula for derivatives of an analytic function, we have

$$\int_C \frac{z^6}{(2z - 1)^6} dz = \int_C \frac{\frac{1}{2^6}z^6}{(z - \frac{1}{2})^6} dz = \frac{2\pi i}{5!} f^{(v)}\left(\frac{1}{2}\right),$$


where  $f(z) = \frac{1}{2^6}z^6$ . We calculate this function's 5th derivative:

$$\begin{aligned} f(z) &= \frac{1}{64}z^6 & f'''(z) &= \frac{120}{64}z^3 \\ f'(z) &= \frac{6}{64}z^5 & f^{(iv)}(z) &= \frac{360}{64}z^2 \\ f''(z) &= \frac{30}{64}z^4 & f^{(v)}(z) &= \frac{720}{64}z \end{aligned}$$

Therefore,

$$\int_C \frac{z^6}{(2z-1)^6} dz = \frac{2\pi i}{5!} \cdot \frac{720}{64} \cdot \frac{1}{2} = \frac{2 \cdot 720\pi i}{2 \cdot 120 \cdot 64} = \frac{3\pi i}{32} \approx 0.2945i$$

**Problem 13.4 #16** We are integrating  $f(z) = \frac{e^{z^2}}{z(z-2i)^2}$  along the path  $C$  consisting of  $|z-i|=3$  (counterclockwise) and  $|z|=1$  (clockwise), as shown in the figure. Note that the function  $f(z)$  has two singularities, at  $z=0$  and at  $z=2i$ . Of these,  $z=2i$  has winding number 1 (it lies only inside  $C_1$ ), and  $z=0$  has winding number 0 (it lies inside both paths, but since the paths are traversed in opposite directions, their effects cancel each other out). The desired integral is therefore equal to integrating around a small circle centered at  $z=2i$ , counterclockwise.



$$\int_C \frac{e^{z^2}}{z(z-2i)^2} dz = \int_C \frac{e^{z^2}/z}{(z-2i)^2} dz = 2\pi i f'(2i),$$

where  $f(z) = \frac{e^{z^2}}{z}$ , thus

$$f'(z) = \frac{2z e^{z^2} z - e^{z^2}}{z^2} = \left(2 - \frac{1}{z^2}\right) e^{z^2},$$

thus  $f'(2i) = \left(2 - \frac{1}{-4}\right) e^{-4} = \frac{9}{4} e^{-4}$ . therefore

$$\int_C \frac{e^{z^2}}{z(z-2i)^2} dz = 2\pi i \frac{9}{4} e^{-4} \approx 0.2589i$$