


# Stability of spikes in the presence of cross-diffusion



Theodore Kolokolnikov

Joint works with Juncheng Wei, Michael Ward



# A model of cross-diffusion

- Cross-diffusion model of Shigesada, Kawasaki and Teramoto (1979):

$$\begin{cases} u_t = \Delta [(d_1 + \rho_{12}v) u] + u(a_1 - b_1u - c_1v) \\ v_t = \Delta [(d_2 + \rho_{21}u) v] + v(a_2 - b_1u - c_1v) \\ \text{Neumann B.C. on } [a, b] \end{cases} \quad (1)$$

- Kinetics are just the classic Lotka-Volterra model;  $d_1, d_2$  represent self-diffusion
- Cross-diffusion ( $\rho_{12}, \rho_{21}$ ) represent inter-species avoidance: abundance of  $v$  will cause  $u$  to diffuse faster and vice-versa.
- Without cross-diffusion, only constant solution is stable [Kishimoto, 1981].
- A well-studied sub-regime [Ni, Wu, Xu] is [after scaling]:

$$\begin{cases} u_t = \rho (vu)_{xx} + u(a_1 - b_1u - c_1v) \\ v_t = dv_{xx} + v(a_2 - b_1u - c_1v) \end{cases} \quad (2)$$

with the following assumptions:

$$d \ll 1; \quad \rho \gg 1; \quad \text{all other parameters are positive and of } O(1). \quad (3)$$

- Biologically, when  $\rho$  is large,  $v$  acts as an inhibitor on  $u$ , so that  $u$  diffuses quickly in the regions of high concentration of  $v$ . This effect is believed to be responsible for the segregation of the two species.

# Construction of steady state in 1D

- Lou, Ni, Yotsutani, 2004: Constructed a steady state *in the form of a spike* for  $u$ , and in the form of an inverted spike for  $v$ .
- More explicit computations [spike height] by Wu, Xu, 2010.
- Define

$$\tau = uv$$

so that

$$0 = dv_{xx} + a_2v - b_2\tau - c_2v^2; \quad 0 = \rho\tau_{xx} + \tau \left( \frac{a_1}{v} - b_1\frac{\tau}{v^2} - c_1 \right); \quad (4)$$

- In the limit  $\rho \rightarrow \infty$  the shadow system is:

$$0 = dv_{xx} + a_2v - b_2\tau + c_2v^2; \quad (5)$$

$$Lc_1 = \int_0^L \left( \frac{a_1}{v} - b_1\frac{\tau}{v^2} \right). \quad (6)$$

- Asymptotic solution is:

$$v(x) \sim \frac{a_2}{2c_2} \left[ \frac{3}{2} \tanh^2 \left( \frac{x}{2\varepsilon} \right) + \delta \left( 2 - 3 \tanh^2 \left( \frac{x}{2\varepsilon} \right) \right) \right];$$

$$u \sim \frac{\tau_0}{v(x)}$$

where

$$\varepsilon := \sqrt{\frac{2d}{a_2}} \quad \text{[spike width scaling]}$$







$$\delta := (\varepsilon/L)^{2/3} \frac{3}{4} \left( \frac{b_1 \pi}{b_2 2} \right)^{2/3} \left( 4 \frac{a_1}{a_2} - \frac{b_1}{b_2} - 3 \frac{c_1}{c_2} \right)^{-2/3} \quad \text{[spike height scaling]}$$

$$\tau_0 := \frac{3}{16} \frac{a_2^2}{b_2 c_2};$$

- Note that  $v(0) \sim \frac{a_2}{c_2} \delta = O(\varepsilon^{2/3})$ ;  $u(0) \sim O(\varepsilon^{-2/3})$ .


- This construction works as long as

$$\left( 4 \frac{a_1}{a_2} - \frac{b_1}{b_2} - 3 \frac{c_1}{c_2} \right) > 0.$$



# Stability of multi-spikes

- Very intricate stability properties are observed. Four examples:



- (a) Two stable spikes. Parameter values are  $d = 10^{-3}$ ,  $\rho = 200$ ,  $(a_1, b_1, c_1) = (5, 1, 1)$ ,  $(a_2, b_2, c_2) = (5, 1, 5)$  and  $L = 1.5$ ,  $K = 2$ .
- (b) Slow instability: two spikes persist as a transient state until  $t \sim 1.2 \times 10^4$ . Parameter values are the same as (a) except that  $L = 1$ .
- (c) Fast instability of two boundary spikes: Parameter values are the same as (b).
- (d) Fast instability of three spikes (note log time scale): the middle spike disappears at  $t \sim 20$ . The remaining two spikes slowly drift towards a symmetric equilibrium. Parameter values are the same as (b) except  $K = 3$ .

# Principal stability result

Define

$$\rho_{K,\text{small}} := d^{-1/3} L^{8/3} \frac{c_2}{2} \left( \frac{b_1 \pi}{b_2 2} \right)^{-2/3} \frac{a_2^{1/3}}{2^{1/3}} \left( 4 \frac{a_1}{a_2} - \frac{b_1}{b_2} - 3 \frac{c_1}{c_2} \right)^{5/3}; \quad (7)$$

$$\rho_b := 0.747 \rho_{K,\text{small}}; \quad (8)$$

$$\rho_{K,\text{large}} := \rho_{K,\text{small}} \frac{2 \times 0.747}{1 - \cos[\pi(1 - 1/K)]}. \quad (9)$$

Then:

- A single boundary spike is stable for all  $\rho$  (not exponentially large in  $\varepsilon$ ).
- A double-boundary steady state is stable if  $\rho < \rho_b$  and is unstable otherwise. The instability is due to a large eigenvalue.
- A  $K$ -interior spike steady state with  $K \geq 2$  is stable if  $\rho < \min(\rho_{K,\text{small}}, \rho_{K,\text{large}})$  and is unstable otherwise. When  $K = 1$ , it is stable provided that  $\rho$  is not exponentially large in  $\varepsilon$ .
- The critical scaling is

$$\rho = O(d^{-1/3}) = O(\varepsilon^{-2/3}) \gg 1.$$

# Stability: small vs. large eigenvalues

- $K$  spikes are always stable whenever  $1 \ll \rho \ll d^{-1/3}$  and unstable when  $K \geq 2$  and  $\rho \gg d^{-1/3}$ .

- Recall that  $\rho_{K,\text{large}} := \rho_{K,\text{small}} \frac{2 \times 0.747}{1 - \cos[\pi(1 - 1/K)]}$  and

$$\frac{2 \times 0.747}{1 - \cos[\pi(1 - 1/K)]} = \begin{cases} 1.494 > 1, & K = 2 \\ 0.996 < 1, & K = 3 \\ 0.875 < 1, & K = 4 \end{cases}$$

- $\rho_{K,\text{large}} > \rho_{K,\text{small}}$  if  $K = 2$  but  $\rho_{K,\text{large}} < \rho_{K,\text{small}}$  if  $K \geq 3$ . It follows that the **primary instability is due to small eigenvalues if  $K = 2$  but is due to large eigenvalues if  $K \geq 3$** . This is in agreement with numerical simulations.



# Linearized problem

- Linearized equations are

$$\lambda\phi = d\phi_{xx} + a_2\phi - b_2\psi - c_22v\phi;$$

$$\lambda\left(\frac{1}{v}\psi - \frac{\tau}{v^2}\phi\right) = \rho\psi_{xx} + \left(\frac{a_1}{v} - b_12\frac{\tau}{v^2} - c_1\right)\psi + \left(-\frac{a_1\tau}{v^2} + 2b_1\frac{\tau^2}{v^3}\right)\phi.$$

- Possible boundary conditions:

Config type	Boundary conditions for $\phi$
Single interior spike on $[-L, L]$ even eigenvalue	$\phi'(0) = 0 = \phi'(L)$
Single interior spike on $[-L, L]$ odd eigenvalue	$\phi(0) = 0 = \phi(L)$
Two half-spikes at $[0, L]$	$\phi'(0) = 0 = \phi(L)$
$K$ spikes on $[-L, (2K - 1)L]$ , Periodic BC	$\phi(L) = z\phi(-L), \quad \phi'(L) = z\phi'(-L),$ $z = \exp(2\pi ik/K), \quad k = 0 \dots K - 1$
$K$ spikes on $[-L, (2K - 1)L]$ , Neumann BC	$\phi(L) = z\phi(-L), \quad \phi'(L) = z\phi'(-L),$ $z = \exp(\pi ik/K), \quad k = 0 \dots K - 1$

(same BC for  $\psi$ )

# Reduced problem, large eigenvalues

- Using asymptotic matching, eventually we get a new **point-weight eigenvalue problem (PWE)**:

$$\begin{cases} \lambda\Phi = \Phi_{yy} - \Phi + 2w\Phi - \chi\Phi(0) \\ \Phi \text{ is even and is bounded as } |y| \rightarrow \infty \end{cases} \quad (\text{PWE})$$

where  $w(y) = \frac{3}{2} \operatorname{sech}^2\left(\frac{y}{2}\right)$  satisfies

$$w_{yy} - w + w^2 = 0; \quad w \rightarrow 0 \text{ as } |y| \rightarrow \infty, \quad w'(0) = 0.$$

- For double-boundary spike,

$$\chi = \chi_b := \frac{\varepsilon^{-2/3}}{4\rho} \left( 4\frac{a_1}{a_2} - \frac{b_1}{b_2} - 3\frac{c_1}{c_2} \right)^{5/3} c_2 \left( \frac{b_1 \pi}{b_2 2} \right)^{-2/3} L^{8/3}.$$

- For  $K$  spikes, Neumann BC, there are  $K$  choices for  $\chi$ , namely

$$\chi = \frac{2}{1 - \cos \frac{\pi k}{K}} \chi_b, \quad k = 0 \dots K - 1 \quad \text{and} \quad \chi = \text{very large positive.}$$

# Analysis of *PWEP* $\lambda\Phi = \Phi_{yy} - \Phi + 2w\Phi - \chi\Phi(0)$

- $\lambda = 0$ ,  $\Phi = w_y$  is a solution [corresponds to translation invariance]
- If  $\chi = 0$  then there is an unstable eigenvalue  $\lambda_1 > 0$  and another eigenvalue  $\lambda_3 < 0$ .
- Decompose:

$$\Phi(y) = \Phi^* + \Phi_0(y); \quad \text{where} \quad \Phi^* = \lim_{y \rightarrow \pm\infty} \Phi(y).$$

Then

$$\lambda\Phi^* = -\Phi^* - \chi(\Phi_0(0) + \Phi^*)$$

and  $\Phi_0$  satisfies

$$\lambda\Phi_0 = \Phi_{0yy} - \Phi_0 + 2w\Phi_0 + 2w\Phi^*$$

so the PWEP becomes

$$\lambda\Phi_0 = \Phi_{0yy} - \Phi_0 + 2w\Phi_0 - \frac{2\chi}{\chi + \lambda + 1}\Phi_0(0)w \quad (10)$$

- Ansatz: if  $\Phi_0 = w$ ,  $\lambda = 0$  then  $\chi = \frac{1}{2}$ .
- Rigorous result: there is an unstable eigenvalue  $\lambda > 0$  for all  $\chi < \frac{1}{2}$
- In the limit  $\chi \rightarrow \infty$ , the limiting problem is

$$\lambda\Phi_0 = \Phi_{0yy} - \Phi_0 + 2w\Phi_0 - 2\Phi_0(0)w \quad (11)$$

# Numerics: Hypergeometric reduction


Theorem: the eigenvalues of  $\lambda\Phi = \Phi_{yy} - \Phi + 2w\Phi - \chi\Phi(0)$  are given implicitly by:

$$\lambda = -1 - \chi + 2\chi\Phi_0(0)$$

where

$$\Phi_0(0) = \frac{6\pi\lambda(\lambda+1)}{\sin(\pi\alpha)(4\lambda-5)(4\lambda+3)} - \frac{31}{2\lambda} {}_3F_2\left(\begin{matrix} 1, 3, -1/2 \\ 2+\alpha, 2-\alpha \end{matrix}; 1\right); \quad \alpha = \sqrt{1+\lambda}$$

- Numerical result: all  $\lambda < 0$  whenever  $\chi > 0.669$ ; **stabilization is via a hopf bifurcation.**



# Small eigenvalues

- Construct asymmetric spike steady states
- These bifurcate from the symmetric branch
- The instability thresholds for the small eigenvalues correspond precisely to this bifurcation point!
- **Main result:** For 2 spikes, small eigenvalues is the dominant instability. For 3 or more, large eigenvalues dominate.

# Radial equilibrium in two dimensions

Consider  $\Omega \in \mathbb{R}^2$ . Let  $w$  be the ground state in 2D:

$$\Delta w - w + w^2 = 0; \quad w \rightarrow 0 \text{ as } |y| \rightarrow \infty, \quad \max w = w(0)$$

and define

$$m := \max w(y) = w(0) \approx 2.39195.$$

Suppose that

$$\frac{a_1}{a_2} (2m - 1) - (m - 1) \frac{b_1}{b_2} - m \frac{c_1}{c_2} > 0 \quad (12)$$

and consider the asymptotic limit

$$d \ll 1; \quad \rho \gg 1. \quad (13)$$

If  $\Omega$  is radially symmetric, there is a steady state at  $x = 0$ , in the form of an inverted spike for  $v$ . More precisely, we have

$$v(x) \sim \frac{1}{2m - 1} \frac{a_2}{c_2} (1 - 2\delta) \left( m - w \left( \frac{1 - \delta}{\varepsilon} x \right) + (2m - 1) \delta \right);$$
$$u \sim \frac{\tau_0}{v(x)}$$

where

$$\varepsilon := \sqrt{\frac{(2m-1)d}{a_2}}; \quad \delta \sim \frac{\varepsilon^2}{|\Omega|} \frac{4\pi b_1 m}{b_2 (2m-1) \left( \frac{a_1}{a_2} (2m-1) - (m-1) \frac{b_1}{b_2} - m \frac{c_1}{c_2} \right)} \frac{1}{\left( \frac{a_1}{a_2} (2m-1) - (m-1) \frac{b_1}{b_2} - m \frac{c_1}{c_2} \right)};$$



$$\tau_0 := \frac{(m-1)m a_2^2}{(2m-1)^2 b_2 c_2}.$$

In particular,

$$v(0) \sim \frac{a_2}{c_2} \delta = O(d); \quad u(0) \sim \frac{(m-1)m a_2}{(2m-1)^2 b_2} \frac{1}{\delta} = O\left(\frac{1}{d}\right). \quad (14)$$


# Nice patterns: $\rho = O(1)$

- Spike insertion, spatio-temporal chaos



Sensitivity to initial conditions. The left and right figure differ only in the initial conditions. On the left, symmetric initial conditions result in an intricate a time-periodic solution. On the right, the initial condition is the same as on the left, except for a shift of 0.1 units to the right. dynamics eventually settle to a 5-spike stable pattern. Parameter values for both figures are  $\rho = 7, d_2 = 0.0005, (a_1, b_1, c_1) = (5, 1, 1); (a_2, b_2, c_2) = (1, 1, 2)$ .





$$\rho = 50, \quad (a_1, b_1, c_1) = (5, 1, 1), \quad (a_2, b_2, c_2) = (5, 1, 5)$$

Row 1:  $\rho = 2$ . Spot splits into three spots. Row 2:  $\rho = 4$ . Initially, spot splits into two, final steady state consists of two boundary and one center spot. Row 3:  $\rho = 6$ . Row 4:  $\rho = 500$ . The interior spike is unstable and slowly drifts to the boundary. Once it reaches the boundary, it starts to oscillate indefinitely.

# UCLA Model of hot-spots in crime

- Recently proposed by Short Brantingham, Bertozzi et.al [2008].
- Very "sexy" math: e.g. *The New York Times*, Dec 2010
- Crime is ubiquitous but not uniformly distributed
  - some neighbourhoods are worse than others, leading to crime "hot spots"
  - Crime hotspots can persist for long time.



Fig. 1. Dynamic changes in residential burglary hotspots for two consecutive three-month periods beginning June 2001 in Long Beach, CA. These density maps were created using ArcGIS.

Figure taken from Short et.al., *A statistical model of criminal behaviour*, 2008.

- Crime is temporally correlated:
  - Criminals often return to the spot of previous crime
  - If a home was broken into in the past, the likelihood of subsequent breakin increases
  - Example: graffitti "tagging"

- Two-component model

$$A_t = \varepsilon^2 A_{xx} - A + \rho A + A_0$$

$$\tau \rho_t = D \left( \rho_x - 2 \frac{\rho}{A} A_x \right)_x - \rho A + \bar{A} - A_0.$$

- $\rho(x, t) \equiv$  density of criminals;
- $A(x, t) \equiv$  "attractiveness" of area to crime
- $A_0 = O(1) \equiv$  "baseline attractiveness"
- $D(-2\frac{\rho}{A}A_x)_x$  models the motion of criminals towards higher attractiveness areas
- $\bar{A} - A_0 > 0$  is the baseline criminal "feed rate"
- We assume here:

$$\varepsilon^2 \ll 1, \quad D \gg 1.$$

# Hot-spot steady state

$$0 = \varepsilon^2 A_{xx} - A + \rho A + A_0; \quad 0 = D \left( \rho_x - 2 \frac{\rho}{A} A_x \right)_x - \rho A + \bar{A} - A_0$$

- Key trick:  $\rho_x - 2 \frac{\rho}{A} A_x = A^2 (\rho A^{-2})_x$ . This suggests the change of variables:

$$v = \frac{\rho}{A^2};$$

so that

$$0 = \varepsilon^2 A_{xx} - A + v A^3 + A_0; \quad 0 = D (A^2 v_x)_x - v A^3 + \bar{A} - A_0.$$

- “Shadow limit” Large  $D$  :  $v(x) \sim v_0$ ;

$$\varepsilon^2 A_{xx} - A + v A^3 + A_0 = 0; \quad v_0 \int_0^L A^3 dx = (\bar{A} - A_0) L.$$

- Ansatz:  $v_0 \ll 1$ ,  $A \sim v_0^{-1/2} w(y)$ ,  $y = x/\varepsilon$  where  $w$  is the ground state,  
 $w_{yy} - w + w^3 = 0$ ,  $w'(0) = 0$ ,  $w \rightarrow 0$  as  $|y| \rightarrow \infty$ ;

then

$$v_0 \sim \frac{\left(\int_{-\infty}^{\infty} w^3 dy\right)^2}{4L^2 (\bar{A} - A_0)^2} \varepsilon^2;$$

$$A(x) \sim \begin{cases} \frac{2L(\bar{A} - A_0)}{\varepsilon \int w^3} w(x/\varepsilon), & x = O(\varepsilon) \\ A_0, & x \gg O(\varepsilon). \end{cases}$$

# Main stability result (1D)

- **Main result:** Consider  $K$  spikes on the domain of size  $2KL$ . Then small eigenvalues become unstable if  $D > D_{c,\text{small}}$ ; large eigenvalues become unstable if  $D > D_{c,\text{small}}$  where

$$D_{c,\text{small}} \sim \frac{L^4 (\bar{A} - A_0)^3}{\varepsilon^2 A_0^2 \pi^2}$$
$$D_{c,\text{large}} \sim D_{c,\text{small}} \left( \frac{2}{1 - \cos \frac{\pi}{K}} \right) > D_{c,\text{small}}$$

- Small eigenvalues become unstable before the large eigenvalues.
- **Example:** Take  $L = 1, \bar{A} = 2, A_0 = 1, K = 2, \varepsilon = 0.07$ . Then  $D_{c,\text{small}} = 20.67, D_{c,\text{large}} = 41.33$ .
  - if  $D = 15 \implies$  two spikes are stable
  - if  $D = 30 \implies$  two spikes have very slow developing instability
  - if  $D = 50 \implies$  two spikes have very fast developing instability

# Stability: large eigenvalues

- **Step 1:** Reduces to the nonlocal eigenvalue problem (NLEP):

$$\lambda\phi = \phi'' - \phi + 3w^2\phi - \chi \left( \int w^2\phi \right) w^3 \quad \text{where } w'' - w + w^3 = 0. \quad (15)$$

with

$$\chi \sim \frac{3}{\int_{-\infty}^{\infty} w^3 dy} \left( 1 + \varepsilon^2 D \left( 1 - \cos \frac{\pi k}{K} \right) \frac{A_0^2 \pi^2}{4L^4 (\bar{A} - A_0)^3} \right)^{-1}$$

- **Step 2: *Key identity*:**  $L_0 w^2 = 3w^2$ , where  $L_0\phi := \phi'' - \phi + 3w^2\phi$ . Multiply (15) by  $w^2$  and integrate to get

$$\lambda = 3 - \chi \int w^5 = 3 - \chi \frac{3}{2} \int w^3$$

Conclusion: (15) is stable iff  $\chi > \frac{2}{\int w^3} \iff D > D_{c,\text{large}}$ .

- This NLEP in 1D can be fully solved!!

# Stability: small eigenvalues

- Compute asymmetric spikes
- They bifurcate from symmetric branch
- The bifurcation point is precisely when  $D = D_{c,\text{small}}$ .
- This is “cheating”... but it gets the correct threshold!!



# Two dimensions

$$\left\{ \begin{array}{l} A_t = \varepsilon^2 \Delta A - A + \hat{v} A^3 + A_0 \\ \tau(A\hat{v})_t = D \nabla \cdot (A^2 \nabla \hat{v}) - \hat{v} A^3 + \bar{A} - A_0, \quad x \in \Omega \\ \text{Neumann BC} \end{array} \right.$$

- **Steady-state:** construction is similar to 1D
- **Stability:** of  $K$  hot-spots:
  - If  $K = 1$ , then a single hot-spot is stable with respect to large eigenvalues, as long as  $D$  is not exponentially large in  $1/\varepsilon$ .
  - If  $K \geq 2$ , then the steady state is stable with respect to large eigenvalues if

$$D < \frac{1}{\varepsilon^4} \ln \frac{1}{\varepsilon} \frac{(\bar{A} - A_0)^3 |\Omega|^3 A_0^{-2}}{4\pi K^3 \left( \int_{\mathbb{R}^2} w^3 dy \right)^2}; \quad (16)$$

and it is unstable otherwise.

- Instability thresholds occur when  $D = O\left(\frac{\ln \varepsilon^{-1}}{\varepsilon^4 K^3}\right) \gg 1$ .

# General remarks

- In both models, the instability thresholds occur close to the "shadow limit", i.e. the cross-diffusion term is very large.
- Steady-state computation is essentially a shadow system, but stability computations require more.
- Consider a general **reaction-diffusion system**

$$\begin{cases} u_t = \varepsilon^2 u_{xx} + f(u, w), & \tau w_t = Dw_{xx} + g(u, w) \\ \text{Neumann B.C. on } [a, b] \end{cases}; \quad (17)$$

in the singular limit  $\varepsilon \ll 1$ .

- If we formally take an additional limit  $D \rightarrow \infty$ , we get a **Shadow-limit PDE with an integral constraint**:

$$u_t = \varepsilon^2 u_{xx} + f(u, w_0); \quad \tau \frac{d}{dt} w_0(t) = \frac{1}{b-a} \int_a^b g(u, w_0) dx \quad (18)$$

- The PDE (18) is simpler than (17), but can preserve some of its properties:
- Equilibrium of (18) is similar to (17) with  $D \gg 1$ .
- Stability can be **dramatically different**:
  - Any non-monotone solution of (18) is unstable [Ni, Poláčik, Yanagida, 2001].
  - Can have multiple non-monotone stable solutions of (17), depending on  $D$  [e.g. stable spikes in GM system or multiple stable layers in FitzHugh-Nagumo model]

# Oscillatory layers near the shadow limit

- FitzHuhg-Nagumo type model:

$$u_t = \varepsilon^2 u_{xx} + 2(u - u^3) + w, \quad \tau w_t = Dw_{xx} - u + \beta$$

*Neumann BC on  $[0, 1]$*

$$\varepsilon \ll 1, \quad D \gg 1$$

- Stationary steady state is an interface computed from the shadow limit

$$w \sim 0; \quad u \sim \tanh\left(\frac{l_0 - x}{\varepsilon}\right); \quad l_0 := (1 + \beta)/2$$

- As  $\tau$  is increased, the interface is destabilized via a Hopf Bifurcation. The critical scaling is:

$$\tau = \frac{D}{\varepsilon} \tau_0, \quad \text{where } \tau_0 = O(1).$$

- The interface position is given by





$$l(t) \sim l_0 + A(t) \cos(\sqrt{3/\tau_0 \varepsilon} D^{-1/2} t + \phi_0)$$

where  $A$  is the oscillation envelope that satisfies

$$\frac{D dA}{\varepsilon dt} = \left( \frac{1}{4}(1 - 3\beta^2) - \frac{1}{8\tau_0} \right) A - \frac{3}{4} A^3.$$

- Hopf bifurcation occurs when

$$\tau_{0h} = \begin{cases} \frac{1}{2(1-3\beta^2)} & \text{if } |\beta| < 3^{-1/2}; \\ \infty & \text{otherwise} \end{cases}.$$



# Concluding remarks

- Cross-diffusion (directed movement) can create **stable multi-spike solutions** even in the absence of spatial heterogeneity.
- Stability thresholds for both SKT model and crime model appear very close to the shadow regime
- Stability analysis leads to novel, interesting eigenvalue problems
- The papers can be downloaded from my website,  
`www.mathstat.dal.ca/~tkolokol`

Thank you!